

<p align="center">Table of Contents</p> <p align="center">Section II – County Soil and Site Information Reports</p>

	Issue Date	Date of Last Review	Responsible Staff
<i>Use and Explanation of Soil Interpretations</i>			
<i>Explanation of Key Phrases Used in Soil Interpretations</i>			
Soils Legends	4/02	4/02	SOI
<i>*Acreage and Proportionate Extent of the Soils</i>			
Soil Descriptions - Nontechnical	4/02	4/02	SOI
<i>Use and Explanation of Nontechnical Descriptions</i>			
<i>*Nontechnical Soils Description Report</i>			
Soil Descriptions - Technical	4/02	4/02	SOI
<i>*Map Unit Description Report</i>			
Cropland Interpretations - Technical	4/02	4/02	SOI
<i>*Land Capability and Yields Per Acre of Crops</i>			
<i>*Prime Farmland Report</i>			
<i>*Soil Rating for Plant Growth Index</i>			
<i>*Soil Properties for Conservation Planning</i>			
Rangeland, Grazed Forestland, Native Pastureland Interpretations	4/02	4/02	SOI
<i>*Rangeland Productivity Report</i>			
<i>*Range Site Descriptions</i>			
Forestland Interpretations	1/93	1/93	SOI
<i>Use and Explanation of Forestland Interpretations</i>			
<i>*Woodland Management and Productivity</i>			
Nonagricultural Interpretations	4/02	4/02	SOI
<i>*Building Site Development Report</i>			
<i>*Construction Materials Report</i>			
Recreation Interpretations	4/02	4/02	SOI
<i>*Recreational Interpretations</i>			
Wildlife Interpretations	4/02	4/02	SOI
<i>*Wildlife Interpretations Report</i>			

Pastureland and Hayland Interpretations	4/02	4/02	SOI
<i>*Yields Per Acre of Pasture and Hayland</i>			
	Issue Date	Date of Last Review	Responsible Staff
<hr/>			
Mined Land Interpretations			
<i>Use and Explanation of Mined Land Interpretations</i>			
Windbreak Interpretations	4/02	4/02	SOI
<i>*Conservation Tree and Shrub Management Report</i>			
Engineering Interpretations	4/02	4/02	SOI
<i>*Engineering Index Properties</i>			
<i>*Physical Properties of the Soils</i>			
<i>*Chemical Properties of the Soils</i>			
<i>*Water Features</i>			
<i>*Soil Features</i>			
<i>*Water Management Report</i>			
Waste Disposal Interpretations	4/02	4/02	SOI
<i>*Sanitary Facilities Report</i>			
<i>*Agricultural Waste Management Report</i>			
Hydric Soil Interpretations	4/02	4/02	SOI
<i>Use and Explanation of Hydric Soil Interpretations</i>			
<i>*Hydric Soils List</i>			
HEL Interpretations	See Report		SOI
<i>Use and Explanation of Highly Erodible Land Interpretations</i>			
<i>*Highly Erodible Lands Report</i>			
1 <i>Highly Erodible Land</i>			
2 <i>Potentially Highly Erodible Land</i>			
3 <i>Nonhighly Erodible Land</i>			

**County specific computer generated reports.*

ACREAGE AND PROPORTIONATE EXTENT OF THE SOILS

Boyd County, Nebraska: Published

Map symbol	Soil name	Acres	Percent
Ab	Albaton Silty Clay, 0 To 2 Percent Slopes-----	660	0.2
AnC	Anselmo Fine Sandy Loam, 2 To 6 Percent Slopes-----	5,770	1.7
AnD	Anselmo Fine Sandy Loam, 6 To 11 Percent Slopes-----	7,280	2.1
AnF	Anselmo Fine Sandy Loam, 11 To 20 Percent Slopes-----	5,720	1.6
ArF	Anselmo-Rock Outcrop Complex, 11 To 20 Percent Slopes-----	860	0.2
Ba	Barney Silt Loam, 0 To 2 Percent Slopes-----	2,020	0.6
Bd	Blake Silty Clay Loam, 0 To 2 Percent Slopes-----	400	0.1
Be	Blendon Fine Sandy Loam, 0 To 2 Percent Slopes-----	530	0.2
BeC	Blendon Fine Sandy Loam, 2 To 6 Percent Slopes-----	1,110	0.3
BoD	Boyd Silty Clay, 6 To 11 Percent Slopes-----	3,240	0.9
BrG	Bristow Silty Clay, 20 To 40 Percent Slopes-----	15,650	4.5
Bs	Brocksburg Fine Sandy Loam, 0 To 2 Percent Slopes-----	1,580	0.5
Bt	Brocksburg Loam, 0 To 2 Percent Slopes-----	2,680	0.8
Cb	Cass Fine Sandy Loam, 0 To 2 Percent Slopes-----	3,080	0.9
CrE2	Crofton Silt Loam, 11 To 15 Percent Slopes, Eroded-----	2,760	0.8
DuB	Dunday Loamy Fine Sand, 0 To 3 Percent Slopes-----	2,200	0.6
DuC	Dunday Loamy Fine Sand, 3 To 6 Percent Slopes-----	5,010	1.4
DuD	Dunday Loamy Fine Sand, 6 To 11 Percent Slopes-----	3,460	1.0
DxB	Dunday Loamy Fine Sand, Loamy Substratum, 0 To 3 Percent Slopes-----	1,320	0.4
Et	Eltree Silt Loam, 0 To 2 Percent Slopes-----	2,100	0.6
Go	Grigston Silt Loam, 0 To 2 Percent Slopes-----	4,310	1.2
GrB	Grigston Silt Loam, Channeled, 0 To 3 Percent Slopes-----	2,460	0.7
Ha	Hall Silt Loam, 0 To 2 Percent Slopes-----	2,390	0.7
He	Haynie Silt Loam, 0 To 2 Percent Slopes-----	1,110	0.3
IfD	Inavale Fine Sand, 3 To 11 Percent Slopes-----	840	0.2
IgB	Inavale Fine Sand, Channeled, 0 To 3 Percent Slopes-----	2,350	0.7
IhB	Inavale Loamy Fine Sand, 0 To 3 Percent Slopes-----	4,470	1.3
In	Inavale Fine Sandy Loam, 0 To 2 Percent Slopes-----	1,660	0.5
Jn	Jansen Loam, 0 To 2 Percent Slopes-----	890	0.3
JnC	Jansen Loam, 2 To 6 Percent Slopes-----	2,610	0.8
JnD	Jansen Loam, 6 To 11 Percent Slopes-----	2,130	0.6
LaD	Labu Silty Clay, 6 To 11 Percent Slopes-----	9,480	2.7
LcF	Labu-Sansarc Silty Clays, 11 To 30 Percent Slopes-----	51,150	14.7
Le	Leshara Silt Loam, 0 To 2 Percent Slopes-----	930	0.3
LsC	Lynch Silty Clay, 2 To 6 Percent Slopes-----	1,350	0.4
LsD	Lynch Silty Clay, 6 To 11 Percent Slopes-----	3,150	0.9
LyD	Lynch-Bristow Silty Clays, 6 To 11 Percent Slopes-----	4,350	1.3
LyF	Lynch-Bristow Silty Clays, 11 To 30 Percent Slopes-----	21,310	6.1
MaG	Mariaville-Paka Loams, 15 To 40 Percent Slopes-----	8,990	2.6
MeE	Meadin Sandy Loam, 3 To 17 Percent Slopes-----	3,730	1.1
NoC	Nora Silt Loam, 2 To 6 Percent Slopes-----	3,660	1.1
NoD	Nora Silt Loam, 6 To 11 Percent Slopes-----	1,280	0.4
Oa	Onawa Silty Clay, 0 To 2 Percent Slopes-----	530	0.2
Oe	O'Neill Fine Sandy Loam, 0 To 2 Percent Slopes-----	1,230	0.4
OeC	O'Neill Fine Sandy Loam, 2 To 6 Percent Slopes-----	630	0.2
OfD	O'Neill-Meadin Fine Sandy Loams, 3 To 9 Percent Slopes-----	2,180	0.6
On	Onita Silt Loam, 0 To 2 Percent Slopes-----	40,710	11.7
Or	Ord Fine Sandy Loam, 0 To 2 Percent Slopes-----	1,250	0.4
PaC	Paka Fine Sandy Loam, 2 To 6 Percent Slopes-----	790	0.2
Ph	Paka Loam, 0 To 2 Percent Slopes-----	670	0.2
PhC	Paka Loam, 2 To 6 Percent Slopes-----	1,940	0.6
PhD	Paka Loam, 6 To 11 Percent Slopes-----	2,110	0.6
PoC	Promise Silty Clay, 2 To 6 Percent Slopes-----	2,320	0.7
RaC	Ree Silt Loam, 2 To 6 Percent Slopes-----	14,040	4.0
RaD	Ree Silt Loam, 6 To 11 Percent Slopes-----	5,900	1.7
RaE	Ree Silt Loam, 11 To 15 Percent Slopes-----	1,310	0.4
ReC	Reliance Silt Loam, 2 To 6 Percent Slopes-----	22,860	6.6
ReD	Reliance Silt Loam, 6 To 11 Percent Slopes-----	4,530	1.3
RfC	Reliance Silty Clay Loam, 2 To 6 Percent Slopes-----	1,940	0.6
Rw	Riverwash-----	920	0.3
SaG	Sansarc Silty Clay, 20 To 40 Percent Slopes-----	9,070	2.6
Sc	Scott Silt Loam, 0 To 1 Percent Slopes-----	1,330	0.4
Sm	Simeon Loamy Sand, 0 To 2 Percent Slopes-----	3,100	0.9
SuC	Simeon-Valentine Loamy Sands, 0 To 6 Percent Slopes-----	2,900	0.8
SvF2	Simeon-Valentine Complex, 3 To 30 Percent Slopes, Eroded-----	8,530	2.5
VaE	Valentine Fine Sand, 6 To 17 Percent Slopes-----	5,730	1.6
VbB	Valentine Loamy Sand, 0 To 3 Percent Slopes-----	2,820	0.8
Ve	Verdel Silty Clay, 0 To 2 Percent Slopes-----	2,040	0.6
WeC	Wewela Fine Sandy Loam, 2 To 6 Percent Slopes-----	590	0.2
zwa	Water > 40 Acres-----	6,720	1.9
	Total-----	350,720	101.0

* Less than 0.1 percent.

Nontechnical Soil Descriptions
Boyd County, Nebraska

Nontechnical soil descriptions describe soil properties or management considerations specific to a soil map unit or group of map units, shown in the NonTechnical Descriptions report. These descriptions are written in terminology that Non-technical users of soil survey information can understand. Nontechnical soil descriptions are a powerful tool for creating reports. These high quality, easy to read reports can be generated by conservation planners and other NRCS employees for distribution to land users. Soil map unit descriptions and National Soil Information System records are the basis for these descriptions.

Ab Albaton Silty Clay, 0 To 2 Percent Slopes

Albaton soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping depression on flood plain on river valley. The runoff class is very high. The parent material consists of clayey alluvium. This soil is poorly drained. The slowest permeability is very slow. It has a moderate available water capacity and a very high shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 6 inches. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Clayey Overflow - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

AnC Anselmo Fine Sandy Loam, 2 To 6 Percent Slopes

Anselmo soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping backslope, shoulder, summit hill on tableland. The runoff class is very low. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 3 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

AnD Anselmo Fine Sandy Loam, 6 To 11 Percent Slopes

Anselmo soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping summit, backslope, shoulder hill on tableland. The runoff class is low. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 3 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

AnF Anselmo Fine Sandy Loam, 11 To 20 Percent Slopes

Anselmo soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep summit, shoulder, backslope hill on tableland. The runoff class is low. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 3 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 6e. It is in the nonirrigated land capability classification 6e.

ArF Anselmo-Rock Outcrop Complex, 11 To 20 Percent Slopes

Anselmo soil makes up 75 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep backslope hill on tableland. The runoff class is low. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 3 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 6e. It is in the nonirrigated land capability classification 6e.

Ba Barney Silt Loam, 0 To 2 Percent Slopes

Barney soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is very poorly drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 6 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Wet Land - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 5w.

Nontechnical Soil Descriptions--Continued
Boyd County, Nebraska

Bd Blake Silty Clay Loam, 0 To 2 Percent Slopes

Blake soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is medium. The parent material consists of alluvium. This soil is somewhat poorly drained. The slowest permeability is very slow. It has a very high available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 36 inches. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Silty Lowland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

Be Blendon Fine Sandy Loam, 0 To 2 Percent Slopes

Blendon soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flat on tableland, swale on tableland. The runoff class is very low. The parent material consists of coarse-loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

BeC Blendon Fine Sandy Loam, 2 To 6 Percent Slopes

Blendon soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping footslope valley side on valley, swale on tableland, hill on tableland. The runoff class is very low. The parent material consists of coarse-loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

BoD Boyd Silty Clay, 6 To 11 Percent Slopes

Boyd soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping summit, shoulder, backslope hill on hills. The runoff class is very high. The parent material consists of residuum weathered from clayey shale. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is very slow. It has a low available water capacity and a very high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Clayey - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 4e.

BrG Bristow Silty Clay, 20 To 40 Percent Slopes

Bristow soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a steep to steep backslope, summit, shoulder hill on hills, valley side on valley. The runoff class is very high. The parent material consists of clayey residuum weathered from calcareous, gypsiferous soft shale. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is very slow. It has a very low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 40 percent calcium carbonate. This soil is in the Shallow Limy - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 7s.

Bs Brocksburg Fine Sandy Loam, 0 To 2 Percent Slopes

Brocksburg soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flat on tableland. The runoff class is medium. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Bt Brocksburg Loam, 0 To 2 Percent Slopes

Brocksburg soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level flat on tableland. The runoff class is medium. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2s. It is in the nonirrigated land capability classification 2s.

Nontechnical Soil Descriptions--Continued
Boyd County, Nebraska

Cb Cass Fine Sandy Loam, 0 To 2 Percent Slopes

Cass soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is very low. The parent material consists of sandy alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy Lowland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

CrE2 Crofton Silt Loam, 11 To 15 Percent Slopes, Eroded

Crofton soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep shoulder, backslope, summit loess hill on tableland. The runoff class is medium. The parent material consists of calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 4e.

DuB Dunday Loamy Fine Sand, 0 To 3 Percent Slopes

Dunday soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping hummock on tableland. The runoff class is negligible. The parent material consists of eolian sands. This soil is somewhat excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

DuC Dunday Loamy Fine Sand, 3 To 6 Percent Slopes

Dunday soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping dune on tableland, valley side on valley. The runoff class is very low. The parent material consists of eolian sands. This soil is somewhat excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

DuD Dunday Loamy Fine Sand, 6 To 11 Percent Slopes

Dunday soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping footslope dune on upland, valley side on valley. The runoff class is very low. The parent material consists of eolian sands. This soil is somewhat excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

DxB Dunday Loamy Fine Sand, Loamy Substratum, 0 To 3 Percent Slopes

Dunday soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping hummock on upland. The runoff class is negligible. The parent material consists of eolian sands. This soil is somewhat excessively drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

Et Eltree Silt Loam, 0 To 2 Percent Slopes

Eltree soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level flat on plains. The runoff class is negligible. The parent material consists of calcareous silty colluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 2c.

Go Grigston Silt Loam, 0 To 2 Percent Slopes

Grigston soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is low. The parent material consists of calcareous alluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Silty Lowland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

Nontechnical Soil Descriptions--Continued
Boyd County, Nebraska

GrB Grigston Silt Loam, Channeled, 0 To 3 Percent Slopes

Aowa soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is low. The parent material consists of stratified calcareous alluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty Overflow - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6w.

Ha Hall Silt Loam, 0 To 2 Percent Slopes

Hall soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level stream terrace on valley. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty Lowland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 2c.

He Haynie Silt Loam, 0 To 2 Percent Slopes

Haynie soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is low. The parent material consists of coarse-silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Silty Lowland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 1.

IfD Inavale Fine Sand, 3 To 11 Percent Slopes

Inavale soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping flood plain on valley. The runoff class is very low. The parent material consists of sandy alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

IgB Inavale Fine Sand, Channeled, 0 To 3 Percent Slopes

Inavale soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is negligible. The parent material consists of sandy alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy Lowland - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6w.

IhB Inavale Loamy Fine Sand, 0 To 3 Percent Slopes

Inavale soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is negligible. The parent material consists of sandy alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy Lowland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

In Inavale Fine Sandy Loam, 0 To 2 Percent Slopes

Inavale soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is very low. The parent material consists of sandy alluvium. This soil is excessively drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy Lowland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Jn Jansen Loam, 0 To 2 Percent Slopes

Jansen soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flat on tableland. The runoff class is medium. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2s. It is in the nonirrigated land capability classification 2s.

Nontechnical Soil Descriptions--Continued
Boyd County, Nebraska

JnC Jansen Loam, 2 To 6 Percent Slopes

Jansen soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping backslope, summit, shoulder hill on tableland. The runoff class is medium. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

JnD Jansen Loam, 6 To 11 Percent Slopes

Jansen soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping summit, shoulder, backslope hill on tableland. The runoff class is high. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

LaD Labu Silty Clay, 6 To 11 Percent Slopes

Labu soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping summit, shoulder, backslope hill on plains. The runoff class is very high. The parent material consists of residuum weathered from clayey shale. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is very slow. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Clayey - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

LcF Labu-Sansarc Silty Clays, 11 To 30 Percent Slopes

Labu soil makes up 65 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a strongly sloping to steep backslope hill on hills, valley side on valley. The runoff class is very high. The parent material consists of residuum weathered from clayey shale. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is very slow. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Clayey - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

Sansarc soil makes up 35 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a strongly sloping to steep summit, shoulder hill on hills, valley side on valley. The runoff class is very high. The parent material consists of clayey residuum weathered from shale. The soil is 4 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is slow. It has a very low available water capacity and a very high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Shallow Clay - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

Le Leshara Silt Loam, 0 To 2 Percent Slopes

Leshara soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is low. The parent material consists of stratified loamy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 24 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Subirrigated - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

LsC Lynch Silty Clay, 2 To 6 Percent Slopes

Lynch soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping summit, shoulder, backslope hill on plains, valley side on valley. The runoff class is very high. The parent material consists of clayey residuum weathered from calcareous shale. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is very slow. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 40 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Nontechnical Soil Descriptions--Continued
Boyd County, Nebraska

LsD Lynch Silty Clay, 6 To 11 Percent Slopes

Lynch soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping summit, shoulder, backslope hill on plains. The runoff class is very high. The parent material consists of clayey residuum weathered from calcareous shale. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is very slow. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 40 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

LyD Lynch-Bristow Silty Clays, 6 To 11 Percent Slopes

Lynch soil makes up 65 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping backslope hill on plains. The runoff class is very high. The parent material consists of clayey residuum weathered from calcareous shale. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is very slow. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 40 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

Bristow soil makes up 35 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping summit, shoulder hill on plains. The runoff class is very high. The parent material consists of clayey residuum weathered from calcareous, gypsiferous soft shale. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is very slow. It has a very low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 40 percent calcium carbonate. This soil is in the Shallow Limy - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

LyF Lynch-Bristow Silty Clays, 11 To 30 Percent Slopes

Lynch soil makes up 55 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a strongly sloping to steep backslope valley side on valley. The runoff class is very high. The parent material consists of clayey residuum weathered from calcareous shale. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is very slow. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 40 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

Bristow soil makes up 45 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a strongly sloping to steep shoulder, summit valley side on valley. The runoff class is very high. The parent material consists of clayey residuum weathered from calcareous, gypsiferous soft shale. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is very slow. It has a very low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 40 percent calcium carbonate. This soil is in the Shallow Limy - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

MaG Mariaville-Paka Loams, 15 To 40 Percent Slopes

Mariaville soil makes up 55 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately steep to steep backslope valley side on valley. The runoff class is very high. The parent material consists of residuum weathered from siltstone. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately slow. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 14 percent calcium carbonate. This soil is in the Shallow Limy - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

Paka soil makes up 45 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately steep to steep summit, shoulder valley side on valley. The runoff class is very high. The parent material consists of loamy residuum weathered from siltstone. The soil is 40 to 60 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

MeE Meadin Sandy Loam, 3 To 17 Percent Slopes

Meadin soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep hill on tableland, valley side on valley. The runoff class is low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

Nontechnical Soil Descriptions--Continued
Boyd County, Nebraska

NoC Nora Silt Loam, 2 To 6 Percent Slopes

Nora soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping backslope, summit, shoulder loess hill on tableland. The runoff class is low. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 2e.

NoD Nora Silt Loam, 6 To 11 Percent Slopes

Nora soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping summit, backslope, shoulder loess hill on tableland. The runoff class is medium. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 3e.

Oa Onawa Silty Clay, 0 To 2 Percent Slopes

Onawa soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on flood plain on valley. The runoff class is negligible. The parent material consists of clayey alluvium over loamy alluvium. This soil is somewhat poorly drained. The slowest permeability is very slow. It has a high available water capacity and a high shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 36 inches. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Clayey Overflow - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

Oe O'Neill Fine Sandy Loam, 0 To 2 Percent Slopes

O'Neill soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flat on tableland. The runoff class is very low. The parent material consists of coarse-loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

OeC O'Neill Fine Sandy Loam, 2 To 6 Percent Slopes

O'Neill soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping summit, backslope, shoulder hill on tableland. The runoff class is very low. The parent material consists of coarse-loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

OFD O'Neill-Meadin Fine Sandy Loams, 3 To 9 Percent Slopes

O'Neill soil makes up 55 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping backslope, summit hill on tableland. The runoff class is low. The parent material consists of coarse-loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

Meadin soil makes up 45 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping shoulder, summit hill on tableland. The runoff class is very low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

On Onita Silt Loam, 0 To 2 Percent Slopes

Onita soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flat on tableland, swale on tableland. The runoff class is low. The parent material consists of loamy colluvium. This soil is well drained. The slowest permeability is very slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 1 It is in the nonirrigated land capability classification 2c.

Nontechnical Soil Descriptions--Continued
Boyd County, Nebraska

Or Ord Fine Sandy Loam, 0 To 2 Percent Slopes

Ord soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping interdune on sandhills, stream terrace on valley. The runoff class is very low. The parent material consists of stratified sandy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 40 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Subirrigated - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

PaC Paka Fine Sandy Loam, 2 To 6 Percent Slopes

Paka soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping backslope, shoulder, summit hill on tableland. The runoff class is medium. The parent material consists of loamy residuum weathered from siltstone. The soil is 40 to 60 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Ph Paka Loam, 0 To 2 Percent Slopes

Paka soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level flat on tableland. The runoff class is medium. The parent material consists of loamy residuum weathered from siltstone. The soil is 40 to 60 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 1 It is in the nonirrigated land capability classification 2c.

PhC Paka Loam, 2 To 6 Percent Slopes

Paka soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping summit, shoulder, backslope hill on tableland. The runoff class is medium. The parent material consists of loamy residuum weathered from siltstone. The soil is 40 to 60 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

PhD Paka Loam, 6 To 11 Percent Slopes

Paka soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping summit, backslope, shoulder hill on tableland. The runoff class is high. The parent material consists of loamy residuum weathered from siltstone. The soil is 40 to 60 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

PoC Promise Silty Clay, 2 To 6 Percent Slopes

Promise soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping hill on plains, valley side on valley. The runoff class is very high. The parent material consists of clayey residuum weathered from shale. This soil is well drained. The slowest permeability is very slow. It has a moderate available water capacity and a very high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil contains a very slightly saline horizon, it has a horizon that is slightly sodic. This soil is in the Clayey - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 3e.

RaC Ree Silt Loam, 2 To 6 Percent Slopes

Ree soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping backslope, shoulder, summit hill on tableland. The runoff class is medium. The parent material consists of loamy alluvial sediments. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Nontechnical Soil Descriptions--Continued
Boyd County, Nebraska

RaD Ree Silt Loam, 6 To 11 Percent Slopes

Ree soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping shoulder, backslope, summit hill on tableland. The runoff class is high. The parent material consists of loamy alluvial sediments. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

RaE Ree Silt Loam, 11 To 15 Percent Slopes

Ree soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep summit, backslope, shoulder hill on tableland. The runoff class is high. The parent material consists of loamy alluvial sediments. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

ReC Reliance Silt Loam, 2 To 6 Percent Slopes

Reliance soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping backslope, summit, shoulder loess hill on tableland. The runoff class is high. The parent material consists of loess. This soil is well drained. The slowest permeability is slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 20 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

ReD Reliance Silt Loam, 6 To 11 Percent Slopes

Reliance soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping summit, shoulder, backslope loess hill on tableland. The runoff class is very high. The parent material consists of loess. This soil is well drained. The slowest permeability is slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 20 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

RfC Reliance Silty Clay Loam, 2 To 6 Percent Slopes

Reliance soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping summit, shoulder, backslope loess hill on tableland. The runoff class is medium. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 20 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Rw Riverwash

Riverwash soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on valley. The runoff class is negligible. The parent material consists of sandy and gravelly alluvium. This soil is very poorly drained. The slowest permeability is rapid. It has a very low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 0 inches. The soil contains a maximum amount of 5 percent calcium carbonate. It is in the nonirrigated land capability classification 8w.

SaG Sansarc Silty Clay, 20 To 40 Percent Slopes

Sansarc soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a steep to steep hill on hills, valley side on valley. The runoff class is very high. The parent material consists of clayey residuum weathered from shale. The soil is 4 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is slow. It has a very low available water capacity and a very high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Shallow Clay - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 7s.

Sc Scott Silt Loam, 0 To 1 Percent Slopes

Scott soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level playa on tableland. The runoff class is negligible. The parent material consists of loess. This soil is poorly drained. The slowest permeability is very slow. It has a moderate available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 0 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Clayey Overflow - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 4w.

Nontechnical Soil Descriptions--Continued
Boyd County, Nebraska

Sm Simeon Loamy Sand, 0 To 2 Percent Slopes

Simeon soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flat on tableland. The runoff class is negligible. The parent material consists of sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4s. It is in the nonirrigated land capability classification 6s.

SuC Simeon-Valentine Loamy Sands, 0 To 6 Percent Slopes

Simeon soil makes up 55 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to moderately sloping swale, upland. The runoff class is negligible. The parent material consists of sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4s. It is in the nonirrigated land capability classification 6s.

Valentine soil makes up 45 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to moderately sloping dune, upland. The runoff class is negligible. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

SvF2 Simeon-Valentine Complex, 3 To 30 Percent Slopes, Eroded

Simeon soil makes up 70 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to steep summit, shoulder hill on upland, valley side on valley. The runoff class is low. The parent material consists of sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

Valentine soil makes up 30 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to steep backslope hill on tableland, valley side on valley. The runoff class is low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

VaE Valentine Fine Sand, 6 To 17 Percent Slopes

Valentine soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep dune on sandhills. The runoff class is very low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

VbB Valentine Loamy Sand, 0 To 3 Percent Slopes

Valentine soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping hummock on upland. The runoff class is negligible. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

Ve Verdel Silty Clay, 0 To 2 Percent Slopes

Verdel soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a nearly level stream terrace on valley. The runoff class is high. The parent material consists of clayey alluvium derived from shale. This soil is well drained. The slowest permeability is very slow. It has a moderate available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Clayey - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2s. It is in the nonirrigated land capability classification 2s.

Nontechnical Soil Descriptions--Continued
Boyd County, Nebraska

WeC Wewela Fine Sandy Loam, 2 To 6 Percent Slopes

Wewela soil makes up 100 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping shoulder, backslope, summit hill on plains. The runoff class is very high. The parent material consists of loamy eolian deposits over clayey residuum weathered from clayey shale. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is very slow. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Ab—Albaton silty clay, 0 to 2 percent slopes

Map Unit Composition

Albaton: 100 percent

Component Descriptions

Albaton

MLRA: 63B - Southern Rolling Pierre Shale
Plains

Landform: Depression on flood plain on river valley

Parent material: Clayey alluvium

Slope: 0 to 2 percent

Drainage class: Poorly drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: Moderate (About 7.1 inches)

Shrink-swell potential: Very high (About 17.0 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: About 0 to 12 inches

Runoff class: Very high

Ecological site: Clayey Overflow - Veg. Zone 3

Land capability (irrigated): 3w

Land capability (nonirrigated): 3w

Typical Profile:

H1—0 to 7 inches; silty clay

H2—7 to 54 inches; stratified silty clay to silt loam

H3—54 to 60 inches; silty clay

AnC—Anselmo fine sandy loam, 2 to 6 percent slopes

Map Unit Composition

Anselmo: 100 percent

Component Descriptions

Anselmo

MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Hill on tableland

Hillslope position: Backslope, shoulder, summit

Parent material: Loamy eolian deposits

Slope: 2 to 6 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 1.98 in/hr)

Available water capacity: Moderate (About 7.2 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Sandy - Veg. Zone 3

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 16 inches; fine sandy loam

H2—16 to 21 inches; fine sandy loam

H3—21 to 34 inches; fine sandy loam

H4—34 to 60 inches; loamy fine sand

AnD—Anselmo fine sandy loam, 6 to 11 percent slopes

Map Unit Composition

Anselmo: 100 percent

Component Descriptions

Anselmo

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Hill on tableland

Hillslope position: Summit, backslope, shoulder

Parent material: Loamy eolian deposits

Slope: 6 to 11 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 1.98 in/hr)

Available water capacity: Moderate (About 7.2 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sandy - Veg. Zone 3

Land capability (irrigated): 4e

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 16 inches; fine sandy loam

H2—16 to 21 inches; fine sandy loam

H3—21 to 34 inches; fine sandy loam

H4—34 to 60 inches; loamy fine sand

AnF—Anselmo fine sandy loam, 11 to 20 percent slopes

Map Unit Composition

Anselmo: 100 percent

Component Descriptions

Anselmo

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Hill on tableland

Hillslope position: Summit, shoulder, backslope

Parent material: Loamy eolian deposits

Slope: 11 to 17 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 1.98 in/hr)

Available water capacity: Moderate (About 7.2 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sandy - Veg. Zone 3

Land capability (irrigated): 6e

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 16 inches; fine sandy loam
H2—16 to 21 inches; fine sandy loam
H3—21 to 34 inches; fine sandy loam
H4—34 to 60 inches; loamy fine sand

Parent material: Loamy eolian deposits

Slope: 11 to 17 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 1.98 in/hr)

Available water capacity: Moderate (About 7.2 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sandy - Veg. Zone 3

Land capability (irrigated): 6e

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 16 inches; fine sandy loam
H2—16 to 21 inches; fine sandy loam
H3—21 to 34 inches; fine sandy loam
H4—34 to 60 inches; loamy fine sand

Rock outcrop

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Hill on tableland

Hillslope position: Shoulder, summit

Slope: 11 to 20 percent

Depth to restrictive feature: 0 inches to bedrock (paralithic)

Available water capacity: Very low (About 0.0 inches)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: No Site - Veg. Zone 3

Land capability (nonirrigated): 8s

ArF—Anselmo-Rock outcrop complex, 11 to 20 percent slopes

Map Unit Composition

Anselmo: 75 percent

Rock outcrop: 25 percent

Component Descriptions

Anselmo

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Hill on tableland

Hillslope position: Backslope

Ba—Barney silt loam, 0 to 2 percent slopes

Map Unit Composition

Barney: 100 percent

Component Descriptions

Barney

MLRA: 63B - Southern Rolling Pierre Shale Plains

Landform: Flood plain on valley

Parent material: Loamy alluvium over sandy and gravelly alluvium

Slope: 0 to 2 percent

Drainage class: Very poorly drained
Slowest permeability: Moderate (About 0.57 in/hr)
Available water capacity: Low (About 4.6 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Frequent
Depth to seasonal water saturation: About 0 to 12 inches
Runoff class: Low
Ecological site: Wet Land - Veg. Zone 3
Land capability (nonirrigated): 5w

Typical Profile:

H1—0 to 5 inches; silt loam
 H2—5 to 10 inches; stratified loam to sand
 H3—10 to 60 inches; fine sand

Bd—Blake silty clay loam, 0 to 2 percent slopes

Map Unit Composition

Blake: 100 percent

Component Descriptions

Blake

MLRA: 63B - Southern Rolling Pierre Shale Plains

Landform: Flood plain on valley

Parent material: Alluvium

Slope: 0 to 2 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Very slow (About 0.06 in/hr)

Available water capacity: Very high (About 12.4 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: About 24 to 48 inches

Runoff class: Medium

Ecological site: Silty Lowland - Veg. Zone 3

Land capability (irrigated): 2w

Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 8 inches; silty clay loam
 H2—8 to 25 inches; stratified silty clay loam to silty clay
 H3—25 to 60 inches; stratified very fine sandy loam to silty clay loam

Minor Components

Albaton

Slope: 0 to 2 percent

Drainage class: Poorly drained

Ecological site: Clayey Overflow - Veg. Zone 3

Be—Blendon fine sandy loam, 0 to 2 percent slopes

Map Unit Composition

Blendon: 100 percent

Component Descriptions

Blendon

MLRA: 63B - Southern Rolling Pierre Shale Plains

Landform: Flat on tableland, swale on tableland

Parent material: Coarse-loamy eolian deposits

Slope: 0 to 2 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 1.98 in/hr)

Available water capacity: Moderate (About 7.4 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Sandy - Veg. Zone 3

Land capability (irrigated): 2e

Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 16 inches; fine sandy loam
 H2—16 to 42 inches; sandy loam
 H3—42 to 60 inches; loamy sand

BeC—Blendon fine sandy loam, 2 to 6 percent slopes

Map Unit Composition

Blendon: 100 percent

Component Descriptions

Blendon

MLRA: 63B - Southern Rolling Pierre Shale
Plains

Landform: Valley side on valley, swale on
tableland, hill on tableland

Hillslope position: Foothills

Parent material: Coarse-loamy eolian deposits

Slope: 2 to 6 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About
1.98 in/hr)

Available water capacity: Moderate (About 7.4
inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6
feet

Runoff class: Very low

Ecological site: Sandy - Veg. Zone 3

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 16 inches; fine sandy loam

H2—16 to 42 inches; sandy loam

H3—42 to 60 inches; loamy sand

Minor Components**Ponded Soils****BoD—Boyd silty clay, 6 to 11
percent slopes**

Map Unit Composition

Boyd: 100 percent

Component Descriptions

Boyd

MLRA: 63B - Southern Rolling Pierre Shale
Plains

Landform: Hill on hills

Hillslope position: Summit, shoulder, backslope

Parent material: Residuum weathered from
clayey shale

Slope: 6 to 11 percent

Depth to restrictive feature: 20 to 40 inches to
bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Very slow (About 0.00
in/hr)

Available water capacity: Low (About 3.1 inches)

Shrink-swell potential: Very high (About 17.0
LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6
feet

Runoff class: Very high

Ecological site: Clayey - Veg. Zone 3

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 6 inches; silty clay

H2—6 to 22 inches; clay

H3—22 to 32 inches; clay

Cr—32 to 60 inches; weathered bedrock

**BrG—Bristow silty clay, 20 to 40
percent slopes**

Map Unit Composition

Bristow: 100 percent

Component Descriptions

Bristow

MLRA: 63B - Southern Rolling Pierre Shale
Plains

Landform: Hill on hills, valley side on valley

Hillslope position: Backslope, summit, shoulder

Parent material: Clayey residuum weathered
from calcareous, gypsiferous soft shale

Slope: 20 to 40 percent

Depth to restrictive feature: 10 to 20 inches to
bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Very slow (About 0.00
in/hr)

Available water capacity: Very low (About 1.5
inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6
feet

Runoff class: Very high

Ecological site: Shallow Limy - Veg. Zone 3

Land capability (nonirrigated): 7s

Typical Profile:

H1—0 to 7 inches; silty clay

H2—7 to 17 inches; clay

Cr—17 to 80 inches; weathered bedrock

Bs—Brocksburg fine sandy loam, 0 to 2 percent slopes

Map Unit Composition

Brocksburg: 100 percent

Component Descriptions

Brocksburg

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Flat on tableland

Parent material: Loamy alluvium over sandy and gravelly alluvium

Slope: 0 to 2 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: Moderate (About 6.2 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Sandy - Veg. Zone 3

Land capability (irrigated): 2e

Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 18 inches; fine sandy loam

H2—18 to 32 inches; clay loam

H3—32 to 60 inches; gravelly sand

Minor Components

Scott

Slope: 0 to 1 percent

Drainage class: Poorly drained

Ecological site: Clayey Overflow - Veg. Zone 3

Bt—Brocksburg loam, 0 to 2 percent slopes

Map Unit Composition

Brocksburg: 100 percent

Component Descriptions

Brocksburg

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Flat on tableland

Parent material: Loamy alluvium over sandy and gravelly alluvium

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: Moderate (About 7.2 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 2s

Land capability (nonirrigated): 2s

Typical Profile:

H1—0 to 14 inches; loam

H2—14 to 34 inches; clay loam

H3—34 to 60 inches; gravelly coarse sand

Minor Components

Scott

Slope: 0 to 1 percent

Drainage class: Poorly drained

Ecological site: Clayey Overflow - Veg. Zone 3

Cb—Cass fine sandy loam, 0 to 2 percent slopes

Map Unit Composition

Cass: 100 percent

Component Descriptions

Cass

MLRA: 63B - Southern Rolling Pierre Shale Plains

Landform: Flood plain on valley

Parent material: Sandy alluvium

Slope: 0 to 2 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Moderate (About 7.3 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Sandy Lowland - Veg. Zone 3

Land capability (irrigated): 2w

Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 12 inches; fine sandy loam

H2—12 to 26 inches; fine sandy loam

H3—26 to 60 inches; stratified coarse sand to fine sandy loam

CrE2—Crofton silt loam, 11 to 15 percent slopes, Eroded

Map Unit Composition

Crofton: 100 percent

Component Descriptions

Crofton

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Loess hill on tableland

Hillslope position: Shoulder, backslope, summit

Parent material: Calcareous loess

Slope: 11 to 15 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 12.0 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Limy Upland - Veg. Zone 3

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 6 inches; silt loam

H2—6 to 60 inches; silt loam

DuB—Dunday loamy fine sand, 0 to 3 percent slopes

Map Unit Composition

Dunday: 100 percent

Component Descriptions

Dunday

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Hummock on tableland

Parent material: Eolian sands

Slope: 0 to 3 percent

Drainage class: Somewhat excessively drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Low (About 5.2 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Sandy - Veg. Zone 3

Land capability (irrigated): 3e

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 15 inches; loamy fine sand

H2—15 to 60 inches; fine sand

DuC—Dunday loamy fine sand, 3 to 6 percent slopes

Map Unit Composition

Dunday: 100 percent

Component Descriptions

Dunday

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Dune on tableland, valley side on valley

Parent material: Eolian sands

Slope: 3 to 6 percent

Drainage class: Somewhat excessively drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Low (About 5.2 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Sandy - Veg. Zone 3

Land capability (irrigated): 4e

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 15 inches; loamy fine sand

H2—15 to 60 inches; fine sand

DuD—Dunday loamy fine sand, 6 to 11 percent slopes

Map Unit Composition

Dunday: 100 percent

Component Descriptions

Dunday

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Dune on upland, valley side on valley

Hillslope position: Footslope

Parent material: Eolian sands

Slope: 6 to 11 percent

Drainage class: Somewhat excessively drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Low (About 5.2 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Sandy - Veg. Zone 3

Land capability (irrigated): 4e

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 15 inches; loamy fine sand

H2—15 to 60 inches; fine sand

DxB—Dunday loamy fine sand, loamy Substratum, 0 to 3 percent slopes

Map Unit Composition

Dunday: 100 percent

Component Descriptions

Dunday

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Hummock on upland

Parent material: Eolian sands

Slope: 0 to 3 percent

Drainage class: Somewhat excessively drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Low (About 5.2 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Sandy - Veg. Zone 3

Land capability (irrigated): 3e

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 14 inches; loamy fine sand

H2—14 to 34 inches; loamy sand

H3—34 to 48 inches; sandy loam

H4—48 to 60 inches; coarse sand

Et—Eltree silt loam, 0 to 2 percent slopes

Map Unit Composition

Eltree: 100 percent

Component Descriptions

Eltree

MLRA: 63B - Southern Rolling Pierre Shale Plains

Landform: Flat on plains

Parent material: Calcareous silty colluvium

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Very high (About 12.1 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 1

Land capability (nonirrigated): 2c

Typical Profile:

H1—0 to 25 inches; silt loam

H2—25 to 38 inches; silt loam

H3—38 to 60 inches; silt loam

Minor Components

Scott

Slope: 0 to 1 percent

Drainage class: Poorly drained
Ecological site: Clayey Overflow - Veg. Zone 3

Go—Grigston silt loam, 0 to 2 percent slopes

Map Unit Composition

Grigston: 100 percent

Component Descriptions

Grigston

MLRA: 63B - Southern Rolling Pierre Shale Plains

Landform: Flood plain on valley

Parent material: Calcareous alluvium

Slope: 0 to 2 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 12.0 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Silty Lowland - Veg. Zone 3

Land capability (irrigated): 2w

Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 15 inches; silt loam

H2—15 to 31 inches; silt loam

H3—31 to 60 inches; stratified loam to fine sandy loam

Minor Components

Wt At 0-1 Foot

GrB—Grigston silt loam, Channeled, 0 to 3 percent slopes

Map Unit Composition

Aowa: 100 percent

Component Descriptions

Aowa

MLRA: 63B - Southern Rolling Pierre Shale Plains

Landform: Flood plain on valley

Parent material: Stratified calcareous alluvium

Slope: 0 to 2 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Very high (About 12.3 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Frequent

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Silty Overflow - Veg. Zone 3

Land capability (nonirrigated): 6w

Typical Profile:

H1—0 to 15 inches; silt loam

H2—15 to 60 inches; stratified silt loam to fine sand

Minor Components

Wt At 0-1 Foot

Ha—Hall silt loam, 0 to 2 percent slopes

Map Unit Composition

Hall: 100 percent

Component Descriptions

Hall

MLRA: 63B - Southern Rolling Pierre Shale Plains

Landform: Stream terrace on valley

Parent material: Loess

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 11.1 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Silty Lowland - Veg. Zone 3
Land capability (irrigated): 1
Land capability (nonirrigated): 2c

Typical Profile:

H1—0 to 5 inches; silt loam
 H2—5 to 36 inches; silty clay loam
 H3—36 to 42 inches; silty clay loam
 H4—42 to 60 inches; stratified silty clay loam
 to fine sandy loam

Minor Components
Wt At 0-1 Foot

He—Haynie silt loam, 0 to 2 percent slopes

Map Unit Composition

Haynie: 100 percent

Component Descriptions

Haynie
 MLRA: 63B - Southern Rolling Pierre Shale
 Plains

Landform: Flood plain on valley
Parent material: Coarse-silty alluvium
Slope: 0 to 2 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Very high (About 12.4 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Occasional
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Silty Lowland - Veg. Zone 3
Land capability (irrigated): 1
Land capability (nonirrigated): 1

Typical Profile:

H1—0 to 7 inches; silt loam
 H2—7 to 60 inches; stratified silt loam to fine sand

IfD—Inavale fine sand, 3 to 11 percent slopes

Map Unit Composition

Inavale: 100 percent

Component Descriptions

Inavale
 MLRA: 63B - Southern Rolling Pierre Shale
 Plains
Landform: Flood plain on valley
Parent material: Sandy alluvium
Slope: 3 to 11 percent
Drainage class: Excessively drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 4.8 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Rare
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sands - Veg. Zone 3
Land capability (irrigated): 4e
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 6 inches; fine sand
 H2—6 to 12 inches; fine sand
 H3—12 to 60 inches; fine sand

IgB—Inavale fine sand, Channeled, 0 to 3 percent slopes

Map Unit Composition

Inavale: 100 percent

Component Descriptions

Inavale
 MLRA: 63B - Southern Rolling Pierre Shale
 Plains
Landform: Flood plain on valley
Parent material: Sandy alluvium
Slope: 0 to 3 percent
Drainage class: Excessively drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 4.8 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Frequent
Depth to seasonal water saturation: More than 6 feet
Runoff class: Negligible
Ecological site: Sandy Lowland - Veg. Zone 3
Land capability (nonirrigated): 6w

Typical Profile:

H1—0 to 6 inches; fine sand
 H2—6 to 12 inches; fine sand
 H3—12 to 60 inches; fine sand

Minor Components**Barney**

Slope: 0 to 2 percent
Drainage class: Very poorly drained
Ecological site: Wet Land - Veg. Zone 3

1hB—Inavale loamy fine sand, 0 to 3 percent slopes

Map Unit Composition

Inavale: 100 percent

Component Descriptions

Inavale

MLRA: 63B - Southern Rolling Pierre Shale
 Plains

Landform: Flood plain on valley

Parent material: Sandy alluvium

Slope: 0 to 3 percent

Drainage class: Excessively drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Low (About 5.1 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Sandy Lowland - Veg. Zone 3

Land capability (irrigated): 3e

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 9 inches; loamy fine sand
 H2—9 to 19 inches; fine sand
 H3—19 to 60 inches; loamy fine sand

1n—Inavale fine sandy loam, 0 to 2 percent slopes

Map Unit Composition

Inavale: 100 percent

Component Descriptions

Inavale

MLRA: 63B - Southern Rolling Pierre Shale
 Plains

Landform: Flood plain on valley

Parent material: Sandy alluvium

Slope: 0 to 3 percent

Drainage class: Excessively drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Moderate (About 6.1 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Sandy Lowland - Veg. Zone 3

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 16 inches; fine sandy loam
 H2—16 to 26 inches; fine sand
 H3—26 to 60 inches; fine sand

Jn—Jansen loam, 0 to 2 percent slopes

Map Unit Composition

Jansen: 100 percent

Component Descriptions

Jansen

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Flat on tableland

Parent material: Loamy alluvium over sandy and gravelly alluvium

Slope: 0 to 2 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: Moderate (About 6.9 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 2s

Land capability (nonirrigated): 2s

Typical Profile:

- H1—0 to 13 inches; loam
- H2—13 to 32 inches; clay loam
- H3—32 to 60 inches; gravelly coarse sand

JnC—Jansen loam, 2 to 6 percent slopes

Map Unit Composition

Jansen: 100 percent

Component Descriptions

Jansen
MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Hill on tableland
Hillslope position: Backslope, summit, shoulder
Parent material: Loamy alluvium over sandy and gravelly alluvium
Slope: 2 to 6 percent
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: Moderate (About 6.9 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Silty - Veg. Zone 3
Land capability (irrigated): 3e
Land capability (nonirrigated): 3e

Typical Profile:

- H1—0 to 13 inches; loam
- H2—13 to 32 inches; clay loam
- H3—32 to 60 inches; gravelly coarse sand

JnD—Jansen loam, 6 to 11 percent slopes

Map Unit Composition

Jansen: 100 percent

Component Descriptions

Jansen

MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Hill on tableland
Hillslope position: Summit, shoulder, backslope
Parent material: Loamy alluvium over sandy and gravelly alluvium
Slope: 6 to 11 percent
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: Moderate (About 6.9 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: High
Ecological site: Silty - Veg. Zone 3
Land capability (irrigated): 4e
Land capability (nonirrigated): 4e

Typical Profile:

- H1—0 to 13 inches; loam
- H2—13 to 32 inches; clay loam
- H3—32 to 60 inches; gravelly coarse sand

LaD—Labu silty clay, 6 to 11 percent slopes

Map Unit Composition

Labu: 100 percent

Component Descriptions

Labu
MLRA: 63B - Southern Rolling Pierre Shale Plains
Landform: Hill on plains
Hillslope position: Summit, shoulder, backslope
Parent material: Residuum weathered from clayey shale
Slope: 6 to 11 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: Low (About 3.3 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high
Ecological site: Clayey - Veg. Zone 3
Land capability (irrigated): 4e
Land capability (nonirrigated): 4e

Typical Profile:
 H1—0 to 6 inches; silty clay
 H2—6 to 34 inches; clay
 Cr—34 to 60 inches; weathered bedrock

Minor Components
Wt At 0-1 Foot

LcF—Labu-Sansarc silty clays, 11 to 30 percent slopes

Map Unit Composition

Labu: 65 percent
 Sansarc: 35 percent

Component Descriptions

Labu

MLRA: 63B - Southern Rolling Pierre Shale Plains

Landform: Hill on hills, valley side on valley

Hillslope position: Backslope

Parent material: Residuum weathered from clayey shale

Slope: 11 to 30 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: Low (About 3.3 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Clayey - Veg. Zone 3

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 6 inches; silty clay
 H2—6 to 34 inches; clay
 Cr—34 to 60 inches; weathered bedrock

Sansarc

MLRA: 63B - Southern Rolling Pierre Shale Plains

Landform: Hill on hills, valley side on valley

Hillslope position: Summit, shoulder
Parent material: Clayey residuum weathered from shale

Slope: 11 to 30 percent

Depth to restrictive feature: 4 to 20 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Slow (About 0.06 in/hr)

Available water capacity: Very low (About 1.1 inches)

Shrink-swell potential: Very high (About 17.0 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Shallow Clay - Veg. Zone 3

Land capability (nonirrigated): 6s

Typical Profile:

H1—0 to 14 inches; silty clay
 Cr—14 to 60 inches; weathered bedrock

Minor Components
Wt At 0-1 Foot

Le—Leshara silt loam, 0 to 2 percent slopes

Map Unit Composition

Leshara: 100 percent

Component Descriptions

Leshara

MLRA: 63B - Southern Rolling Pierre Shale Plains

Landform: Flood plain on valley

Parent material: Stratified loamy alluvium

Slope: 0 to 2 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Very high (About 12.6 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: About 12 to 36 inches

Runoff class: Low

Ecological site: Subirrigated - Veg. Zone 3

Land capability (irrigated): 2w

Land capability (nonirrigated): 2w

Typical Profile:

- H1—0 to 21 inches; silt loam
- H2—21 to 32 inches; silt loam
- H3—32 to 60 inches; stratified silt loam to fine sand

Minor Components
Poorly Drained

LsC—Lynch silty clay, 2 to 6 percent slopes

Map Unit Composition

Lynch: 100 percent

Component Descriptions

Lynch

MLRA: 63B - Southern Rolling Pierre Shale
Plains

Landform: Hill on plains, valley side on valley

Hillslope position: Summit, shoulder, backslope

Parent material: Clayey residuum weathered from calcareous shale

Slope: 2 to 6 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: Low (About 3.5 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Limy Upland - Veg. Zone 3

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

- H1—0 to 8 inches; silty clay
- H2—8 to 28 inches; silty clay
- H3—28 to 36 inches; silty clay
- Cr—36 to 60 inches; weathered bedrock

LsD—Lynch silty clay, 6 to 11 percent slopes

Map Unit Composition

Lynch: 100 percent

Component Descriptions

Lynch

MLRA: 63B - Southern Rolling Pierre Shale
Plains

Landform: Hill on plains

Hillslope position: Summit, shoulder, backslope

Parent material: Clayey residuum weathered from calcareous shale

Slope: 6 to 11 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: Low (About 3.5 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Limy Upland - Veg. Zone 3

Land capability (irrigated): 4e

Land capability (nonirrigated): 4e

Typical Profile:

- H1—0 to 8 inches; silty clay
- H2—8 to 28 inches; silty clay
- H3—28 to 36 inches; silty clay
- Cr—36 to 60 inches; weathered bedrock

LyD—Lynch-Bristow silty clays, 6 to 11 percent slopes

Map Unit Composition

Lynch: 65 percent

Bristow: 35 percent

Component Descriptions

Lynch

MLRA: 63B - Southern Rolling Pierre Shale
Plains

Landform: Hill on plains

Hillslope position: Backslope

Parent material: Clayey residuum weathered from calcareous shale

Slope: 6 to 11 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: Low (About 3.5 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very high
Ecological site: Limy Upland - Veg. Zone 3
Land capability (irrigated): 4e
Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 8 inches; silty clay
 H2—8 to 28 inches; silty clay
 H3—28 to 36 inches; silty clay
 Cr—36 to 60 inches; weathered bedrock

Bristow

MLRA: 63B - Southern Rolling Pierre Shale Plains
Landform: Hill on plains
Hillslope position: Summit, shoulder
Parent material: Clayey residuum weathered from calcareous, gypsiferous soft shale
Slope: 6 to 11 percent
Depth to restrictive feature: 10 to 20 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: Very low (About 1.5 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very high
Ecological site: Shallow Limy - Veg. Zone 3
Land capability (nonirrigated): 6s

Typical Profile:

H1—0 to 7 inches; silty clay
 H2—7 to 17 inches; clay
 Cr—17 to 60 inches; weathered bedrock

LyF—Lynch-Bristow silty clays, 11 to 30 percent slopes

Map Unit Composition

Lynch: 55 percent
 Bristow: 45 percent

Component Descriptions

Lynch

MLRA: 63B - Southern Rolling Pierre Shale Plains
Landform: Valley side on valley
Hillslope position: Backslope
Parent material: Clayey residuum weathered from calcareous shale
Slope: 11 to 30 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: Low (About 3.5 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very high
Ecological site: Limy Upland - Veg. Zone 3
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 8 inches; silty clay
 H2—8 to 28 inches; silty clay
 H3—28 to 36 inches; silty clay
 Cr—36 to 60 inches; weathered bedrock

Bristow

MLRA: 63B - Southern Rolling Pierre Shale Plains
Landform: Valley side on valley
Hillslope position: Shoulder, summit
Parent material: Clayey residuum weathered from calcareous, gypsiferous soft shale
Slope: 11 to 30 percent
Depth to restrictive feature: 10 to 20 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: Very low (About 1.5 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very high
Ecological site: Shallow Limy - Veg. Zone 3
Land capability (nonirrigated): 6s

Typical Profile:

H1—0 to 7 inches; silty clay
 H2—7 to 17 inches; clay
 Cr—17 to 60 inches; weathered bedrock

MaG—Mariaville-Paka loams, 15 to 40 percent slopes

Map Unit Composition

Mariaville: 55 percent
 Paka: 45 percent

Component Descriptions

Mariaville

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Valley side on valley

Hillslope position: Backslope

Parent material: Residuum weathered from siltstone

Slope: 15 to 40 percent

Depth to restrictive feature: 10 to 20 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: Low (About 3.0 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Shallow Limy - Veg. Zone 3

Land capability (nonirrigated): 6s

Typical Profile:

H1—0 to 5 inches; loam
 H2—5 to 17 inches; silty clay loam
 Cr—17 to 60 inches; weathered bedrock

Paka

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Valley side on valley

Hillslope position: Summit, shoulder

Parent material: Loamy residuum weathered from siltstone

Slope: 15 to 40 percent

Depth to restrictive feature: 40 to 60 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 9.7 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Silty - Veg. Zone 3

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 8 inches; loam
 H2—8 to 26 inches; clay loam
 H3—26 to 48 inches; silt loam
 Cr—48 to 60 inches; weathered bedrock

Minor Components

Wt At 0-1 Foot

MeE—Meadin sandy loam, 3 to 17 percent slopes

Map Unit Composition

Meadin: 100 percent

Component Descriptions

Meadin

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Hill on tableland, valley side on valley

Parent material: Loamy alluvium over sandy and gravelly alluvium

Slope: 3 to 17 percent

Drainage class: Excessively drained

Slowest permeability: Moderately rapid (About 1.98 in/hr)

Available water capacity: Low (About 3.6 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Shallow To Gravel - Veg. Zone 3

Land capability (nonirrigated): 6s

Typical Profile:

H1—0 to 8 inches; sandy loam
 H2—8 to 13 inches; gravelly sandy loam
 H3—13 to 60 inches; very gravelly coarse sand

NoC—Nora silt loam, 2 to 6 percent slopes

Map Unit Composition

Nora: 100 percent

Component Descriptions

Nora

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Loess hill on tableland

Hillslope position: Backslope, summit, shoulder

Parent material: Fine-silty calcareous loess

Slope: 2 to 6 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 11.4 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 3e

Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 7 inches; silt loam

H2—7 to 34 inches; silt loam

H3—34 to 60 inches; silt loam

NoD—Nora silt loam, 6 to 11 percent slopes

Map Unit Composition

Nora: 100 percent

Component Descriptions

Nora

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Loess hill on tableland

Hillslope position: Summit, backslope, shoulder

Parent material: Fine-silty calcareous loess

Slope: 6 to 11 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 11.4 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 4e

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 7 inches; silt loam

H2—7 to 34 inches; silt loam

H3—34 to 60 inches; silt loam

Oa—Onawa silty clay, 0 to 2 percent slopes

Map Unit Composition

Onawa: 100 percent

Component Descriptions

Onawa

MLRA: 63B - Southern Rolling Pierre Shale Plains

Landform: Swale on flood plain on valley

Parent material: Clayey alluvium over loamy alluvium

Slope: 0 to 2 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: High (About 10.2 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: About 24 to 48 inches

Runoff class: Negligible

Ecological site: Clayey Overflow - Veg. Zone 3

Land capability (irrigated): 2w

Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 7 inches; silty clay

H2—7 to 28 inches; silty clay

H3—28 to 60 inches; stratified silt loam to clay

Minor Components**Albaton**

Slope: 0 to 2 percent
Drainage class: Poorly drained
Ecological site: Clayey Overflow - Veg. Zone 3

Oe—O'Neill fine sandy loam, 0 to 2 percent slopes

Map Unit Composition

O'Neill: 100 percent

Component Descriptions

O'Neill

MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Flat on tableland
Parent material: Coarse-loamy alluvium over sandy and gravelly alluvium
Slope: 0 to 2 percent
Drainage class: Well drained
Slowest permeability: Moderately rapid (About 1.98 in/hr)
Available water capacity: Low (About 5.2 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sandy - Veg. Zone 3
Land capability (irrigated): 3e
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 9 inches; fine sandy loam
 H2—9 to 34 inches; sandy loam
 H3—34 to 60 inches; coarse sand

OeC—O'Neill fine sandy loam, 2 to 6 percent slopes

Map Unit Composition

O'Neill: 100 percent

Component Descriptions

O'Neill

MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Hill on tableland
Hillslope position: Summit, backslope, shoulder
Parent material: Coarse-loamy alluvium over sandy and gravelly alluvium
Slope: 2 to 6 percent
Drainage class: Well drained
Slowest permeability: Moderately rapid (About 1.98 in/hr)
Available water capacity: Low (About 5.2 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sandy - Veg. Zone 3
Land capability (irrigated): 4e
Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 9 inches; fine sandy loam
 H2—9 to 34 inches; fine sandy loam
 H3—34 to 60 inches; gravelly sand

OfD—O'Neill-Meadin fine sandy loams, 3 to 9 percent slopes

Map Unit Composition

O'Neill: 55 percent
 Meadin: 45 percent

Component Descriptions

O'Neill

MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Hill on tableland
Hillslope position: Backslope, summit
Parent material: Coarse-loamy alluvium over sandy and gravelly alluvium
Slope: 3 to 9 percent
Drainage class: Well drained
Slowest permeability: Moderately rapid (About 1.98 in/hr)
Available water capacity: Low (About 5.2 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Sandy - Veg. Zone 3
Land capability (irrigated): 4e
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 9 inches; fine sandy loam
 H2—9 to 34 inches; fine sandy loam
 H3—34 to 60 inches; gravelly sand

Meadin

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Hill on tableland

Hillslope position: Shoulder, summit

Parent material: Loamy alluvium over sandy and gravelly alluvium

Slope: 3 to 9 percent

Drainage class: Excessively drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Low (About 3.6 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Shallow To Gravel - Veg. Zone 3

Land capability (nonirrigated): 6s

Typical Profile:

H1—0 to 8 inches; fine sandy loam
 H2—8 to 13 inches; gravelly sandy loam
 H3—13 to 60 inches; very gravelly coarse sand

On—Onita silt loam, 0 to 2 percent slopes

Map Unit Composition

Onita: 100 percent

Poorly Drained:

Component Descriptions

Onita

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Flat on tableland, swale on tableland

Parent material: Loamy colluvium

Slope: 0 to 2 percent

Drainage class: Well drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: High (About 10.9 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 1

Land capability (nonirrigated): 2c

Typical Profile:

H1—0 to 18 inches; silt loam
 H2—18 to 32 inches; silty clay loam
 H3—32 to 60 inches; silt loam

Minor Components**Scott**

Slope: 0 to 1 percent

Drainage class: Poorly drained

Ecological site: Clayey Overflow - Veg. Zone 3

Component Descriptions

Poorly Drained

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Swale

Depth to seasonal water saturation: More than 6 feet

*Typical Profile:***Or—Ord fine sandy loam, 0 to 2 percent slopes**

Map Unit Composition

Ord: 100 percent

Component Descriptions

Ord

MLRA: 63B - Southern Rolling Pierre Shale Plains

Landform: Interdune on sandhills, stream terrace on valley

Parent material: Stratified sandy alluvium

Slope: 0 to 2 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Moderate (About 6.1 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: About 18 to 36 inches

Runoff class: Very low
Ecological site: Subirrigated - Veg. Zone 3
Land capability (irrigated): 2w
Land capability (nonirrigated): 2w

Typical Profile:
 H1—0 to 11 inches; fine sandy loam
 H2—11 to 30 inches; fine sandy loam
 H3—30 to 60 inches; stratified sand to loamy fine sand

Minor Components

Barney

Slope: 0 to 2 percent
Drainage class: Very poorly drained
Ecological site: Wet Land - Veg. Zone 3

PaC—Paka fine sandy loam, 2 to 6 percent slopes

Map Unit Composition

Paka: 100 percent

Component Descriptions

Paka

MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Hill on tableland
Hillslope position: Backslope, shoulder, summit
Parent material: Loamy residuum weathered from siltstone
Slope: 2 to 6 percent
Depth to restrictive feature: 40 to 60 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: Moderate (About 8.9 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Sandy - Veg. Zone 3
Land capability (irrigated): 3e
Land capability (nonirrigated): 3e

Typical Profile:
 H1—0 to 9 inches; fine sandy loam
 H2—9 to 27 inches; clay loam
 H3—27 to 46 inches; silt loam
 Cr—46 to 60 inches; weathered bedrock

Minor Components

Scott

Slope: 0 to 1 percent
Drainage class: Poorly drained
Ecological site: Clayey Overflow - Veg. Zone 3

Ph—Paka loam, 0 to 2 percent slopes

Map Unit Composition

Paka: 100 percent

Component Descriptions

Paka

MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Flat on tableland
Parent material: Loamy residuum weathered from siltstone
Slope: 0 to 1 percent
Depth to restrictive feature: 40 to 60 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: High (About 9.7 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Silty - Veg. Zone 3
Land capability (irrigated): 1
Land capability (nonirrigated): 2c

Typical Profile:

H1—0 to 8 inches; loam
 H2—8 to 26 inches; silty clay loam
 H3—26 to 48 inches; silt loam
 Cr—48 to 60 inches; weathered bedrock

Minor Components

Scott

Slope: 0 to 1 percent
Drainage class: Poorly drained
Ecological site: Clayey Overflow - Veg. Zone 3

PhC—Paka loam, 2 to 6 percent slopes

Map Unit Composition

Paka: 100 percent

Component Descriptions

Paka

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Hill on tableland

Hillslope position: Summit, shoulder, backslope

Parent material: Loamy residuum weathered from siltstone

Slope: 2 to 6 percent

Depth to restrictive feature: 40 to 60 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 9.7 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 8 inches; loam

H2—8 to 26 inches; silty clay loam

H3—26 to 48 inches; silt loam

Cr—48 to 60 inches; weathered bedrock

PhD—Paka loam, 6 to 11 percent slopes

Map Unit Composition

Paka: 100 percent

Component Descriptions

Paka

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Hill on tableland

Hillslope position: Summit, backslope, shoulder

Parent material: Loamy residuum weathered from siltstone

Slope: 6 to 11 percent

Depth to restrictive feature: 40 to 60 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 9.7 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: High

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 4e

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 8 inches; loam

H2—8 to 26 inches; clay loam

H3—26 to 48 inches; silt loam

Cr—48 to 60 inches; weathered bedrock

PoC—Promise silty clay, 2 to 6 percent slopes

Map Unit Composition

Promise: 100 percent

Component Descriptions

Promise

MLRA: 63B - Southern Rolling Pierre Shale Plains

Landform: Hill on plains, valley side on valley

Parent material: Clayey residuum weathered from shale

Slope: 2 to 6 percent

Drainage class: Well drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: Moderate (About 6.7 inches)

Shrink-swell potential: Very high (About 17.0 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Clayey - Veg. Zone 3

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 18 inches; silty clay
 H2—18 to 33 inches; clay
 H3—33 to 60 inches; clay

RaC—Ree silt loam, 2 to 6 percent slopes

Map Unit Composition

Ree: 100 percent

Component Descriptions

Ree

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Hill on tableland

Hillslope position: Backslope, shoulder, summit

Parent material: Loamy alluvial sediments

Slope: 2 to 6 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 10.0 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 7 inches; silt loam
 H2—7 to 22 inches; clay loam
 H3—22 to 60 inches; sandy loam

RaD—Ree silt loam, 6 to 11 percent slopes

Map Unit Composition

Ree: 100 percent

Component Descriptions

Ree

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Hill on tableland

Hillslope position: Shoulder, backslope, summit

Parent material: Loamy alluvial sediments

Slope: 6 to 11 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 10.0 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: High

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 4e

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 7 inches; silt loam
 H2—7 to 22 inches; silty clay loam
 H3—22 to 60 inches; sandy loam

RaE—Ree silt loam, 11 to 15 percent slopes

Map Unit Composition

Ree: 100 percent

Component Descriptions

Ree

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Hill on tableland

Hillslope position: Summit, backslope, shoulder

Parent material: Loamy alluvial sediments

Slope: 11 to 15 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 10.0 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: High

Ecological site: Silty - Veg. Zone 3

Land capability (nonirrigated): 6e

Typical Profile:

- H1—0 to 7 inches; silt loam
- H2—7 to 22 inches; sandy clay loam
- H3—22 to 60 inches; fine sandy loam

ReC—Reliance silt loam, 2 to 6 percent slopes

Map Unit Composition

Reliance: 100 percent
 Minor components: 100 percent

Component Descriptions

Reliance

MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Loess hill on tableland
Hillslope position: Backslope, summit, shoulder
Parent material: Loess
Slope: 2 to 6 percent

Drainage class: Well drained

Slowest permeability: Slow (About 0.06 in/hr)

Available water capacity: High (About 10.1 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: High

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

- H1—0 to 10 inches; silt loam
- H2—10 to 28 inches; silty clay loam
- H3—28 to 60 inches; silt loam

Minor Components**Scott**

Composition: About 100 percent

Slope: 0 to 1 percent

Drainage class: Poorly drained

Ecological site: Clayey Overflow - Veg. Zone 3

ReD—Reliance silt loam, 6 to 11 percent slopes

Map Unit Composition

Reliance: 100 percent

Component Descriptions

Reliance

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Loess hill on tableland

Hillslope position: Summit, shoulder, backslope

Parent material: Loess

Slope: 6 to 11 percent

Drainage class: Well drained

Slowest permeability: Slow (About 0.06 in/hr)

Available water capacity: High (About 10.1 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 4e

Land capability (nonirrigated): 4e

Typical Profile:

- H1—0 to 10 inches; silt loam
- H2—10 to 28 inches; silty clay loam
- H3—28 to 60 inches; silt loam

RfC—Reliance silty clay loam, 2 to 6 percent slopes

Map Unit Composition

Reliance: 100 percent

Component Descriptions

Reliance

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Loess hill on tableland

Hillslope position: Summit, shoulder, backslope

Parent material: Loess

Slope: 2 to 6 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 10.1 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 3e
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 10 inches; silty clay loam
 H2—10 to 28 inches; silty clay loam
 H3—28 to 60 inches; silty clay loam

Minor Components

Scott

Slope: 0 to 1 percent
Drainage class: Poorly drained
Ecological site: Clayey Overflow - Veg. Zone 3

Rw—Riverwash

Map Unit Composition

Riverwash: 100 percent

Component Descriptions

Riverwash

MLRA: 63B - Southern Rolling Pierre Shale
 Plains

Landform: Flood plain on valley
Parent material: Sandy and gravelly alluvium
Slope: 0 to 1 percent
Drainage class: Very poorly drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Very low (About 2.1 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Frequent
Depth to seasonal water saturation: About 0 to 0 inches
Runoff class: Negligible
Land capability (nonirrigated): 8w

Typical Profile:

H1—0 to 6 inches; fine sand
 H2—6 to 60 inches; stratified sand to very gravelly coarse sand

SaG—Sansarc silty clay, 20 to 40 percent slopes

Map Unit Composition

Sansarc: 100 percent

Component Descriptions

Sansarc

MLRA: 63B - Southern Rolling Pierre Shale
 Plains

Landform: Hill on hills, valley side on valley
Parent material: Clayey residuum weathered from shale
Slope: 30 to 40 percent
Depth to restrictive feature: 4 to 20 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Slow (About 0.06 in/hr)
Available water capacity: Very low (About 1.1 inches)
Shrink-swell potential: Very high (About 17.0 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very high
Ecological site: Shallow Clay - Veg. Zone 3
Land capability (nonirrigated): 7s

Typical Profile:

H1—0 to 16 inches; silty clay
 Cr—16 to 60 inches; weathered bedrock

Sc—Scott silt loam, 0 to 1 percent slopes

Map Unit Composition

Scott: 100 percent

Component Descriptions

Scott

MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Playa on tableland
Parent material: Loess
Slope: 0 to 1 percent
Drainage class: Poorly drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: Moderate (About 8.7 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: About 0 to 0 inches
Runoff class: Negligible

Ecological site: Clayey Overflow - Veg. Zone 3
Land capability (nonirrigated): 4w

Typical Profile:

H1—0 to 8 inches; silt loam
 H2—8 to 48 inches; silty clay
 H3—48 to 56 inches; silty clay loam
 H4—56 to 64 inches; clay loam

Sm—Simeon loamy sand, 0 to 2 percent slopes

Map Unit Composition

Simeon: 100 percent

Component Descriptions

Simeon

MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Flat on tableland
Parent material: Sandy and gravelly alluvium
Slope: 0 to 2 percent
Drainage class: Excessively drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 4.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Negligible
Ecological site: Shallow To Gravel - Veg. Zone 3
Land capability (irrigated): 4s
Land capability (nonirrigated): 6s

Typical Profile:

H1—0 to 5 inches; loamy sand
 H2—5 to 60 inches; sand

SuC—Simeon-Valentine loamy sands, 0 to 6 percent slopes

Map Unit Composition

Simeon: 55 percent
 Valentine: 45 percent

Component Descriptions

Simeon

MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Swale, upland
Parent material: Sandy and gravelly alluvium
Slope: 0 to 6 percent
Drainage class: Excessively drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 5.2 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Negligible
Ecological site: Shallow To Gravel - Veg. Zone 3
Land capability (irrigated): 4s
Land capability (nonirrigated): 6s

Typical Profile:

H1—0 to 15 inches; loamy sand
 H2—15 to 60 inches; coarse sand

Valentine

MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Dune, upland
Parent material: Eolian sands
Slope: 0 to 6 percent
Drainage class: Excessively drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 4.0 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Negligible
Ecological site: Sands - Veg. Zone 3
Land capability (irrigated): 4e
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 5 inches; loamy sand
 H2—5 to 11 inches; fine sand
 H3—11 to 60 inches; fine sand

SvF2—Simeon-Valentine complex, 3 to 30 percent slopes, Eroded

Map Unit Composition

Simeon: 70 percent
 Valentine: 30 percent

Component Descriptions

Simeon

MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Hill on upland, valley side on valley
Hillslope position: Summit, shoulder

Parent material: Sandy and gravelly alluvium
Slope: 3 to 30 percent
Drainage class: Excessively drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 5.2 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Shallow To Gravel - Veg. Zone 3
Land capability (nonirrigated): 6s

Typical Profile:
 H1—0 to 15 inches; loamy sand
 H2—15 to 60 inches; coarse sand

Valentine
MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Hill on tableland, valley side on valley
Hillslope position: Backslope
Parent material: Eolian sands
Slope: 3 to 30 percent
Drainage class: Excessively drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 3.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Sands - Veg. Zone 3
Land capability (nonirrigated): 6e

Typical Profile:
 H1—0 to 5 inches; fine sand
 H2—5 to 11 inches; fine sand
 H3—11 to 60 inches; fine sand

VaE—Valentine fine sand, 6 to 17 percent slopes

Map Unit Composition

Valentine: 100 percent

Component Descriptions
Valentine
MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Dune on sandhills
Parent material: Eolian sands
Slope: 6 to 17 percent
Drainage class: Excessively drained

Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 3.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sands - Veg. Zone 3
Land capability (nonirrigated): 6e

Typical Profile:
 H1—0 to 5 inches; fine sand
 H2—5 to 11 inches; fine sand
 H3—11 to 60 inches; fine sand

VbB—Valentine loamy sand, 0 to 3 percent slopes

Map Unit Composition

Valentine: 100 percent

Component Descriptions
Valentine
MLRA: 66 - Dakota-Nebraska Eroded Tableland
Landform: Hummock on upland
Parent material: Eolian sands
Slope: 0 to 3 percent
Drainage class: Excessively drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 4.0 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Negligible
Ecological site: Sandy - Veg. Zone 3
Land capability (irrigated): 4e
Land capability (nonirrigated): 4e

Typical Profile:
 H1—0 to 5 inches; loamy sand
 H2—5 to 11 inches; loamy sand
 H3—11 to 60 inches; fine sand

Ve—Verdel silty clay, 0 to 2 percent slopes

Map Unit Composition

Verdel: 100 percent

Component Descriptions

Verdel

MLRA: 63B - Southern Rolling Pierre Shale
Plains

Landform: Stream terrace on valley

Parent material: Clayey alluvium derived from shale

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: Moderate (About 7.4 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: High

Ecological site: Clayey - Veg. Zone 3

Land capability (irrigated): 2s

Land capability (nonirrigated): 2s

Typical Profile:

H1—0 to 14 inches; silty clay

H2—14 to 60 inches; silty clay

Hillslope position: Shoulder, backslope, summit

Parent material: Loamy eolian deposits over clayey residuum weathered from clayey shale

Slope: 2 to 6 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: Low (About 3.2 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Sandy - Veg. Zone 3

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 8 inches; fine sandy loam

H2—8 to 15 inches; sandy clay loam

H3—15 to 22 inches; clay

Cr—22 to 60 inches; weathered bedrock

WeC—Wewela fine sandy loam, 2 to 6 percent slopes

Map Unit Composition

Wewela: 100 percent

Component Descriptions

Wewela

MLRA: 63B - Southern Rolling Pierre Shale
Plains

Landform: Hill on plains

zwa—Water > 40 Acres

Map Unit Composition

Water: 100 percent

Component Descriptions

Water

MLRA: 63B - Southern Rolling Pierre Shale
Plains

Depth to seasonal water saturation: More than 6 feet

Land capability classification shows, in a general way, the suitability of soils for most kinds of field crops. Crops that require special management are excluded. The soils are grouped according to their limitations for field crops, the risk of damage if they are used for crops, and the way they respond to management. The criteria used in grouping the soils do not include major and generally expensive land-forming that would change slope, depth, or other characteristics of the soils, nor do they include possible but unlikely major reclamation projects. Capability classification is not a substitute for interpretations designed to show suitability and limitations of groups of soils for rangeland, for forestland, or for engineering purposes. In the capability system, soils are generally grouped at three levels: capability class, subclass, and unit.

Capability classes, the broadest groups, are designated by the numbers 1 through 8. The numbers indicate progressively greater limitations and narrower choices for practical use. The classes are defined as follows:

(Class 1) soils have slight limitations that restrict their use.

(Class 2) soils have moderate limitations that restrict the choice of plants or that require moderate conservation practices.

(Class 3) soils have severe limitations that restrict the choice of plants or that require special conservation practices, or both.

(Class 4) soils have very severe limitations that restrict the choice of plants or that require very careful management, or both.

(Class 5) soils are subject to little or no erosion but have other limitations, impractical to remove, that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

(Class 6) soils have severe limitations that make them generally unsuitable for cultivation and that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

(Class 7) soils have very severe limitations that make them unsuitable for cultivation and that restrict their use mainly to grazing, forestland, or wildlife habitat.

(Class 8) soils and miscellaneous areas have limitations that preclude commercial plant production and that restrict their use to recreational purposes, wildlife habitat, watershed, or esthetic purposes.

Capability subclasses are soil groups within one class. They are designated by adding a small letter, e, w, s, or c, to the class numeral, for example, 2e. The letter e shows that the main hazard is the risk of erosion unless close-growing plant cover is maintained; w shows that water in or on the soil interferes with plant growth or cultivation (in some soils the wetness can be partly corrected by artificial drainage); s shows that the soil is limited mainly because it is shallow, droughty, or stony; and c, used in only some parts of the United States, shows that the chief limitation is climate that is very cold or very dry.

In class 1 there are no subclasses because the soils of this class have few limitations. Class 5 contains only the subclasses indicated by w, s, or c because the soils in class 5 are subject to little or no erosion. They have other limitations that restrict their use to pasture, rangeland, forestland, wildlife habitat, or recreation.

Capability units are soil groups within a subclass. The soils in a capability unit are enough alike to be suited to the same crops and pasture plants, to require similar management, and to have similar productivity. Capability units are generally designated by adding an Arabic numeral to the subclass symbol, for example, 2e-4 and 3e-6. These units are not given in all soil surveys.

The capability classification of map units in this survey area is given in the section "Detailed Soil Map Units" and in the Land Capability and Component Yields table.

Crop Yield Estimates

The average yields per acre that can be expected of the principal crops under a high level of management are shown in "Land Capability and Component Yields" table. In any given year, yields may be higher or lower than those indicated in the table because of variations in rainfall and other climatic factors. The land capability classification of map units in the survey area also is shown in the table.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, animal waste manure, and green manure crops; and harvesting that ensures the smallest possible loss.

For yields of irrigated crops, it is assumed that the irrigation system is adapted to the soils and to the crops grown, that good-quality irrigation water is uniformly applied as needed, and that tillage is kept to a minimum.

The estimated yields reflect the productive capacity of each soil for each of the principal crops. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Crops other than those shown in this table, are grown in the survey area, but estimated yields are not listed because the acreage of such crops is small. The local office of the Natural Resources Conservation Service (NRCS) or the Cooperative Extension Service (CES) can provide information about the management and productivity of the soils for those crops.

LAND CAPABILITY AND YIELDS PER ACRE OF CROPS--Continued
Boyd County, Nebraska

PAGE 2 of 4

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)

Map symbol and soil name	Land Capability		Corn		Grain sorghum		Oats	
	N	I	N	I	N	I	N	I
	Bu		Bu		Bu		Bu	
Ab: ALBATON-----	3w	3w	80.00	116.00	65.00	100.00	55.00	---
AnC: ANSELMO-----	3e	3e	41.00	137.00	45.00	110.00	44.00	---
AnD: ANSELMO-----	4e	4e	35.00	---	40.00	---	39.00	---
AnF: ANSELMO-----	6e	6e	30.00	95.00	30.00	---	33.00	---
ArF: ANSELMO-----	6e	6e	25.00	84.00	25.00	---	28.00	---
ROCK OUTCROP----	8s	---	---	---	25.00	---	28.00	---
Ba: BARNEY-----	5w	---	---	---	---	---	---	---
Bd: BLAKE-----	2w	2w	85.00	147.00	80.00	120.00	72.00	---
Be: BLENDON-----	2e	2e	50.00	152.00	55.00	120.00	50.00	75.00
BeC: BLENDON-----	3e	3e	45.00	147.00	50.00	110.00	46.00	---
BoD: BOYD-----	4e	---	26.00	---	40.00	---	33.00	---
BrG: BRISTOW-----	7s	---	---	---	---	---	---	---
Bs: BROCKSBURG-----	2e	2e	30.00	137.00	40.00	110.00	33.00	---
Bt: BROCKSBURG-----	2s	2s	40.00	142.00	45.00	115.00	39.00	---
Cb: CASS-----	2w	2w	45.00	142.00	55.00	115.00	44.00	---
CrE2: CROFTON-----	4e	---	35.00	---	40.00	---	33.00	---
DuB: DUNDAY-----	4e	3e	30.00	131.00	35.00	100.00	33.00	---
DuC: DUNDAY-----	4e	4e	24.00	126.00	25.00	95.00	28.00	---
DuD: DUNDAY-----	6e	4e	20.00	84.00	20.00	70.00	22.00	---
DxB: DUNDAY-----	4e	3e	35.00	137.00	40.00	110.00	39.00	---
Et: ELTREE-----	2c	1	60.00	152.00	70.00	120.00	66.00	80.00
Go: GRIGSTON-----	2w	2w	65.00	152.00	75.00	120.00	72.00	---
GrB: AOWA-----	6w	---	---	---	---	---	---	---
Ha: HALL-----	2c	1	60.00	152.00	70.00	120.00	66.00	80.00
He: HAYNIE-----	1	1	80.00	147.00	80.00	120.00	72.00	---
IfD: INAFALE-----	6e	4e	---	---	---	---	---	---
IgB: INAFALE-----	6w	---	---	---	---	---	---	---

LAND CAPABILITY AND YIELDS PER ACRE OF CROPS--Continued

Boyd County, Nebraska

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)

Map symbol and soil name	Land Capability		Corn		Grain sorghum		Oats	
	N	I	N	I	N	I	N	I
	Bu		Bu		Bu		Bu	
IhB: INAVALE-----	4e	3e	30.00	116.00	35.00	100.00	33.00	---
In: INAVALE-----	3e	3e	35.00	121.00	40.00	110.00	39.00	---
Jn: JANSEN-----	2s	2s	35.00	142.00	40.00	110.00	33.00	---
JnC: JANSEN-----	3e	3e	30.00	131.00	40.00	110.00	33.00	---
JnD: JANSEN-----	4e	4e	30.00	---	31.00	---	28.00	---
LaD: LABU-----	4e	4e	24.00	---	38.00	---	28.00	---
LcF: LABU-----	6e	---	---	---	---	---	---	---
SANSARC-----	6s	---	---	---	---	---	---	---
Le: LESHARA-----	2w	2w	60.00	152.00	75.00	120.00	55.00	---
LsC: LYNCH-----	3e	3e	32.00	---	40.00	---	40.00	---
LsD: LYNCH-----	4e	4e	25.00	---	30.00	---	31.00	---
LyD: LYNCH-----	4e	4e	25.00	---	30.00	---	31.00	---
BRISTOW-----	6s	---	25.00	---	30.00	---	31.00	---
LyF: LYNCH-----	6e	---	---	---	---	---	---	---
BRISTOW-----	6s	---	---	---	---	---	---	---
MaG: MARIAVILLE-----	6s	---	---	---	---	---	---	---
PAKA-----	6e	---	---	---	---	---	---	---
MeE: MEADIN-----	6s	---	---	---	---	---	---	---
NoC: NORA-----	2e	3e	55.00	137.00	65.00	110.00	55.00	---
NoD: NORA-----	3e	4e	50.00	---	60.00	---	50.00	---
Oa: ONAWA-----	2w	2w	80.00	116.00	75.00	110.00	61.00	---
Oe: O'NEILL-----	3e	3e	30.00	137.00	35.00	105.00	33.00	---
OeC: O'NEILL-----	4e	4e	25.00	116.00	30.00	100.00	31.00	---
OfD: O'NEILL-----	6e	4e	20.00	105.00	25.00	80.00	28.00	---
MEADIN-----	6s	---	---	---	25.00	80.00	28.00	---
On: ONITA-----	2c	1	55.00	152.00	70.00	125.00	66.00	80.00
POORLY DRAINED--	---	---	---	---	---	---	---	---
Or: ORD-----	2w	2w	45.00	121.00	50.00	115.00	44.00	---
PaC: PAKA-----	3e	3e	45.00	142.00	50.00	110.00	55.00	---

LAND CAPABILITY AND YIELDS PER ACRE OF CROPS--Continued
Boyd County, Nebraska

PAGE 4 of 4

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)

Map symbol and soil name	Land Capability		Corn		Grain sorghum		Oats	
	N	I	N	I	N	I	N	I
			Bu		Bu		Bu	
Ph: PAKA-----	2c	1	55.00	152.00	70.00	125.00	66.00	80.00
PhC: PAKA-----	3e	3e	50.00	137.00	65.00	110.00	55.00	---
PhD: PAKA-----	4e	4e	40.00	---	45.00	---	50.00	---
PoC: PROMISE-----	3e	---	35.00	---	45.00	---	39.00	---
RaC: REE-----	3e	3e	47.00	137.00	60.00	110.00	52.00	---
RaD: REE-----	4e	4e	40.00	---	55.00	---	44.00	---
RaE: REE-----	6e	---	28.00	---	40.00	---	33.00	---
ReC: RELIANCE-----	3e	3e	50.00	137.00	65.00	115.00	55.00	---
ReD: RELIANCE-----	4e	4e	45.00	---	58.00	---	47.00	---
RfC: RELIANCE-----	3e	3e	45.00	131.00	60.00	110.00	50.00	---
Rw: RIVERWASH-----	8w	---	---	---	---	---	---	---
SaG: SANSARC-----	7s	---	---	---	---	---	---	---
Sc: SCOTT-----	4w	---	---	---	---	---	---	---
Sm: SIMEON-----	6s	4s	---	84.00	---	---	---	---
SuC: SIMEON-----	6s	4s	---	95.00	---	---	---	---
VALENTINE-----	6e	4e	---	95.00	---	---	---	---
SvF2: SIMEON-----	6s	---	---	---	---	---	---	---
VALENTINE-----	6e	---	---	---	---	---	---	---
VaE: VALENTINE-----	6e	---	---	---	---	---	---	---
VbB: VALENTINE-----	4e	4e	---	105.00	---	---	---	---
Ve: VERDEL-----	2s	2s	45.00	121.00	55.00	110.00	50.00	---
WeC: WEWELA-----	3e	3e	35.00	---	45.00	---	39.00	---
zwa: WATER-----	---	---	---	---	---	---	---	---

Farmland Classification
Boyd County, Nebraska : Published

Prime farmland is one of several kinds of important farmland defined by the U.S. Department of Agriculture. It is of major importance in meeting the Nation's short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation's prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil qualities, growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. It is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local office of the Natural Resources Conservation Service.

A recent trend in land use in some parts of the survey area has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

The map units in the survey area that are considered prime farmland are listed in the following table. This list does not constitute a recommendation for a particular land use. On some soils included in the list, measures that overcome a hazard or limitation, such as flooding, wetness, and droughtiness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures. The extent of each listed map unit is shown in the "Acres and Proportionate Extent of Soils" table. The location is shown on the detailed soil maps. The soil qualities that affect use and management are described in other tables in this document."

Map symbol	Mapunit name	Farmland Classification
AnC	Anselmo fine sandy loam, 2 to 6 percent slopes	All areas are prime farmland
Be	Blendon fine sandy loam, 0 to 2 percent slopes	All areas are prime farmland
BeC	Blendon fine sandy loam, 2 to 6 percent slopes	All areas are prime farmland
Bs	Brocksburg fine sandy loam, 0 to 2 percent slopes	All areas are prime farmland
Bt	Brocksburg loam, 0 to 2 percent slopes	All areas are prime farmland
Cb	Cass fine sandy loam, 0 to 2 percent slopes	All areas are prime farmland
Et	Eltree silt loam, 0 to 2 percent slopes	All areas are prime farmland
Go	Grigston silt loam, 0 to 2 percent slopes	All areas are prime farmland
Ha	Hall silt loam, 0 to 2 percent slopes	All areas are prime farmland
He	Haynie silt loam, 0 to 2 percent slopes	All areas are prime farmland
Jn	Jansen loam, 0 to 2 percent slopes	All areas are prime farmland
JnC	Jansen loam, 2 to 6 percent slopes	All areas are prime farmland
NoC	Nora silt loam, 2 to 6 percent slopes	All areas are prime farmland
On	Onita silt loam, 0 to 2 percent slopes	All areas are prime farmland
PaC	Paka fine sandy loam, 2 to 6 percent slopes	All areas are prime farmland
Ph	Paka loam, 0 to 2 percent slopes	All areas are prime farmland
PhC	Paka loam, 2 to 6 percent slopes	All areas are prime farmland
ReC	Reliance silt loam, 2 to 6 percent slopes	All areas are prime farmland
RfC	Reliance silty clay loam, 2 to 6 percent slopes	All areas are prime farmland
Ve	Verdel silty clay, 0 to 2 percent slopes	All areas are prime farmland
Bd	Blake silty clay loam, 0 to 2 percent slopes	Prime farmland if drained
Le	Leshara silt loam, 0 to 2 percent slopes	Prime farmland if drained
Oa	Onawa silty clay, 0 to 2 percent slopes	Prime farmland if drained
Or	Ord fine sandy loam, 0 to 2 percent slopes	Prime farmland if drained

The "Soil Rating for Plant Growth, modified 1998" (SRPG) is a relative rating of the capacity of a soil to produce a specific plant under a defined management system. The index is determined from yield data on a few benchmark soils and is used to calculate yields, the net returns from crops, land assessment values, and taxes and to perform risk analysis when land management decisions are made. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

Map symbol	Soil name	Crop Index
Ab	Albaton Silty Clay, 0 To 2 Percent Slopes-----	31
AnC	Anselmo Fine Sandy Loam, 2 To 6 Percent Slopes-----	50
AnD	Anselmo Fine Sandy Loam, 6 To 11 Percent Slopes-----	46
AnF	Anselmo Fine Sandy Loam, 11 To 20 Percent Slopes-----	41
ArF	Anselmo-Rock Outcrop Complex, 11 To 20 Percent Slopes-----	31
Ba	Barney Silt Loam, 0 To 2 Percent Slopes-----	23
Bd	Blake Silty Clay Loam, 0 To 2 Percent Slopes-----	34
Be	Blendon Fine Sandy Loam, 0 To 2 Percent Slopes-----	49
BeC	Blendon Fine Sandy Loam, 2 To 6 Percent Slopes-----	47
BoD	Boyd Silty Clay, 6 To 11 Percent Slopes-----	20
BrG	Bristow Silty Clay, 20 To 40 Percent Slopes-----	1
Bs	Brocksburg Fine Sandy Loam, 0 To 2 Percent Slopes-----	54
Bt	Brocksburg Loam, 0 To 2 Percent Slopes-----	59
Cb	Cass Fine Sandy Loam, 0 To 2 Percent Slopes-----	45
CrE2	Crofton Silt Loam, 11 To 15 Percent Slopes, Eroded-----	44
DuB	Dunday Loamy Fine Sand, 0 To 3 Percent Slopes-----	31
DuC	Dunday Loamy Fine Sand, 3 To 6 Percent Slopes-----	30
DuD	Dunday Loamy Fine Sand, 6 To 11 Percent Slopes-----	28
DxB	Dunday Loamy Fine Sand, Loamy Substratum, 0 To 3 Percent Slopes-----	37
Et	Eltree Silt Loam, 0 To 2 Percent Slopes-----	61
Go	Grigston Silt Loam, 0 To 2 Percent Slopes-----	56
GrB	Grigston Silt Loam, Channeled, 0 To 3 Percent Slopes-----	44
Ha	Hall Silt Loam, 0 To 2 Percent Slopes-----	70
He	Haynie Silt Loam, 0 To 2 Percent Slopes-----	42
IfD	Inavale Fine Sand, 3 To 11 Percent Slopes-----	28
IgB	Inavale Fine Sand, Channeled, 0 To 3 Percent Slopes-----	23
IhB	Inavale Loamy Fine Sand, 0 To 3 Percent Slopes-----	32
In	Inavale Fine Sandy Loam, 0 To 2 Percent Slopes-----	37
Jn	Jansen Loam, 0 To 2 Percent Slopes-----	58
JnC	Jansen Loam, 2 To 6 Percent Slopes-----	56
JnD	Jansen Loam, 6 To 11 Percent Slopes-----	51
LaD	Labu Silty Clay, 6 To 11 Percent Slopes-----	22
LcF	Labu-Sansarc Silty Clays, 11 To 30 Percent Slopes-----	9
Le	Leshara Silt Loam, 0 To 2 Percent Slopes-----	61
LsC	Lynch Silty Clay, 2 To 6 Percent Slopes-----	11
LsD	Lynch Silty Clay, 6 To 11 Percent Slopes-----	10
LyD	Lynch-Bristow Silty Clays, 6 To 11 Percent Slopes-----	9
LyF	Lynch-Bristow Silty Clays, 11 To 30 Percent Slopes-----	5
MaG	Mariaville-Paka Loams, 15 To 40 Percent Slopes-----	8
MeE	Meadin Sandy Loam, 3 To 17 Percent Slopes-----	26
NoC	Nora Silt Loam, 2 To 6 Percent Slopes-----	66
NoD	Nora Silt Loam, 6 To 11 Percent Slopes-----	61
Oa	Onawa Silty Clay, 0 To 2 Percent Slopes-----	33
Oe	O'Neill Fine Sandy Loam, 0 To 2 Percent Slopes-----	44
OeC	O'Neill Fine Sandy Loam, 2 To 6 Percent Slopes-----	42
OfD	O'Neill-Meadin Fine Sandy Loams, 3 To 9 Percent Slopes-----	35
On	Onita Silt Loam, 0 To 2 Percent Slopes-----	65
Or	Ord Fine Sandy Loam, 0 To 2 Percent Slopes-----	26
PaC	Paka Fine Sandy Loam, 2 To 6 Percent Slopes-----	63
Ph	Paka Loam, 0 To 2 Percent Slopes-----	66
PhC	Paka Loam, 2 To 6 Percent Slopes-----	64
PhD	Paka Loam, 6 To 11 Percent Slopes-----	58
PoC	Promise Silty Clay, 2 To 6 Percent Slopes-----	33
RaC	Ree Silt Loam, 2 To 6 Percent Slopes-----	61
RaD	Ree Silt Loam, 6 To 11 Percent Slopes-----	56
RaE	Ree Silt Loam, 11 To 15 Percent Slopes-----	51
ReC	Reliance Silt Loam, 2 To 6 Percent Slopes-----	37
ReD	Reliance Silt Loam, 6 To 11 Percent Slopes-----	55
RfC	Reliance Silty Clay Loam, 2 To 6 Percent Slopes-----	60
Rw	Riverwash-----	0
SaG	Sansarc Silty Clay, 20 To 40 Percent Slopes-----	0
Sc	Scott Silt Loam, 0 To 1 Percent Slopes-----	14
Sm	Simeon Loamy Sand, 0 To 2 Percent Slopes-----	27
SuC	Simeon-Valentine Loamy Sands, 0 To 6 Percent Slopes-----	27
SvF2	Simeon-Valentine Complex, 3 To 30 Percent Slopes, Eroded-----	20
VaE	Valentine Fine Sand, 6 To 17 Percent Slopes-----	22
VbB	Valentine Loamy Sand, 0 To 3 Percent Slopes-----	27
Ve	Verdel Silty Clay, 0 To 2 Percent Slopes-----	46
WeC	Wewela Fine Sandy Loam, 2 To 6 Percent Slopes-----	30
zwa	Water > 40 Acres-----	0

Boyd County, Nebraska: Published
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Ab:ALBATON-----	100	3w-	3w	Not prime farmland	D	Clayey Overflow - Veg. Zone 3		.28	.28	5	4	86
AnC:ANSELMO-----	100	3e-	3e	All areas are prime farmland	B	Sandy - Veg. Zone 3		.20	.20	5	3	86
AnD:ANSELMO-----	100	4e-	4e	Not prime farmland	B	Sandy - Veg. Zone 3		.20	.20	5	3	86
AnF:ANSELMO-----	100	6e-	6e	Not prime farmland	B	Sandy - Veg. Zone 3		.20	.20	5	3	86
ArF:ANSELMO-----	75	6e-	6e	Not prime farmland	B	Sandy - Veg. Zone 3		.20	.20	5	3	86
ArF:ROCK OUTCROP	25	N/A	8s	Not prime farmland	D	No Site - Veg. Zone 3		---	---	-	8	0
Ba:BARNEY-----	100	N/A	5w	Not prime farmland	D	Wet Land - Veg. Zone 3		.32	.32	5	8	0
Bd:BLAKE-----	100	2w-	2w	Prime farmland if drained	B	Silty Lowland - Veg. Zone 3		.32	.32	5	4L	86
Be:BLENDON-----	100	2e-	2e	All areas are prime farmland	B	Sandy - Veg. Zone 3		.20	.20	5	3	86
BeC:BLENDON-----	100	3e-	3e	All areas are prime farmland	B	Sandy - Veg. Zone 3		.20	.20	5	3	86
BoD:BOYD-----	100	N/A	4e	Not prime farmland	D	Clayey - Veg. Zone 3		.37	.37	3	4	86
BrG:BRISTOW-----	100	N/A	7s	Not prime farmland	D	Shallow Limy - Veg. Zone 3		.43	.43	2	4	86
Bs:BROCKSBURG---	100	2e-	2e	All areas are prime farmland	B	Sandy - Veg. Zone 3		.20	.20	4	3	86
Bt:BROCKSBURG---	100	2s-	2s	All areas are prime farmland	B	Silty - Veg. Zone 3		.28	.28	4	5	56
Cb:CASS-----	100	2w-	2w	All areas are prime farmland	B	Sandy Lowland - Veg. Zone 3		.20	.20	5	3	86
CrE2:CROFTON----	100	N/A	4e	Not prime farmland	B	Limy Upland - Veg. Zone 3		.43	.43	5	4L	86
DuB:DUNDAY-----	100	3e-	4e	Not prime farmland	A	Sandy - Veg. Zone 3		.15	.15	5	2	134
DuC:DUNDAY-----	100	4e-	4e	Not prime farmland	A	Sandy - Veg. Zone 3		.17	.17	5	2	134

Boyd County, Nebraska: Published
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
DuD:DUNDAY-----	100	4e-	6e	Not prime farmland	A	Sandy - Veg. Zone 3		.17	.17	5	2	134
DxB:DUNDAY-----	100	3e-	4e	Not prime farmland	A	Sandy - Veg. Zone 3		.17	.17	5	2	134
Et:ELTREE-----	100	1-	2c	All areas are prime farmland	B	Silty - Veg. Zone 3		.32	.32	5	6	48
Go:GRIGSTON-----	100	2w-	2w	All areas are prime farmland	B	Silty Lowland - Veg. Zone 3		.32	.32	5	6	48
GrB:AOWA-----	100	N/A	6w	Not prime farmland	B	Silty Overflow - Veg. Zone 3		.32	.32	5	4L	86
Ha:HALL-----	100	1-	2c	All areas are prime farmland	B	Silty Lowland - Veg. Zone 3		.32	.32	5	6	48
He:HAYNIE-----	100	1-	1	All areas are prime farmland	B	Silty Lowland - Veg. Zone 3		.32	.32	5	4L	86
IfD:INAVALE-----	100	4e-	6e	Not prime farmland	A	Sands - Veg. Zone 3		.15	.15	5	1	220
IgB:INAVALE-----	100	N/A	6w	Not prime farmland	A	Sandy Lowland - Veg. Zone 3		.15	.15	5	1	220
IhB:INAVALE-----	100	3e-	4e	Not prime farmland	A	Sandy Lowland - Veg. Zone 3		.17	.17	5	2	134
In:INAVALE-----	100	3e-	3e	Not prime farmland	A	Sandy Lowland - Veg. Zone 3		.24	.24	5	3	86
Jn:JANSEN-----	100	2s-	2s	All areas are prime farmland	B	Silty - Veg. Zone 3		.28	.28	4	5	56
JnC:JANSEN-----	100	3e-	3e	All areas are prime farmland	B	Silty - Veg. Zone 3		.28	.28	4	5	56
JnD:JANSEN-----	100	4e-	4e	Not prime farmland	B	Silty - Veg. Zone 3		.28	.28	4	5	56
LaD:LABU-----	100	4e-	4e	Not prime farmland	D	Clayey - Veg. Zone 3		.32	.32	3	4	86
LcF:LABU-----	65	N/A	6e	Not prime farmland	D	Clayey - Veg. Zone 3		.32	.32	3	4	86
LcF:SANSARC-----	35	N/A	6s	Not prime farmland	D	Shallow Clay - Veg. Zone 3		.37	.37	2	4	86
Le:LESHARA-----	100	2w-	2w	Prime farmland if drained	B	Subirrigated - Veg. Zone 3		.32	.32	5	6	48

Boyd County, Nebraska: Published
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
LsC:LYNCH-----	100	3e-	3e	Not prime farmland	D	Limy Upland - Veg. Zone 3		.32	.32	3	4	86
LsD:LYNCH-----	100	4e-	4e	Not prime farmland	D	Limy Upland - Veg. Zone 3		.32	.32	3	4	86
LyD:LYNCH-----	65	4e-	4e	Not prime farmland	D	Limy Upland - Veg. Zone 3		.32	.32	3	4	86
LyD:BRISTOW----	35	N/A	6s	Not prime farmland	D	Shallow Limy - Veg. Zone 3		.43	.43	2	4	86
LyF:LYNCH-----	55	N/A	6e	Not prime farmland	D	Limy Upland - Veg. Zone 3		.32	.32	3	4	86
LyF:BRISTOW----	45	N/A	6s	Not prime farmland	D	Shallow Limy - Veg. Zone 3		.43	.43	2	4	86
MaG:MARIAVILLE--	55	N/A	6s	Not prime farmland	D	Shallow Limy - Veg. Zone 3		.37	.37	2	4L	86
MaG:PAKA-----	45	N/A	6e	Not prime farmland	B	Silty - Veg. Zone 3		.28	.28	4	5	56
MeE:MEADIN-----	100	N/A	6s	Not prime farmland	A	Shallow To Gravel - Veg. Zone 3		.20	.20	5	3	86
NoC:NORA-----	100	3e-	2e	All areas are prime farmland	B	Silty - Veg. Zone 3		.32	.32	5	6	48
NoD:NORA-----	100	4e-	3e	Not prime farmland	B	Silty - Veg. Zone 3		.32	.32	5	6	48
Oa:ONAWA-----	100	2w-	2w	Prime farmland if drained	D	Clayey Overflow - Veg. Zone 3		.32	.32	5	4	86
Oe:O'NEILL-----	100	3e-	3e	Not prime farmland	B	Sandy - Veg. Zone 3		.20	.20	4	3	86
OeC:O'NEILL-----	100	4e-	4e	Not prime farmland	B	Sandy - Veg. Zone 3		.20	.20	4	3	86
OfD:O'NEILL-----	55	4e-	6e	Not prime farmland	B	Sandy - Veg. Zone 3		.20	.20	4	3	86
OfD:MEADIN-----	45	N/A	6s	Not prime farmland	A	Shallow To Gravel - Veg. Zone 3		.20	.20	5	3	86
On:ONITA-----	100	1-	2c	All areas are prime farmland	C	Silty - Veg. Zone 3		.28	.28	5	6	48
On:POORLY DRAINED-----		N/A	N/A	All areas are prime farmland		Unspecified		---	---	-	---	---

Boyd County, Nebraska: Published
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Or:ORD-----	100	2w-	2w	Prime farmland if drained	B	Subirrigated - Veg. Zone 3		.20	.20	4	3	86
PaC:PAKA-----	100	3e-	3e	All areas are prime farmland	B	Sandy - Veg. Zone 3		.20	.20	4	3	86
Ph:PAKA-----	100	1-	2c	All areas are prime farmland	B	Silty - Veg. Zone 3		.28	.28	4	5	56
PhC:PAKA-----	100	3e-	3e	All areas are prime farmland	B	Silty - Veg. Zone 3		.28	.28	4	5	56
PhD:PAKA-----	100	4e-	4e	Not prime farmland	B	Silty - Veg. Zone 3		.28	.28	4	5	56
PoC:PROMISE----	100	N/A	3e	Not prime farmland	D	Clayey - Veg. Zone 3		.37	.37	5	4	86
RaC:REE-----	100	3e-	3e	Not prime farmland	B	Silty - Veg. Zone 3		.32	.32	5	6	48
RaD:REE-----	100	4e-	4e	Not prime farmland	B	Silty - Veg. Zone 3		.32	.32	5	6	48
RaE:REE-----	100	N/A	6e	Not prime farmland	B	Silty - Veg. Zone 3		.32	.32	5	6	48
ReC:RELIANCE----	100	3e-	3e	All areas are prime farmland	C	Silty - Veg. Zone 3		.32	.32	4	6	48
ReD:RELIANCE----	100	4e-	4e	Not prime farmland	C	Silty - Veg. Zone 3		.32	.32	4	6	48
RfC:RELIANCE----	100	3e-	3e	All areas are prime farmland	C	Silty - Veg. Zone 3		.32	.32	4	7	38
Rw:RIVERWASH----	100	N/A	8w	Not prime farmland	D	Unspecified		.15	.15	5	8	0
SaG:SANSARC----	100	N/A	7s	Not prime farmland	D	Shallow Clay - Veg. Zone 3		.37	.37	2	4	86
Sc:SCOTT-----	100	N/A	4w	Not prime farmland	D	Clayey Overflow - Veg. Zone 3		.37	.37	3	6	48
Sm:SIMEON-----	100	4s-	6s	Not prime farmland	A	Shallow To Gravel - Veg. Zone 3		.17	.17	5	2	134
SuC:SIMEON-----	55	4s-	6s	Not prime farmland	A	Shallow To Gravel - Veg. Zone 3		.17	.17	5	2	134
SuC:VALENTINE---	45	4e-	6e	Not prime farmland	A	Sands - Veg. Zone 3		.17	.17	5	2	134

Boyd County, Nebraska: Published
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
SvF2:SIMEON----	70	N/A	6s	Not prime farmland	A	Shallow To Gravel - Veg. Zone 3		.17	.17	5	2	134
SvF2:VALENTINE--	30	N/A	6e	Not prime farmland	A	Sands - Veg. Zone 3		.15	.15	5	1	250
VaE:VALENTINE---	100	N/A	6e	Not prime farmland	A	Sands - Veg. Zone 3		.15	.15	5	1	250
VbB:VALENTINE---	100	4e-	4e	Not prime farmland	A	Sandy - Veg. Zone 3		.17	.17	5	2	134
Ve:VERDEL-----	100	2s-	2s	All areas are prime farmland	D	Clayey - Veg. Zone 3		.28	.28	5	4	86
WeC:WEWELA-----	100	3e-	3e	Not prime farmland	B	Sandy - Veg. Zone 3		.24	.24	3	3	86
zwa:WATER-----	100	N/A	N/A	Not prime farmland		Unspecified		---	---	-	---	---

RANGELAND PRODUCTIVITY
Boyd County, Nebraska

Use and Explanation of Rangeland, Grazed Forest Land, Native Pastureland Interpretations

Information in this subsection can be used to plan the use and management of soils for rangeland, grazed forest land, and native pasture. Different kinds of soils vary in their capacity to produce native grasses and other plants suitable for grazing. Information in this subsection provides groupings of similar soils and estimates of potential forage production, which can be used to determine livestock stocking rates.

Rangeland. Range is land on which the native vegetation (climax or natural potential plant community) is predominantly grasses, grasslike plants, forbs, and shrubs suitable for grazing and browsing. Range includes natural grasslands, savannas, many wetlands, some deserts, tundra, and certain shrub and forb communities. Rangeland receives no regular or frequent cultural treatment. The composition and production of the plant community are determined by soil, climate, topography, overstory canopy, and grazing management.

Grazed Forest Land. Includes land on which the understory includes, as an integral part of the forest plant community, plants that can be grazed without significantly impairing other forest values.

Native Pasture. Includes land on which the native vegetation (climax or natural potential plant community) is forest but which is used and managed primarily for production of native plants for forage. Native pasture includes cut-over forest land and forest land cleared and now managed for native or naturalized forage plants.

Rangeland

In areas that have similar climate and topography, differences in the kind and amount of vegetation produced on rangeland are closely related to the kind of soil. Effective management based on the relationship between the soils and vegetation and water.

The Rangeland, Grazed Forest land, Native Pastureland Interpretations shows, for each soil that supports rangeland vegetation, the ecological site and the potential annual production of vegetation in favorable, normal, unfavorable years. An explanation of the column headings in this table follows.

An ecological site is the product of all the environmental factors responsible for its development. It has characteristic soils that have developed over time throughout the soil development process; a characteristic hydrology, particularly infiltration and runoff, that has developed over time; and a characteristic plant community (kind and amount of vegetation). The hydrology of a site is influenced by development of the soil and plant community. The vegetation, soils, and hydrology are all interrelated. Each is influenced by the others and influences the development of the others. The plant community on an ecological site is typified by an association of species that differs from that of other ecological sites in the kind and/or proportion of species or in total production. Descriptions of ecological sites are provided in the Field Office Technical Guide, which is available in local offices of the Natural Resources Conservation Service.

Total dry-weight production is the amount of vegetation that can be expected to grow annually on well managed rangeland that is supporting the potential natural plant community. It includes all vegetation, whether or not it is palatable to grazing animals. It includes the current year's growth of leaves, twigs, and fruits of woody plants. It does not include the increase in stem diameter of trees and shrubs. It is expressed in pounds per acre of air-dry vegetation for favorable, average, and unfavorable years. In a favorable year, the amount and distribution of precipitation and the temperatures make growing conditions substantially better than average. In a normal year, growing conditions are about average. In an unfavorable year, growing conditions are well below average, generally because of low available soil moisture. Yields are adjusted to a common percent of air-dry moisture content.

Range management requires a knowledge of the kinds of soil and of the potential natural plant community. It also requires an evaluation of the present range similarity index and rangeland trend. Range similarity index is determined by comparing the present plant community with the potential natural plant community on a particular rangeland ecological site. The more closely the existing community resembles the potential community, the higher the range similarity index. Rangeland trend is defined as the direction of change in an existing plant community relative to the potential natural plant community. Further information about the range similarity index and rangeland trend is available in chapter 4 of the National Range and Pasture Handbook, which is available in local offices of the Natural Resources Conservation Service. The objective in range management is to control grazing so that the plants growing on a site are about the same in kind and amount as the potential natural plant community for that site. Such management generally results in the optimum production of vegetation, control of undesirable brush species, conservation of water, and control of erosion. Sometimes, however, an area with a range similarity index somewhat below the potential meets grazing needs, provides wildlife habitat, and protects soil and water resources.

RANGELAND PRODUCTIVITY--Continued
Boyd County, Nebraska

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
Ab:				
Albaton-----	Clayey Overflow - Veg. Zone 3	3,200	2,700	2,200
AnC:				
Anselmo-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
AnD:				
Anselmo-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
AnF:				
Anselmo-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
ArF:				
Anselmo-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
Rock Outcrop-----	No Site - Veg. Zone 3	0	0	0
Ba:				
Barney-----	Wet Land - Veg. Zone 3	6,000	5,400	5,000
Bd:				
Blake-----	Silty Lowland - Veg. Zone 3	3,500	3,100	2,700
Be:				
Blendon-----	Sandy - Veg. Zone 3	3,500	2,900	2,000
BeC:				
Blendon-----	Sandy - Veg. Zone 3	3,500	2,900	2,000
BoD:				
Boyd-----	Clayey - Veg. Zone 3	3,200	2,700	1,900
BrG:				
Bristow-----	Shallow Limy - Veg. Zone 3	2,300	2,000	1,800
Bs:				
Brocksburg-----	Sandy - Veg. Zone 3	3,300	3,000	2,600
Bt:				
Brocksburg-----	Silty - Veg. Zone 3	3,700	3,200	2,700
Cb:				
Cass-----	Sandy Lowland - Veg. Zone 3	3,500	3,200	3,000
CrE2:				
Crofton-----	Limy Upland - Veg. Zone 3	3,300	3,000	2,700
DuB:				
Dunday-----	Sandy - Veg. Zone 3	3,300	3,000	2,600
DuC:				
Dunday-----	Sandy - Veg. Zone 3	3,300	3,000	2,600
DuD:				
Dunday-----	Sandy - Veg. Zone 3	3,300	3,000	2,600
DxB:				
Dunday-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
Et:				
Eltree-----	Silty - Veg. Zone 3	4,000	3,500	2,500
Go:				
Grigston-----	Silty Lowland - Veg. Zone 3	5,000	4,000	3,000
GrB:				
Aowa-----	Silty Overflow - Veg. Zone 3	3,500	3,300	3,000
Ha:				
Hall-----	Silty Lowland - Veg. Zone 3	4,000	3,600	3,300
He:				
Haynie-----	Silty Lowland - Veg. Zone 3	4,500	4,200	2,800
IfD:				
Inavale-----	Sands - Veg. Zone 3	3,300	2,800	2,200
IgB:				
Inavale-----	Sandy Lowland - Veg. Zone 3	3,300	2,800	2,200
IhB:				
Inavale-----	Sandy Lowland - Veg. Zone 3	3,500	3,000	2,200
In:				
Inavale-----	Sandy Lowland - Veg. Zone 3	3,200	2,900	2,600
Jn:				
Jansen-----	Silty - Veg. Zone 3	3,700	3,200	2,700
JnC:				
Jansen-----	Silty - Veg. Zone 3	3,700	3,200	2,700
JnD:				
Jansen-----	Silty - Veg. Zone 3	3,700	3,200	2,700
LaD:				
Labu-----	Clayey - Veg. Zone 3	3,400	3,000	2,600
LcF:				
Labu-----	Clayey - Veg. Zone 3	3,400	3,000	2,600
Sansarc-----	Shallow Clay - Veg. Zone 3	2,500	2,100	1,500
Le:				
Leshara-----	Subirrigated - Veg. Zone 3	5,500	5,300	5,000
LsC:				
Lynch-----	Limy Upland - Veg. Zone 3	3,000	2,700	2,400
LsD:				
Lynch-----	Limy Upland - Veg. Zone 3	3,000	2,700	2,400
LyD:				
Lynch-----	Limy Upland - Veg. Zone 3	3,000	2,700	2,400
Bristow-----	Shallow Limy - Veg. Zone 3	2,300	2,000	1,800
LyF:				
Lynch-----	Limy Upland - Veg. Zone 3	3,000	2,700	2,400
Bristow-----	Shallow Limy - Veg. Zone 3	2,300	2,000	1,800
MaG:				
Mariaville-----	Shallow Limy - Veg. Zone 3	2,300	2,000	1,800
Paka-----	Silty - Veg. Zone 3	4,000	3,600	3,300
MeE:				

RANGELAND PRODUCTIVITY--Continued
Boyd County, Nebraska

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
Meadin-----	Shallow To Gravel - Veg. Zone 3	1,500	1,300	1,100
NoC:				
Nora-----	Silty - Veg. Zone 3	4,000	3,600	3,300
NoD:				
Nora-----	Silty - Veg. Zone 3	4,000	3,600	3,300
Oa:				
Onawa-----	Clayey Overflow - Veg. Zone 3	3,500	3,100	2,700
Oe:				
O'Neill-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
OeC:				
O'Neill-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
OfD:				
O'Neill-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
Meadin-----	Shallow To Gravel - Veg. Zone 3	1,500	1,300	1,100
On:				
Onita-----	Silty - Veg. Zone 3	3,200	2,800	2,000
Poorly Drained-----	---	---	---	---
Or:				
Ord-----	Subirrigated - Veg. Zone 3	5,500	5,300	5,000
PaC:				
Paka-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
Ph:				
Paka-----	Silty - Veg. Zone 3	4,000	3,600	3,300
PhC:				
Paka-----	Silty - Veg. Zone 3	4,000	3,600	3,300
PhD:				
Paka-----	Silty - Veg. Zone 3	4,000	3,600	3,300
PoC:				
Promise-----	Clayey - Veg. Zone 3	3,300	2,800	2,000
RaC:				
Ree-----	Silty - Veg. Zone 3	4,000	3,200	2,200
RaD:				
Ree-----	Silty - Veg. Zone 3	4,000	3,200	2,200
RaE:				
Ree-----	Silty - Veg. Zone 3	4,000	3,200	2,200
ReC:				
Reliance-----	Silty - Veg. Zone 3	4,000	3,200	2,200
ReD:				
Reliance-----	Silty - Veg. Zone 3	4,000	3,200	2,200
RfC:				
Reliance-----	Silty - Veg. Zone 3	4,000	3,200	2,200
Rw:				
Riverwash-----	---	---	---	---
SaG:				
Sansarc-----	Shallow Clay - Veg. Zone 3	2,500	2,100	1,500
Sc:				
Scott-----	Clayey Overflow - Veg. Zone 3	3,900	3,300	2,300
Sm:				
Simeon-----	Shallow To Gravel - Veg. Zone 3	1,800	1,600	1,100
SuC:				
Simeon-----	Shallow To Gravel - Veg. Zone 3	1,400	1,200	700
Valentine-----	Sands - Veg. Zone 3	3,000	2,600	2,200
SvF2:				
Simeon-----	Shallow To Gravel - Veg. Zone 3	1,400	1,200	700
Valentine-----	Sands - Veg. Zone 3	3,000	2,600	2,200
VaE:				
Valentine-----	Sands - Veg. Zone 3	3,000	2,600	2,200
VbB:				
Valentine-----	Sandy - Veg. Zone 3	3,300	3,000	2,600
Ve:				
Verdel-----	Clayey - Veg. Zone 3	3,800	3,400	3,000
WeC:				
Wewela-----	Sandy - Veg. Zone 3	3,600	3,000	2,100
zwa:				
Water-----	---	---	---	---

BUILDING SITE DEVELOPMENT
Boyd County, Nebraska

Building Site Development

Soil properties influence the development of building sites, including the selection of the site, the design of the structure, construction, performance after construction, and maintenance. These tables show the degree and kind of soil limitations that affect dwellings with and without basements, small commercial buildings, local roads and streets, shallow excavations, and lawns and landscaping.

The ratings in the tables are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect building site development. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Dwellings are single-family houses of three stories or less. For dwellings without basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. For dwellings with basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of about 7 feet. The ratings for dwellings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility. Compressibility is inferred from the Unified classification. The properties that affect the ease and amount of excavation include depth to a water table, ponding, flooding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Small commercial buildings are structures that are less than three stories high and do not have basements. The foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. The ratings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility (which is inferred from the Unified classification). The properties that affect the ease and amount of excavation include flooding, depth to a water table, ponding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Local roads and streets have an all-weather surface and carry automobile and light truck traffic all year. They have a subgrade of cut or fill soil material; a base of gravel, crushed rock, or soil material stabilized by lime or cement; and a surface of flexible material (asphalt), rigid material (concrete), or gravel with a binder. The ratings are based on the soil properties that affect the ease of excavation and grading and the traffic-supporting capacity. The properties that affect the ease of excavation and grading are depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, depth to a water table, ponding, flooding, the amount of large stones, and slope. The properties that affect the traffic-supporting capacity are soil strength (as inferred from the AASHTO group index number), subsidence, linear extensibility (shrink-swell potential), the potential for frost action, depth to a water table, and ponding.

Shallow excavations are trenches or holes dug to a maximum depth of 5 or 6 feet for graves, utility lines, open ditches, or other purposes. The ratings are based on the soil properties that influence the ease of digging and the resistance to sloughing. Depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, the amount of large stones, and dense layers influence the ease of digging, filling, and compacting. Depth to the seasonal high water table, flooding, and ponding may restrict the period when excavations can be made. Slope influences the ease of using machinery. Soil texture, depth to the water table, and linear extensibility (shrink-swell potential) influence the resistance to sloughing.

Lawns and landscaping require soils on which turf and ornamental trees and shrubs can be established and maintained. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer.

BUILDING SITE DEVELOPMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00
AnC: Anselmo-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
AnD: Anselmo-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
AnF: Anselmo-----	100	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
ArF: Anselmo-----	75	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
Ba: Barney-----	100	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Bd: Blake-----	100	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Depth to saturated zone	1.00 0.95	Very limited Flooding Shrink-swell	1.00 0.50
Be: Blendon-----	100	Not limited		Not limited		Not limited	
BeC: Blendon-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
BoD: Boyd-----	100	Very limited Shrink-swell Slope	1.00 0.04	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 0.42 0.04	Very limited Shrink-swell Slope	1.00 1.00
BrG: Bristow-----	100	Very limited Slope Depth to soft bedrock Shrink-swell	1.00 1.00 1.00	Very limited Slope Shrink-swell Depth to soft bedrock	1.00 1.00 1.00	Very limited Slope Depth to soft bedrock Shrink-swell	1.00 1.00 1.00
Bs: Brocksburg-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell	0.50
Bt: Brocksburg-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell	0.50
Cb: Cass-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
CrE2: Crofton-----	100	Somewhat limited Slope	0.84	Somewhat limited Slope	0.84	Very limited Slope	1.00
DuB: Dunday-----	100	Not limited		Not limited		Not limited	
DuC: Dunday-----	100	Not limited		Not limited		Somewhat limited Slope	0.12
DuD: Dunday-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
DxB: Dunday-----	100	Not limited		Not limited		Not limited	
Et: Eltree-----	100	Not limited		Not limited		Not limited	
Go: Grigston-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00

BUILDING SITE DEVELOPMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
GrB: Aowa-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Ha: Hall-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
He: Haynie-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
IfD: Inavale-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding Slope	1.00 0.86
IgB: Inavale-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
IhB: Inavale-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
In: Inavale-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Jn: Jansen-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell	0.50
JnC: Jansen-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell Slope	0.50 0.00
JnD: Jansen-----	100	Somewhat limited Shrink-swell Slope	0.50 0.04	Somewhat limited Slope	0.04	Very limited Slope Shrink-swell	1.00 0.50
LaD: Labu-----	100	Very limited Shrink-swell Slope	1.00 0.04	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 0.42 0.04	Very limited Shrink-swell Slope	1.00 1.00
LcF: Labu-----	65	Very limited Shrink-swell Slope	1.00 1.00	Very limited Shrink-swell Slope Depth to soft bedrock	1.00 1.00 0.42	Very limited Slope Shrink-swell	1.00 1.00
Sansarc-----	35	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 1.00 1.00	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 1.00 1.00	Very limited Slope Shrink-swell Depth to soft bedrock	1.00 1.00 1.00
Le: Leshara-----	100	Very limited Flooding Depth to saturated zone	1.00 0.39	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.39
LsC: Lynch-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell Depth to soft bedrock	1.00 0.42	Very limited Shrink-swell Slope	1.00 0.00
LsD: Lynch-----	100	Very limited Shrink-swell Slope	1.00 0.04	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 0.42 0.04	Very limited Shrink-swell Slope	1.00 1.00

BUILDING SITE DEVELOPMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
LyD: Lynch-----	65	Very limited Shrink-swell Slope	1.00 0.04	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 0.42 0.04	Very limited Shrink-swell Slope	1.00 1.00
Bristow-----	35	Very limited Depth to soft bedrock Shrink-swell Slope	1.00 1.00 0.04	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 1.00 0.04	Very limited Depth to soft bedrock Shrink-swell Slope	1.00 1.00 1.00
LyF: Lynch-----	55	Very limited Shrink-swell Slope	1.00 1.00	Very limited Shrink-swell Slope Depth to soft bedrock	1.00 1.00 0.42	Very limited Slope Shrink-swell	1.00 1.00
Bristow-----	45	Very limited Depth to soft bedrock Shrink-swell Slope	1.00 1.00 1.00	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 1.00 1.00	Very limited Slope Depth to soft bedrock Shrink-swell	1.00 1.00 1.00
MaG: Mariaville-----	55	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Slope Depth to soft bedrock	1.00 1.00
Paka-----	45	Very limited Slope Shrink-swell	1.00 0.50	Very limited Slope Shrink-swell	1.00 0.50	Very limited Slope Shrink-swell	1.00 0.50
MeE: Meadin-----	100	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
NoC: Nora-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.00
NoD: Nora-----	100	Somewhat limited Shrink-swell Slope	0.50 0.04	Somewhat limited Shrink-swell Slope	0.50 0.04	Very limited Slope Shrink-swell	1.00 0.50
Oa: Onawa-----	100	Very limited Flooding Shrink-swell	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.95	Very limited Flooding Shrink-swell	1.00 1.00
Oe: O'Neill-----	100	Not limited		Not limited		Not limited	
OeC: O'Neill-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
OfD: O'Neill-----	55	Not limited		Not limited		Somewhat limited Slope	0.48
Meadin-----	45	Not limited		Not limited		Somewhat limited Slope	0.48
On: Onita-----	100	Very limited Shrink-swell	1.00	Somewhat limited Shrink-swell	0.50	Very limited Shrink-swell	1.00
Poorly Drained-----		Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Or: Ord-----	100	Very limited Flooding Depth to saturated zone	1.00 0.07	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.07
PaC: Paka-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.00
Ph: Paka-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50

BUILDING SITE DEVELOPMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
PhC: Paka-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.00
PhD: Paka-----	100	Somewhat limited Shrink-swell Slope	0.50 0.04	Somewhat limited Shrink-swell Slope	0.50 0.04	Very limited Slope Shrink-swell	1.00 0.50
PoC: Promise-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell Slope	1.00 0.00
RaC: Ree-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
RaD: Ree-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
RaE: Ree-----	100	Somewhat limited Slope	0.84	Somewhat limited Slope	0.84	Very limited Slope	1.00
ReC: Reliance-----	100	Very limited Shrink-swell	1.00	Somewhat limited Shrink-swell	0.50	Very limited Shrink-swell Slope	1.00 0.00
ReD: Reliance-----	100	Very limited Shrink-swell Slope	1.00 0.04	Somewhat limited Shrink-swell Slope	0.50 0.04	Very limited Shrink-swell Slope	1.00 1.00
RfC: Reliance-----	100	Very limited Shrink-swell	1.00	Somewhat limited Shrink-swell	0.50	Very limited Shrink-swell Slope	1.00 0.00
Rw: Riverwash-----	100	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
SaG: Sansarc-----	100	Very limited Slope Shrink-swell Depth to soft bedrock	1.00 1.00 1.00	Very limited Slope Shrink-swell Depth to soft bedrock	1.00 1.00 1.00	Very limited Slope Shrink-swell Depth to soft bedrock	1.00 1.00 1.00
Sc: Scott-----	100	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Shrink-swell	1.00 1.00
Sm: Simeon-----	100	Not limited		Not limited		Not limited	
SuC: Simeon-----	55	Not limited		Not limited		Not limited	
SvF2: Valentine-----	45	Not limited		Not limited		Not limited	
SvF2: Simeon-----	70	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
VaE: Valentine-----	30	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
VaE: Valentine-----	100	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
VbB: Valentine-----	100	Not limited		Not limited		Not limited	
Ve: Verdel-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00
WeC: Wewela-----	100	Not limited		Somewhat limited Depth to soft bedrock	0.42	Somewhat limited Slope	0.00
zwa: Water-----	100	Not rated		Not rated		Not rated	

BUILDING SITE DEVELOPMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Very limited Shrink-swell	1.00	Very limited Depth to saturated zone	1.00	Very limited Too clayey	1.00
		Depth to saturated zone	1.00	Too clayey	1.00	Depth to saturated zone	1.00
		Flooding	1.00	Flooding	0.60	Flooding	0.60
		Low strength	1.00	Cutbanks cave	0.10		
		Frost action	0.50				
AnC: Anselmo-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
AnD: Anselmo-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Somewhat limited Slope	0.04
		Slope	0.04	Slope	0.04		
AnF: Anselmo-----	100	Somewhat limited Slope	0.96	Very limited Cutbanks cave	1.00	Somewhat limited Slope	0.96
		Frost action	0.50	Slope	0.96		
ArF: Anselmo-----	75	Somewhat limited Slope	0.96	Very limited Cutbanks cave	1.00	Somewhat limited Slope	0.96
		Frost action	0.50	Slope	0.96		
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
Ba: Barney-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Flooding	1.00	Cutbanks cave	1.00	Depth to saturated zone	1.00
		Frost action	0.50	Flooding	0.80	Droughty	0.13
				Depth to dense layer	0.50		
Bd: Blake-----	100	Very limited Frost action	1.00	Somewhat limited Depth to saturated zone	0.95	Somewhat limited Flooding	0.60
		Flooding	1.00	Flooding	0.60		
		Shrink-swell	0.50	Cutbanks cave	0.10		
Be: Blendon-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
BeC: Blendon-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
BoD: Boyd-----	100	Very limited Shrink-swell	1.00	Somewhat limited Too clayey	0.88	Very limited Too clayey	1.00
		Slope	0.04	Depth to soft bedrock	0.42	Depth to bedrock	0.42
				Cutbanks cave	0.10	Droughty	0.34
				Slope	0.04	Slope	0.04
BrG: Bristow-----	100	Very limited Slope	1.00	Very limited Depth to soft bedrock	1.00	Very limited Depth to bedrock	1.00
		Depth to soft bedrock	1.00	Slope	1.00	Slope	1.00
		Shrink-swell	1.00	Too clayey	0.72	Droughty	1.00
		Frost action	0.50	Cutbanks cave	0.10	Too clayey	1.00
Bs: Brocksburg-----	100	Somewhat limited Shrink-swell	0.50	Very limited Cutbanks cave	1.00	Not limited	
		Frost action	0.50				
Bt: Brocksburg-----	100	Somewhat limited Shrink-swell	0.50	Very limited Cutbanks cave	1.00	Not limited	
		Frost action	0.50				
Cb: Cass-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave	1.00	Somewhat limited Flooding	0.60
		Frost action	0.50	Flooding	0.60		

BUILDING SITE DEVELOPMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
CrE2: Crofton-----	100	Somewhat limited Slope Frost action	0.84 0.50	Somewhat limited Slope Cutbanks cave	0.84 0.10	Somewhat limited Slope	0.84
DuB: Dunday-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.07
DuC: Dunday-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.07
DuD: Dunday-----	100	Somewhat limited Slope	0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Droughty Slope	0.07 0.04
DxB: Dunday-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.02
Et: Eltree-----	100	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Go: Grigston-----	100	Very limited Flooding Frost action	1.00 0.50	Somewhat limited Flooding Cutbanks cave	0.60 0.10	Somewhat limited Flooding	0.60
GrB: Aowa-----	100	Very limited Frost action Flooding	1.00 1.00	Very limited Cutbanks cave Flooding	1.00 0.80	Very limited Flooding	1.00
Ha: Hall-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
He: Haynie-----	100	Very limited Frost action Flooding	1.00 1.00	Very limited Cutbanks cave Flooding	1.00 0.60	Somewhat limited Flooding	0.60
IfD: Inavale-----	100	Somewhat limited Flooding	0.40	Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.29
IgB: Inavale-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave Flooding	1.00 0.80	Very limited Flooding Droughty	1.00 0.29
IhB: Inavale-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave Flooding	1.00 0.60	Somewhat limited Flooding Droughty	0.60 0.10
In: Inavale-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave Flooding	1.00 0.60	Somewhat limited Flooding	0.60
Jn: Jansen-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Very limited Cutbanks cave	1.00	Not limited	
JnC: Jansen-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Very limited Cutbanks cave	1.00	Not limited	
JnD: Jansen-----	100	Somewhat limited Shrink-swell Frost action Slope	0.50 0.50 0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Slope	0.04
LaD: Labu-----	100	Very limited Shrink-swell Slope	1.00 0.04	Somewhat limited Too clayey Depth to soft bedrock Cutbanks cave Slope	0.72 0.42 0.10 0.04	Very limited Too clayey Depth to bedrock Droughty Slope	1.00 0.42 0.20 0.04

BUILDING SITE DEVELOPMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
LcF: Labu-----	65	Very limited Shrink-swell Slope	1.00 1.00	Very limited Slope Too clayey Depth to soft bedrock Cutbanks cave	1.00 0.72 0.42 0.10	Very limited Too clayey Slope Depth to bedrock	1.00 1.00 0.42
Sansarc-----	35	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 1.00 1.00	Very limited Depth to soft bedrock Slope Too clayey Cutbanks cave	1.00 1.00 1.00 0.82 0.10	Droughty Very limited Too clayey Depth to bedrock Droughty Slope	0.20 1.00 1.00 1.00
Le: Leshara-----	100	Very limited Frost action Flooding Depth to saturated zone	1.00 1.00 0.19	Very limited Cutbanks cave Depth to saturated zone Flooding	1.00 1.00 0.60	Somewhat limited Flooding Depth to saturated zone	0.60 0.19
LsC: Lynch-----	100	Very limited Shrink-swell	1.00	Somewhat limited Too clayey Depth to soft bedrock Cutbanks cave	0.88 0.42 0.10	Very limited Too clayey Depth to bedrock Droughty	1.00 0.42 0.09
LsD: Lynch-----	100	Very limited Shrink-swell Slope	1.00 0.04	Somewhat limited Too clayey Depth to soft bedrock Cutbanks cave Slope	0.88 0.42 0.10 0.04	Very limited Too clayey Depth to bedrock Droughty Slope	1.00 0.42 0.09 0.04
LyD: Lynch-----	65	Very limited Shrink-swell Slope	1.00 0.04	Somewhat limited Too clayey Depth to soft bedrock Cutbanks cave Slope	0.88 0.42 0.10 0.04	Very limited Too clayey Depth to bedrock Droughty Slope	1.00 0.42 0.09 0.04
Bristow-----	35	Very limited Depth to soft bedrock Shrink-swell Frost action Slope	1.00 1.00 0.50 0.04	Very limited Depth to soft bedrock Too clayey Cutbanks cave Slope	1.00 0.72 0.10 0.04	Very limited Depth to bedrock Droughty Too clayey Slope	1.00 1.00 1.00 0.04
LyF: Lynch-----	55	Very limited Shrink-swell Slope	1.00 1.00	Very limited Slope Too clayey Depth to soft bedrock Cutbanks cave	1.00 0.88 0.42 0.10	Very limited Slope Too clayey Depth to bedrock Droughty	1.00 1.00 0.42 0.09
Bristow-----	45	Very limited Depth to soft bedrock Shrink-swell Slope Frost action	1.00 1.00 1.00 0.50	Very limited Depth to soft bedrock Slope Too clayey Cutbanks cave	1.00 1.00 0.72 0.10	Very limited Depth to bedrock Droughty Slope Too clayey	1.00 1.00 1.00 1.00
MaG: Mariaville-----	55	Very limited Depth to soft bedrock Slope Frost action	1.00 1.00 0.50	Very limited Depth to soft bedrock Slope Cutbanks cave	1.00 1.00 0.10	Very limited Depth to bedrock Slope Droughty	1.00 0.43
Paka-----	45	Very limited Slope Shrink-swell Frost action	1.00 0.50 0.50	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
MeE: Meadin-----	100	Somewhat limited Slope	0.16	Very limited Cutbanks cave Slope	1.00 0.16	Somewhat limited Droughty Slope	0.63 0.16
NoC: Nora-----	100	Very limited Frost action Shrink-swell	1.00 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	

BUILDING SITE DEVELOPMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
NoD: Nora-----	100	Very limited Frost action Shrink-swell Slope	1.00 0.50 0.04	Somewhat limited Cutbanks cave Slope	0.10 0.04	Somewhat limited Slope	0.04
Oa: Onawa-----	100	Very limited Frost action Flooding Shrink-swell	1.00 1.00 1.00	Somewhat limited Depth to saturated zone Too clayey Flooding Cutbanks cave	0.95 0.88 0.60 0.10	Very limited Too clayey Flooding	1.00 0.60
Oe: O'Neill-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
OeC: O'Neill-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
OfD: O'Neill-----	55	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
Meadin-----	45	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.63
On: Onita-----	100	Very limited Shrink-swell Frost action	1.00 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Poorly Drained-----		Very limited Slope Low strength	1.00 1.00	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
Or: Ord-----	100	Very limited Frost action Flooding Depth to saturated zone	1.00 1.00 0.03	Very limited Cutbanks cave Depth to saturated zone Flooding	1.00 1.00 0.60	Somewhat limited Flooding Depth to saturated zone	0.60 0.03
PaC: Paka-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Ph: Paka-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
PhC: Paka-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
PhD: Paka-----	100	Somewhat limited Shrink-swell Frost action Slope	0.50 0.50 0.04	Somewhat limited Cutbanks cave Slope	0.10 0.04	Somewhat limited Slope	0.04
PoC: Promise-----	100	Very limited Shrink-swell	1.00	Very limited Too clayey Cutbanks cave	1.00 1.00	Very limited Too clayey	1.00
RaC: Ree-----	100	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
RaD: Ree-----	100	Somewhat limited Frost action Slope	0.50 0.04	Somewhat limited Cutbanks cave Slope	0.10 0.04	Somewhat limited Slope	0.04
RaE: Ree-----	100	Somewhat limited Slope Frost action	0.84 0.50	Somewhat limited Slope Cutbanks cave	0.84 0.10	Somewhat limited Slope	0.84
ReC: Reliance-----	100	Very limited Shrink-swell	1.00	Somewhat limited Cutbanks cave	0.10	Not limited	
ReD: Reliance-----	100	Very limited Shrink-swell Slope	1.00 0.04	Somewhat limited Cutbanks cave Slope	0.10 0.04	Somewhat limited Slope	0.04

BUILDING SITE DEVELOPMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
RfC: Reliance-----	100	Very limited Shrink-swell	1.00	Somewhat limited Cutbanks cave	0.10	Not limited	
Rw: Riverwash-----	100	Very limited Depth to saturated zone Flooding	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Flooding	1.00 1.00 0.80	Very limited Flooding Depth to saturated zone Droughty	1.00 1.00 1.00
SaG: Sansarc-----	100	Very limited Shrink-swell Slope Depth to soft bedrock	1.00 1.00 1.00	Very limited Depth to soft bedrock Slope Too clayey Cutbanks cave	1.00 1.00 1.00 0.10	Very limited Too clayey Depth to bedrock Slope Droughty	1.00 1.00 1.00 1.00
Sc: Scott-----	100	Very limited Depth to saturated zone Frost action Shrink-swell	1.00 1.00 1.00	Very limited Depth to saturated zone Too clayey Cutbanks cave	1.00 0.28 0.10	Very limited Depth to saturated zone	1.00
Sm: Simeon-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.22
SuC: Simeon-----	55	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.07
Valentine-----	45	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.61
SvF2: Simeon-----	70	Very limited Slope	1.00	Very limited Cutbanks cave Slope	1.00 1.00	Very limited Slope Droughty	1.00 0.07
Valentine-----	30	Very limited Slope	1.00	Very limited Cutbanks cave Slope	1.00 1.00	Very limited Slope Droughty	1.00 0.73
VaE: Valentine-----	100	Somewhat limited Slope	0.63	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Droughty Slope	0.73 0.63
VbB: Valentine-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.61
Ve: Verdel-----	100	Very limited Shrink-swell Frost action	1.00 0.50	Somewhat limited Too clayey Cutbanks cave	0.50 0.10	Very limited Too clayey	1.00
WeC: Wewela-----	100	Somewhat limited Frost action	0.50	Somewhat limited Too clayey Depth to soft bedrock Cutbanks cave	0.50 0.42 0.10	Somewhat limited Depth to bedrock Droughty	0.42 0.32
zwa: Water-----	100	Not rated		Not rated		Not rated	

CONSTRUCTION MATERIALS
Boyd County, Nebraska

Construction Materials

These tables give information about the soils as potential sources of gravel, sand, topsoil, reclamation material, and roadfill. Normal compaction, minor processing, and other standard construction practices are assumed.

The soils are rated good, fair, or poor as potential sources of topsoil, reclamation material, and roadfill. The features that limit the soils as sources of these materials are specified in the tables. The numerical ratings given after the specified features indicate the degree to which the features limit the soils as sources of topsoil, reclamation material, or roadfill. The lower the number, the greater the limitation.

The soils are rated as a probable or improbable source of sand and gravel. A rating of probable means that the source material is likely to be in or below the soil. The numerical ratings in these columns indicate the degree of probability. The number 0.00 indicates that the soil is an improbable source. A number between 0.00 and 1.00 indicates the degree to which the soil is a probable source of sand or gravel.

Sand and gravel are natural aggregates suitable for commercial use with a minimum of processing. They are used in many kinds of construction. Specifications for each use vary widely. In the first table, only the probability of finding material in suitable quantity is evaluated. The suitability of the material for specific purposes is not evaluated, nor are factors that affect excavation of the material. The properties used to evaluate the soil as a source of sand or gravel are gradation of grain sizes (as indicated by the Unified classification of the soil), the thickness of suitable material, and the content of rock fragments. If the lowest layer of the soil contains sand or gravel, the soil is rated as a probable source regardless of thickness. The assumption is that the sand or gravel layer below the depth of observation exceeds the minimum thickness.

Topsoil is used to cover an area so that vegetation can be established and maintained. The upper 40 inches of a soil is evaluated for use as topsoil. Also evaluated is the reclamation potential of the borrow area. The ratings are based on the soil properties that affect plant growth; the ease of excavating, loading, and spreading the material; and reclamation of the borrow area. Toxic substances, soil reaction, and the properties that are inferred from soil texture, such as available water capacity and fertility, affect plant growth. The ease of excavating, loading, and spreading is affected by rock fragments, slope, depth to a water table, soil texture, and thickness of suitable material. Reclamation of the borrow area is affected by slope, depth to a water table, rock fragments, depth to bedrock or a cemented pan, and toxic material.

The surface layer of most soils is generally preferred for topsoil because of its organic matter content. Organic matter greatly increases the absorption and retention of moisture and nutrients for plant growth.

Reclamation material is used in areas that have been drastically disturbed by surface mining or similar activities. When these areas are reclaimed, layers of soil material or unconsolidated geological material, or both, are replaced in a vertical sequence. The reconstructed soil favors plant growth. The ratings in the table do not apply to quarries and other mined areas that require an offsite source of reconstruction material. The ratings are based on the soil properties that affect erosion and stability of the surface and the productive potential of the reconstructed soil. These properties include the content of sodium, salts, and calcium carbonate; reaction; available water capacity; erodibility; texture; content of rock fragments; and content of organic matter and other features that affect fertility.

Roadfill is soil material that is excavated in one place and used in road embankments in another place. In this table, the soils are rated as a source of roadfill for low embankments, generally less than 6 feet high and less exacting in design than higher embankments.

The ratings are for the whole soil, from the surface to a depth of about 5 feet. It is assumed that soil layers will be mixed when the soil material is excavated and spread.

The ratings are based on the amount of suitable material and on soil properties that affect the ease of excavation and the performance of the material after it is in place. The thickness of the suitable material is a major consideration. The ease of excavation is affected by large stones, depth to a water table, and slope. How well the soil performs in place after it has been compacted and drained is determined by its strength (as inferred from the AASHTO classification of the soil) and linear extensibility (shrink-swell potential).

CONSTRUCTION MATERIALS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Ab: Albaton-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
AnC: Anselmo-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.57
AnD: Anselmo-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.57
AnF: Anselmo-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.57
ArF: Anselmo-----	75	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.57
Rock Outcrop-----	25	Not rated		Not rated	
Ba: Barney-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.95
Bd: Blake-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.05
Be: Blendon-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.07 0.09
BeC: Blendon-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.07 0.09
BoD: Boyd-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
BrG: Bristow-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Bs: Brocksburg-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.08
Bt: Brocksburg-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.00
Cb: Cass-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.08 0.95
CrE2: Crofton-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
DuB: Dunday-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.65
DuC: Dunday-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.65
DuD: Dunday-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.65
DxB: Dunday-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.29 0.91
Et: Eltree-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Go: Grigston-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
GrB: Aowa-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ha: Hall-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
He: Haynie-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
IfD: Inavale-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Bottom layer	0.98
IgB: Inavale-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Bottom layer	0.98
IhB: Inavale-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.98 0.98
In: Inavale-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.07 0.98
Jn: Jansen-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.97
JnC: Jansen-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.97
JnD: Jansen-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.97

CONSTRUCTION MATERIALS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
LaD: Labu-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
LcF: Labu-----	65	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Sansarc-----	35	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Le: Leshara-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
LsC: Lynch-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
LsD: Lynch-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
LyD: Lynch-----	65	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Bristow-----	35	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
LyF: Lynch-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Bristow-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MaG: Mariaville-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Paka-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MeE: Meadin-----	100	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.91
NoC: Nora-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
NoD: Nora-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Oa: Onawa-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Oe: O'Neill-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.09
OeC: O'Neill-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.09
OfD: O'Neill-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.09
Meadin-----	45	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.08 0.91
On: Onita-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Poorly Drained-----		Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Or: Ord-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.07 0.97
PaC: Paka-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ph: Paka-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
PhC: Paka-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
PhD: Paka-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
PoC: Promise-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
RaC: Ree-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.09
RaD: Ree-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.09
RaE: Ree-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.03 0.04
ReC: Reliance-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
ReD: Reliance-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
RfC: Reliance-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Rw: Riverwash-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.86 0.86
SaG: Sansarc-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Sc: Scott-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Sm: Simeon-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.39 0.98
SuC: Simeon-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.39 0.98
Valentine-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.99
SvF2: Simeon-----	70	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.39 0.98
Valentine-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.99
VaE: Valentine-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.99
VbB: Valentine-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.70
Ve: Verdel-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
WeC: Wewela-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
zwa: Water-----	100	Not rated		Not rated	

CONSTRUCTION MATERIALS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Poor Too clayey	0.00	Poor Depth to saturated zone Shrink-swell Low strength	0.00 0.00 0.00	Poor Too Clayey Depth to saturated zone	0.00 0.00
AnC: Anselmo-----	100	Fair Low content of organic matter	0.12	Good		Good	
AnD: Anselmo-----	100	Fair Low content of organic matter	0.12	Good		Fair Slope	0.96
AnF: Anselmo-----	100	Fair Low content of organic matter	0.12	Good		Fair Slope	0.04
ArF: Anselmo-----	75	Fair Low content of organic matter	0.12	Good		Fair Slope	0.04
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
Ba: Barney-----	100	Poor Too sandy Low content of organic matter Droughty	0.00 0.12 0.66	Poor Depth to saturated zone	0.00	Poor Hard to reclaim Too sandy Depth to saturated zone Rock fragments	0.00 0.00 0.00 0.97
Bd: Blake-----	100	Fair Low content of organic matter Water erosion Carbonate content	0.12 0.90 0.97	Good		Fair Carbonate content	0.97
Be: Blendon-----	100	Fair Low content of organic matter	0.88	Good		Good	
BeC: Blendon-----	100	Fair Low content of organic matter	0.88	Good		Good	
BoD: Boyd-----	100	Poor Too clayey Droughty Depth to bedrock Water erosion	0.00 0.01 0.58 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.00	Poor Too Clayey Depth to bedrock Slope	0.00 0.58 0.96
BrG: Bristow-----	100	Poor Too clayey Droughty Depth to bedrock Low content of organic matter Carbonate content Water erosion	0.00 0.00 0.00 0.12 0.46 0.90	Poor Depth to bedrock Slope Shrink-swell	0.00 0.00 0.12	Poor Slope Too Clayey Depth to bedrock Carbonate content	0.00 0.00 0.00 0.46
Bs: Brocksburg-----	100	Fair Low content of organic matter	0.12	Good		Fair Hard to reclaim	0.92

CONSTRUCTION MATERIALS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Bt: Brocksburg-----	100	Fair Low content of organic matter	0.12	Good		Fair Hard to reclaim	0.92
Cb: Cass-----	100	Fair Low content of organic matter	0.12	Good		Good	
CrE2: Crofton-----	100	Fair Low content of organic matter Water erosion	0.12 0.90	Good		Fair Slope	0.16
DuB: Dunday-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.90	Good		Poor Too sandy	0.00
DuC: Dunday-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.90	Good		Poor Too sandy	0.00
DuD: Dunday-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.90	Good		Poor Too sandy Slope	0.00 0.96
DxB: Dunday-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.88 0.88	Good		Poor Too sandy Hard to reclaim	0.00 0.92
Et: Eltree-----	100	Fair Water erosion	0.90	Good		Good	
Go: Grigston-----	100	Fair Low content of organic matter	0.12	Good		Good	
GrB: Aowa-----	100	Fair Low content of organic matter Water erosion	0.12 0.90	Good		Good	
Ha: Hall-----	100	Fair Water erosion	0.90	Fair Shrink-swell	0.99	Good	
He: Haynie-----	100	Fair Low content of organic matter Water erosion Carbonate content	0.50 0.90 0.97	Good		Fair Carbonate content	0.97

CONSTRUCTION MATERIALS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
IfD: Inavale-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.74	Good		Poor Too sandy	0.00
IgB: Inavale-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.74	Good		Poor Too sandy	0.00
IhB: Inavale-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.87	Good		Poor Too sandy	0.00
In: Inavale-----	100	Fair Low content of organic matter	0.12	Good		Good	
Jn: Jansen-----	100	Fair Low content of organic matter Too acid Water erosion	0.12 0.97 0.99	Good		Fair Hard to reclaim	0.98
JnC: Jansen-----	100	Fair Low content of organic matter Too acid Water erosion	0.12 0.97 0.99	Good		Fair Hard to reclaim	0.98
JnD: Jansen-----	100	Fair Low content of organic matter Too acid Water erosion	0.12 0.97 0.99	Good		Fair Slope Hard to reclaim	0.96 0.98
LaD: Labu-----	100	Poor Too clayey Droughty Depth to bedrock Low content of organic matter	0.00 0.03 0.58 0.88	Poor Depth to bedrock Shrink-swell	0.00 0.12	Poor Too Clayey Depth to bedrock Slope	0.00 0.58 0.96
LcF: Labu-----	65	Poor Too clayey Droughty Depth to bedrock Low content of organic matter	0.00 0.03 0.58 0.88	Poor Depth to bedrock Shrink-swell Slope	0.00 0.12 0.32	Poor Too Clayey Slope Depth to bedrock	0.00 0.00 0.58
Sansarc-----	35	Poor Too clayey Droughty Depth to bedrock Low content of organic matter Water erosion	0.00 0.00 0.00 0.60 0.99	Poor Depth to bedrock Shrink-swell Slope	0.00 0.00 0.32	Poor Too Clayey Depth to bedrock Slope	0.00 0.00 0.00
Le: Leshara-----	100	Fair Low content of organic matter Water erosion	0.12 0.90	Fair Depth to saturated zone	0.53	Fair Depth to saturated zone	0.53

CONSTRUCTION MATERIALS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
LsC: Lynch-----	100	Poor Too clayey Droughty Carbonate content Depth to bedrock Low content of organic matter	0.00 0.08 0.08 0.58 0.88	Poor Depth to bedrock Shrink-swell	0.00 0.12	Poor Too Clayey Carbonate content Depth to bedrock	0.00 0.08 0.58
LsD: Lynch-----	100	Poor Too clayey Droughty Carbonate content Depth to bedrock Low content of organic matter	0.00 0.08 0.08 0.58 0.88	Poor Depth to bedrock Shrink-swell	0.00 0.12	Poor Too Clayey Carbonate content Depth to bedrock Slope	0.00 0.08 0.58 0.96
LyD: Lynch-----	65	Poor Too clayey Droughty Carbonate content Depth to bedrock Low content of organic matter	0.00 0.08 0.08 0.58 0.88	Poor Depth to bedrock Shrink-swell	0.00 0.12	Poor Too Clayey Carbonate content Depth to bedrock Slope	0.00 0.08 0.58 0.96
Bristow-----	35	Poor Too clayey Droughty Depth to bedrock Low content of organic matter Carbonate content Water erosion	0.00 0.00 0.00 0.12 0.46 0.90	Poor Depth to bedrock Shrink-swell	0.00 0.12	Poor Too Clayey Depth to bedrock Carbonate content Slope	0.00 0.00 0.46 0.96
LyF: Lynch-----	55	Poor Too clayey Droughty Carbonate content Depth to bedrock Low content of organic matter	0.00 0.08 0.08 0.58 0.88	Poor Depth to bedrock Shrink-swell Slope	0.00 0.12 0.32	Poor Too Clayey Slope Carbonate content Depth to bedrock	0.00 0.00 0.08 0.58
Bristow-----	45	Poor Too clayey Droughty Depth to bedrock Low content of organic matter Carbonate content Water erosion	0.00 0.00 0.00 0.12 0.46 0.90	Poor Depth to bedrock Shrink-swell Slope	0.00 0.12 0.32	Poor Too Clayey Depth to bedrock Slope Carbonate content	0.00 0.00 0.00 0.46
MaG: Mariaville-----	55	Poor Depth to bedrock Droughty Low content of organic matter Water erosion	0.00 0.00 0.88 0.90	Poor Depth to bedrock Slope	0.00 0.00	Poor Depth to bedrock Slope	0.00 0.00
Paka-----	45	Fair Low content of organic matter Water erosion	0.88 0.90	Poor Slope Depth to bedrock Shrink-swell	0.00 0.58 0.91	Poor Slope	0.00
MeE: Meadin-----	100	Poor Too sandy Droughty Low content of organic matter Too acid	0.00 0.11 0.12 0.97	Good		Poor Too sandy Rock fragments Hard to reclaim Slope	0.00 0.00 0.00 0.84

CONSTRUCTION MATERIALS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
NoC: Nora-----	100	Fair Low content of organic matter Water erosion	0.88 0.90	Fair Shrink-swell	0.87	Good	
NoD: Nora-----	100	Fair Low content of organic matter Water erosion	0.88 0.90	Fair Shrink-swell	0.87	Fair Slope	0.96
Oa: Onawa-----	100	Poor Too clayey Low content of organic matter Water erosion Carbonate content	0.00 0.12 0.90 0.97	Fair Shrink-swell	0.97	Poor Too Clayey Carbonate content	0.00 0.97
Oe: O'Neill-----	100	Fair Low content of organic matter Too acid Droughty	0.12 0.84 0.88	Good		Fair Hard to reclaim	0.92
OeC: O'Neill-----	100	Fair Low content of organic matter Too acid Droughty	0.12 0.84 0.88	Good		Fair Hard to reclaim	0.92
OfD: O'Neill-----	55	Fair Low content of organic matter Too acid Droughty	0.12 0.84 0.88	Good		Fair Hard to reclaim	0.92
Meadin-----	45	Poor Too sandy Droughty Low content of organic matter Too acid	0.00 0.11 0.12 0.97	Good		Poor Too sandy Rock fragments Hard to reclaim	0.00 0.00 0.00
On: Onita-----	100	Fair Low content of organic matter Water erosion	0.12 0.90	Fair Shrink-swell	0.81	Good	
Poorly Drained-----		Poor Low content of organic matter	0.00	Poor Slope Low strength	0.00 0.00	Poor Slope	0.00
Or: Ord-----	100	Fair Low content of organic matter Carbonate content	0.12 0.80	Fair Depth to saturated zone	0.76	Fair Depth to saturated zone	0.76
PaC: Paka-----	100	Fair Low content of organic matter Water erosion	0.88 0.90	Fair Depth to bedrock Shrink-swell	0.58 0.94	Good	
Ph: Paka-----	100	Fair Low content of organic matter Water erosion	0.88 0.90	Fair Depth to bedrock Shrink-swell	0.58 0.91	Good	

CONSTRUCTION MATERIALS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
PhC: Paka-----	100	Fair Low content of organic matter Water erosion	0.88 0.90	Fair Depth to bedrock Shrink-swell	0.58 0.91	Good	
PhD: Paka-----	100	Fair Low content of organic matter Water erosion	0.88 0.90	Fair Depth to bedrock Shrink-swell	0.58 0.91	Fair Slope	0.96
PoC: Promise-----	100	Poor Too clayey Low content of organic matter Water erosion	0.00 0.12 0.99	Poor Shrink-swell	0.00	Poor Too Clayey	0.00
RaC: Ree-----	100	Fair Low content of organic matter	0.12	Good		Good	
RaD: Ree-----	100	Fair Low content of organic matter	0.12	Good		Fair Slope	0.96
RaE: Ree-----	100	Fair Low content of organic matter	0.12	Good		Fair Slope	0.16
ReC: Reliance-----	100	Poor Too clayey Low content of organic matter Water erosion	0.00 0.50 0.90	Fair Shrink-swell	0.62	Poor Too Clayey	0.00
ReD: Reliance-----	100	Poor Too clayey Low content of organic matter Water erosion	0.00 0.50 0.90	Fair Shrink-swell	0.62	Poor Too Clayey Slope	0.00 0.96
RFC: Reliance-----	100	Poor Too clayey Low content of organic matter Water erosion	0.00 0.50 0.90	Fair Shrink-swell	0.62	Poor Too Clayey	0.00
Rw: Riverwash-----	100	Poor Too sandy Droughty Low content of organic matter	0.00 0.00 0.12	Poor Depth to saturated zone	0.00	Poor Hard to reclaim Too sandy Depth to saturated zone Rock fragments Hard to reclaim	0.00 0.00 0.00 0.00 0.32
SaG: Sansarc-----	100	Poor Too clayey Droughty Depth to bedrock Low content of organic matter Water erosion	0.00 0.00 0.00 0.60 0.99	Poor Depth to bedrock Slope Shrink-swell	0.00 0.00 0.00	Poor Slope Too Clayey Depth to bedrock	0.00 0.00 0.00

CONSTRUCTION MATERIALS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Sc: Scott-----	100	Poor Too clayey Too acid Water erosion	0.00 0.84 0.99	Poor Depth to saturated zone Shrink-swell	0.00 0.16	Poor Too Clayey Depth to saturated zone	0.00 0.00
Sm: Simeon-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.79	Good		Poor Too sandy	0.00
SuC: Simeon-----	55	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.90	Good		Poor Too sandy	0.00
Valentine-----	45	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.29	Good		Poor Too sandy	0.00
SvF2: Simeon-----	70	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.90	Fair Slope	0.92	Poor Too sandy Slope	0.00 0.00
Valentine-----	30	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.22	Fair Slope	0.92	Poor Too sandy Slope	0.00 0.00
VaE: Valentine-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.22	Good		Poor Too sandy Slope	0.00 0.37
VbB: Valentine-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.29	Good		Poor Too sandy	0.00
Ve: Verdel-----	100	Poor Too clayey Low content of organic matter	0.00 0.88	Fair Shrink-swell	0.12	Poor Too Clayey	0.00
WeC: Wewela-----	100	Fair Droughty Low content of organic matter Depth to bedrock Water erosion	0.01 0.12 0.58 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.97	Fair Depth to bedrock	0.58
zwa: Water-----	100	Not rated		Not rated		Not rated	

RECREATIONAL INTERPRETATIONS
Boyd County, Nebraska

Recreation

The soils of the survey area are rated in the following tables according to limitations that affect their suitability for recreation. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the recreational uses. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

The ratings in the tables are based on restrictive soil features, such as wetness, slope, and texture of the surface layer. Susceptibility to flooding is considered. Not considered in the ratings, but important in evaluating a site, are the location and accessibility of the area, the size and shape of the area and its scenic quality, vegetation, access to water, potential water impoundment sites, and access to public sewer lines. The capacity of the soil to absorb septic tank effluent and the ability of the soil to support vegetation also are important. Soils that are subject to flooding are limited for recreational uses by the duration and intensity of flooding and the season when flooding occurs. In planning recreational facilities, onsite assessment of the height, duration, intensity, and frequency of flooding is essential.

The information in this table can be supplemented by other information in this survey, for example, interpretations for building site development, construction materials, sanitary facilities, and water management.

Camp areas require site preparation, such as shaping and leveling the tent and parking areas, stabilizing roads and intensively used areas, and installing sanitary facilities and utility lines. Camp areas are subject to heavy foot traffic and some vehicular traffic. The ratings are based on the soil properties that affect the ease of developing camp areas and the performance of the areas after development. Slope, stoniness, and depth to bedrock or a cemented pan are the main concerns affecting the development of camp areas.

The soil properties that affect the performance of the areas after development are those that influence trafficability and promote the growth of vegetation, especially in heavily used areas. For good trafficability, the surface of camp areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Picnic areas are subject to heavy foot traffic. Most vehicular traffic is confined to access roads and parking areas. The ratings are based on the soil properties that affect the ease of developing picnic areas and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of picnic areas. For good trafficability, the surface of picnic areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Playgrounds require soils that are nearly level, are free of stones, and can withstand intensive foot traffic. The ratings are based on the soil properties that affect the ease of developing playgrounds and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of playgrounds. For good trafficability, the surface of the playgrounds should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Paths and trails for hiking and horseback riding should require little or no slope modification through cutting and filling. The ratings are based on the soil properties that affect trafficability and erodibility. These properties are stoniness, depth to a water table, ponding, flooding, slope, and texture of the surface layer.

Golf fairways are subject to heavy foot traffic and some light vehicular traffic. Cutting or filling may be required. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer. The suitability of the soil for traps, tees, roughs, and greens is not considered in the ratings.

RECREATIONAL INTERPRETATIONS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Very limited Depth to saturated zone Flooding Too clayey Restricted permeability	1.00 1.00 1.00 1.00	Very limited Too clayey Depth to saturated zone Restricted permeability	1.00 1.00 1.00	Very limited Depth to saturated zone Too clayey Restricted permeability Flooding	1.00 1.00 1.00 0.60
AnC: Anselmo-----	100	Not limited		Not limited		Somewhat limited Slope	0.50
AnD: Anselmo-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
AnF: Anselmo-----	100	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
ArF: Anselmo-----	75	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
Ba: Barney-----	100	Very limited Depth to saturated zone Flooding	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Depth to saturated zone Flooding	1.00 1.00
Bd: Blake-----	100	Very limited Flooding Restricted permeability	1.00 0.22	Somewhat limited Restricted permeability	0.22	Somewhat limited Flooding Restricted permeability	0.60 0.22
Be: Blendon-----	100	Not limited		Not limited		Not limited	
BeC: Blendon-----	100	Not limited		Not limited		Somewhat limited Slope	0.50
BoD: Boyd-----	100	Somewhat limited Too clayey Restricted permeability Slope	0.50 0.45 0.04	Somewhat limited Too clayey Restricted permeability Slope	0.50 0.45 0.04	Very limited Slope Too clayey Restricted permeability Depth to bedrock	1.00 0.50 0.45 0.42
BrG: Bristow-----	100	Very limited Slope Depth to bedrock Too clayey Restricted permeability	1.00 1.00 0.50 0.45	Very limited Slope Depth to bedrock Too clayey Restricted permeability	1.00 1.00 0.50 0.45	Very limited Slope Depth to bedrock Too clayey Restricted permeability	1.00 1.00 0.50 0.45
Bs: Brocksburg-----	100	Not limited		Not limited		Not limited	
Bt: Brocksburg-----	100	Not limited		Not limited		Not limited	
Cb: Cass-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
CrE2: Crofton-----	100	Somewhat limited Slope	0.84	Somewhat limited Slope	0.84	Very limited Slope	1.00
DuB: Dunday-----	100	Somewhat limited Too sandy	0.95	Somewhat limited Too sandy	0.95	Somewhat limited Too sandy Slope	0.95 0.00
DuC: Dunday-----	100	Somewhat limited Too sandy	0.95	Somewhat limited Too sandy	0.95	Somewhat limited Too sandy Slope	0.95 0.87
DuD: Dunday-----	100	Somewhat limited Too sandy Slope	0.95 0.04	Somewhat limited Too sandy Slope	0.95 0.04	Very limited Slope Too sandy	1.00 0.95
DxB: Dunday-----	100	Somewhat limited Too sandy	0.34	Somewhat limited Too sandy	0.34	Somewhat limited Too sandy	0.34

RECREATIONAL INTERPRETATIONS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Et: Eltree-----	100	Not limited		Not limited		Slope	0.00
Go: Grigston-----	100	Very limited Flooding	1.00	Not limited		Not limited	
GrB: Aowa-----	100	Very limited Flooding	1.00	Somewhat limited Flooding	0.40	Somewhat limited Flooding	0.60
Ha: Hall-----	100	Not limited		Not limited		Very limited Flooding	1.00
He: Haynie-----	100	Very limited Flooding	1.00	Not limited		Not limited	
IfD: Inavale-----	100	Very limited Flooding Too sandy	1.00 1.00	Not limited		Somewhat limited Flooding	0.60
IgB: Inavale-----	100	Very limited Flooding Too sandy	1.00 1.00	Very limited Too sandy	1.00	Very limited Too sandy Slope	1.00 1.00
IhB: Inavale-----	100	Very limited Flooding Too sandy	1.00 0.96	Very limited Too sandy Flooding	1.00 0.40	Very limited Too sandy Flooding Slope	1.00 1.00 0.00
In: Inavale-----	100	Very limited Flooding	1.00	Somewhat limited Too sandy	0.96	Somewhat limited Too sandy Flooding Slope	0.96 0.60 0.00
Jn: Jansen-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding Slope	0.60 0.00
JnC: Jansen-----	100	Not limited		Not limited		Not limited	
JnD: Jansen-----	100	Not limited		Not limited		Somewhat limited Slope	0.50
LaD: Labu-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
LcF: Labu-----	65	Somewhat limited Too clayey Restricted permeability Slope	0.50 0.45 0.04	Somewhat limited Too clayey Restricted permeability Slope	0.50 0.45 0.04	Very limited Slope Too clayey	1.00 0.50
						Restricted permeability Depth to bedrock	0.45 0.42
Sansarc-----	35	Very limited Slope Too clayey Restricted permeability	1.00 1.00 0.50 0.39	Very limited Slope Too clayey Restricted permeability	1.00 1.00 0.50 0.39	Very limited Slope Depth to bedrock Too clayey Restricted permeability	1.00 1.00 0.50 0.39
Le: Leshara-----	100	Very limited Flooding	1.00	Very limited Depth to bedrock Slope	1.00	Very limited Slope Depth to bedrock	1.00 1.00
		Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Too clayey Restricted permeability	0.50 0.39
LsC: Lynch-----	100	Somewhat limited Too clayey Restricted permeability	0.50 0.45	Somewhat limited Too clayey Restricted permeability	0.50 0.45	Somewhat limited Flooding	0.60
						Depth to saturated zone	0.39
LsD: Lynch-----	100	Somewhat limited Too clayey Restricted permeability Slope	0.50 0.45 0.04	Somewhat limited Too clayey Restricted permeability Slope	0.50 0.45 0.04	Somewhat limited Slope Too clayey	0.50 0.50
						Restricted permeability Depth to bedrock	0.45 0.42
						Very limited Slope Too clayey	1.00 0.50
						Restricted permeability	0.45

RECREATIONAL INTERPRETATIONS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
LyD: Lynch-----	65	Somewhat limited		Somewhat limited		Depth to bedrock	0.42
		Too clayey	0.50	Too clayey	0.50	Very limited	
		Restricted permeability	0.45	Restricted permeability	0.45	Slope	1.00
Bristow-----	35	Slope	0.04	Slope	0.04	Too clayey	0.50
						Restricted permeability	0.45
						Depth to bedrock	0.42
LyF: Lynch-----	55	Very limited		Very limited		Very limited	
		Depth to bedrock	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Too clayey	0.50	Too clayey	0.50	Slope	1.00
Bristow-----	45	Restricted permeability	0.45	Restricted permeability	0.45	Too clayey	0.50
		Slope	0.04	Slope	0.04	Restricted permeability	0.45
MaG: Mariaville-----	55	Very limited		Very limited		Very limited	
		Slope	1.00	Slope	1.00	Slope	1.00
		Too clayey	0.50	Too clayey	0.50	Too clayey	0.50
Paka-----	45	Restricted permeability	0.45	Restricted permeability	0.45	Restricted permeability	0.45
						Depth to bedrock	0.42
MeE: Meadin-----	100	Very limited		Very limited		Very limited	
		Depth to bedrock	1.00	Depth to bedrock	1.00	Slope	1.00
		Slope	1.00	Slope	1.00	Depth to bedrock	1.00
NoC: Nora-----	100	Very limited		Very limited		Too clayey	0.50
		Slope	1.00	Slope	1.00	Restricted permeability	0.45
NoD: Nora-----	100	Somewhat limited		Somewhat limited		Very limited	
		Slope	0.16	Slope	0.16	Slope	1.00
						Gravel content	0.22
Oa: Onawa-----	100	Not limited		Not limited		Somewhat limited	
						Slope	0.50
Oe: O'Neill-----	100	Somewhat limited		Somewhat limited		Very limited	
		Restricted permeability	0.04	Restricted permeability	0.04	Slope	1.00
		Slope	1.00	Slope	1.00	Gravel content	0.22
OfD: O'Neill-----	55	Very limited		Very limited		Somewhat limited	
		Flooding	1.00	Restricted permeability	1.00	Restricted permeability	0.40
		Restricted permeability	1.00	Too clayey	1.00	Too clayey	1.00
On: Onita-----	100	Too clayey	1.00			Flooding	0.60
Or: Ord-----	100	Not limited		Not limited		Not limited	
						Somewhat limited	
						Slope	0.50
PaC: Paka-----	100	Not limited		Not limited		Very limited	
						Slope	1.00
						Gravel content	0.22
Ph: Paka-----	100	Not limited		Not limited		Somewhat limited	
						Flooding	0.60
						Depth to saturated zone	0.07

RECREATIONAL INTERPRETATIONS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
PhC: Paka-----	100	Not limited		Not limited		Somewhat limited Slope	0.50
PhD: Paka-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
PoC: Promise-----	100	Somewhat limited Too clayey Restricted permeability	0.50 0.45	Somewhat limited Too clayey Restricted permeability	0.50 0.45	Somewhat limited Slope Too clayey Restricted permeability	0.50 0.50 0.45
RaC: Ree-----	100	Not limited		Not limited		Somewhat limited Slope	0.50
RaD: Ree-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
RaE: Ree-----	100	Somewhat limited Slope	0.84	Somewhat limited Slope	0.84	Very limited Slope	1.00
ReC: Reliance-----	100	Somewhat limited Restricted permeability	0.40	Somewhat limited Restricted permeability	0.40	Somewhat limited Slope Restricted permeability	0.50 0.40
ReD: Reliance-----	100	Somewhat limited Restricted permeability Slope	0.40 0.04	Somewhat limited Restricted permeability Slope	0.40 0.04	Very limited Slope Restricted permeability	1.00 0.40
RfC: Reliance-----	100	Not limited		Not limited		Somewhat limited Slope	0.50
Rw: Riverwash-----	100	Very limited Depth to saturated zone Flooding Too sandy	1.00 1.00 1.00	Very limited Too sandy Depth to saturated zone Flooding	1.00 1.00 0.40	Very limited Depth to saturated zone Too sandy Flooding	1.00 1.00 1.00
SaG: Sansarc-----	100	Very limited Slope Depth to bedrock Too clayey Restricted permeability	1.00 1.00 0.50 0.39	Very limited Slope Depth to bedrock Too clayey Restricted permeability	1.00 1.00 0.50 0.39	Very limited Slope Depth to bedrock Too clayey Restricted permeability	1.00 1.00 0.50 0.39
Sc: Scott-----	100	Very limited Depth to saturated zone Restricted permeability	1.00 1.00	Very limited Depth to saturated zone Restricted permeability	1.00 1.00	Very limited Depth to saturated zone Restricted permeability	1.00 1.00
Sm: Simeon-----	100	Somewhat limited Too sandy	0.79	Somewhat limited Too sandy	0.79	Somewhat limited Too sandy	0.79
SuC: Simeon-----	55	Somewhat limited Too sandy	0.79	Somewhat limited Too sandy	0.79	Somewhat limited Too sandy Slope	0.79 0.13
Valentine-----	45	Somewhat limited Too sandy	0.87	Somewhat limited Too sandy	0.87	Somewhat limited Too sandy Slope	0.87 0.13
SvF2: Simeon-----	70	Very limited Slope Too sandy	1.00 0.79	Very limited Slope Too sandy	1.00 0.79	Very limited Slope Too sandy	1.00 0.79
Valentine-----	30	Very limited Too sandy Slope	1.00 1.00	Very limited Too sandy Slope	1.00 1.00	Very limited Too sandy Slope	1.00 1.00
VaE: Valentine-----	100	Very limited Too sandy Slope	1.00 0.63	Very limited Too sandy Slope	1.00 0.63	Very limited Too sandy Slope	1.00 1.00
VbB: Valentine-----	100	Somewhat limited Too sandy	0.87	Somewhat limited Too sandy	0.87	Somewhat limited Too sandy Slope	0.87 0.00

RECREATIONAL INTERPRETATIONS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ve: Verdel-----	100	Somewhat limited Too clayey Restricted permeability	0.50 0.45	Somewhat limited Too clayey Restricted permeability	0.50 0.45	Somewhat limited Too clayey Restricted permeability	0.50 0.45
WeC: Wewela-----	100	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45	Somewhat limited Slope Restricted permeability Depth to bedrock	0.50 0.45 0.42
zwa: Water-----	100	Not rated		Not rated		Not rated	

RECREATIONAL INTERPRETATIONS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Very limited Depth to saturated zone Too clayey	1.00 1.00	Very limited Too clayey Depth to saturated zone Flooding	1.00 1.00 0.60
AnC: Anselmo-----	100	Not limited		Not limited	
AnD: Anselmo-----	100	Not limited		Somewhat limited Slope	0.04
AnF: Anselmo-----	100	Not limited		Somewhat limited Slope	0.96
ArF: Anselmo-----	75	Not limited		Somewhat limited Slope	0.96
Rock Outcrop-----	25	Not rated		Not rated	
Ba: Barney-----	100	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone Droughty	1.00 1.00 0.13
Bd: Blake-----	100	Not limited		Somewhat limited Flooding	0.60
Be: Blendon-----	100	Not limited		Not limited	
BeC: Blendon-----	100	Not limited		Not limited	
BoD: Boyd-----	100	Somewhat limited Too clayey	0.50	Very limited Too clayey Depth to bedrock Droughty Slope	1.00 0.42 0.34 0.04
BrG: Bristow-----	100	Very limited Slope Too clayey	1.00 0.50	Very limited Depth to bedrock Slope Droughty Too clayey	1.00 1.00 1.00 1.00
Bs: Brocksburg-----	100	Not limited		Not limited	
Bt: Brocksburg-----	100	Not limited		Not limited	
Cb: Cass-----	100	Not limited		Somewhat limited Flooding	0.60
CrE2: Crofton-----	100	Not limited		Somewhat limited Slope	0.84
DuB: Dunday-----	100	Somewhat limited Too sandy	0.95	Somewhat limited Droughty	0.07
DuC: Dunday-----	100	Somewhat limited Too sandy	0.95	Somewhat limited Droughty	0.07
DuD: Dunday-----	100	Somewhat limited Too sandy	0.95	Somewhat limited Droughty Slope	0.07 0.04
DxB: Dunday-----	100	Somewhat limited Too sandy	0.34	Somewhat limited Droughty	0.02
Et: Eltree-----	100	Not limited		Not limited	
Go: Grigston-----	100	Not limited		Somewhat limited Flooding	0.60
GrB: Aowa-----	100	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
Ha: Hall-----	100	Not limited		Not limited	

RECREATIONAL INTERPRETATIONS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
He: Haynie-----	100	Not limited		Somewhat limited Flooding	0.60
IfD: Inavale-----	100	Very limited Too sandy	1.00	Somewhat limited Droughty	0.29
IgB: Inavale-----	100	Very limited Too sandy Flooding	1.00 0.40	Very limited Flooding Droughty	1.00 0.29
IhB: Inavale-----	100	Somewhat limited Too sandy	0.96	Somewhat limited Flooding Droughty	0.60 0.10
In: Inavale-----	100	Not limited		Somewhat limited Flooding	0.60
Jn: Jansen-----	100	Not limited		Not limited	
JnC: Jansen-----	100	Not limited		Not limited	
JnD: Jansen-----	100	Not limited		Somewhat limited Slope	0.04
LaD: Labu-----	100	Somewhat limited Too clayey	0.50	Very limited Too clayey Depth to bedrock Droughty Slope	1.00 0.42 0.20 0.04
LcF: Labu-----	65	Somewhat limited Slope Too clayey	0.68 0.50	Very limited Too clayey Slope Depth to bedrock Droughty	1.00 1.00 0.42 0.20
Sansarc-----	35	Somewhat limited Slope Too clayey	0.68 0.50	Very limited Too clayey Depth to bedrock Droughty Slope	1.00 1.00 1.00 1.00
Le: Leshara-----	100	Not limited		Somewhat limited Flooding Depth to saturated zone	0.60 0.19
LsC: Lynch-----	100	Somewhat limited Too clayey	0.50	Very limited Too clayey Depth to bedrock Droughty	1.00 0.42 0.09
LsD: Lynch-----	100	Somewhat limited Too clayey	0.50	Very limited Too clayey Depth to bedrock Droughty Slope	1.00 0.42 0.09 0.04
LyD: Lynch-----	65	Somewhat limited Too clayey	0.50	Very limited Too clayey Depth to bedrock Droughty Slope	1.00 0.42 0.09 0.04
Bristow-----	35	Somewhat limited Too clayey	0.50	Very limited Depth to bedrock Droughty Too clayey Slope	1.00 1.00 1.00 0.04
LyF: Lynch-----	55	Somewhat limited Slope Too clayey	0.68 0.50	Very limited Slope Too clayey Depth to bedrock Droughty	1.00 1.00 0.42 0.09
Bristow-----	45	Somewhat limited Slope Too clayey	0.68 0.50	Very limited Depth to bedrock Droughty Slope Too clayey	1.00 1.00 1.00 1.00
MaG: Mariaville-----	55	Very limited		Very limited	

RECREATIONAL INTERPRETATIONS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
		Slope	1.00	Depth to bedrock	1.00
				Slope	1.00
				Droughty	0.43
Paka-----	45	Very limited		Very limited	
MeE:		Slope	1.00	Slope	1.00
Meadin-----	100	Not limited		Somewhat limited	
				Droughty	0.63
				Slope	0.16
NoC:				Not limited	
Nora-----	100	Not limited			
NoD:				Somewhat limited	
Nora-----	100	Not limited		Slope	0.04
Oa:					
Onawa-----	100	Very limited		Very limited	
		Too clayey	1.00	Too clayey	1.00
				Flooding	0.60
Oe:					
O'Neill-----	100	Not limited		Not limited	
OeC:					
O'Neill-----	100	Not limited		Not limited	
OFD:					
O'Neill-----	55	Not limited		Not limited	
Meadin-----	45	Not limited		Somewhat limited	
				Droughty	0.63
On:					
Onita-----	100	Not limited		Not limited	
Poorly Drained-----		Very limited		Very limited	
		Slope	1.00	Slope	1.00
		Water erosion	1.00		
Or:					
Ord-----	100	Not limited		Somewhat limited	
				Flooding	0.60
				Depth to	0.03
				saturated zone	
PaC:					
Paka-----	100	Not limited		Not limited	
Ph:					
Paka-----	100	Not limited		Not limited	
PhC:					
Paka-----	100	Not limited		Not limited	
PhD:					
Paka-----	100	Not limited		Somewhat limited	
				Slope	0.04
PoC:					
Promise-----	100	Somewhat limited		Very limited	
		Too clayey	0.50	Too clayey	1.00
RaC:					
Ree-----	100	Not limited		Not limited	
RaD:					
Ree-----	100	Not limited		Somewhat limited	
				Slope	0.04
RaE:					
Ree-----	100	Not limited		Somewhat limited	
				Slope	0.84
ReC:					
Reliance-----	100	Not limited		Not limited	
ReD:					
Reliance-----	100	Not limited		Somewhat limited	
				Slope	0.04
RfC:					
Reliance-----	100	Not limited		Not limited	
Rw:					
Riverwash-----	100	Very limited		Very limited	
		Depth to	1.00	Flooding	1.00
		saturated zone			
		Too sandy	1.00	Depth to	1.00
				saturated zone	
		Flooding	0.40	Droughty	1.00
SaG:					
Sansarc-----	100	Very limited		Very limited	
		Slope	1.00	Too clayey	1.00
		Too clayey	0.50	Depth to bedrock	1.00
				Slope	1.00
				Droughty	1.00
Sc:					
Scott-----	100	Very limited		Very limited	
		Depth to	1.00	Depth to	1.00
		saturated zone		saturated zone	

RECREATIONAL INTERPRETATIONS--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Sm: Simeon-----	100	Somewhat limited Too sandy	0.79	Somewhat limited Droughty	0.22
SuC: Simeon-----	55	Somewhat limited Too sandy	0.79	Somewhat limited Droughty	0.07
Valentine-----	45	Somewhat limited Too sandy	0.87	Somewhat limited Droughty	0.61
SvF2: Simeon-----	70	Somewhat limited Too sandy Slope	0.79 0.08	Very limited Slope Droughty	1.00 0.07
Valentine-----	30	Very limited Too sandy Slope	1.00 0.08	Very limited Slope Droughty	1.00 0.73
VaE: Valentine-----	100	Very limited Too sandy	1.00	Somewhat limited Droughty Slope	0.73 0.63
VbB: Valentine-----	100	Somewhat limited Too sandy	0.87	Somewhat limited Droughty	0.61
Ve: Verdel-----	100	Somewhat limited Too clayey	0.50	Very limited Too clayey	1.00
WeC: Wewela-----	100	Not limited		Somewhat limited Depth to bedrock Droughty	0.42 0.32
zwa: Water-----	100	Not rated		Not rated	

WILDLIFE INTERPRETATIONS Boyd County, Nebraska

Use and Explanation of Wildlife Interpretations

Soils directly affect the kind and amount of vegetation that is available to wildlife as food and cover. They also affect the development of water impoundments. The kind and abundance of wildlife that populate an area depend largely on the amount and distribution of food, cover, water, and living space. If any one of these elements is missing, inadequate, or inaccessible, wildlife will be scarce or will not inhabit the area. If the soils have the potential, wildlife habitat can be created or improved by planting appropriate vegetation, properly managing the existing plant cover, and fostering the natural establishment of desirable plants.

In the Wildlife Interpretations table, the soils in the survey area are rated according to their potential for providing habitat for various kinds of wildlife. This information can be used in planning parks, wildlife refuges, nature study areas, and other developments for wildlife; in selecting soils that are suitable for establishing, improving, or maintaining specific elements of wildlife habitat; and in determining the intensity of management needed for each element of the habitat.

Suitability Ratings

The potential of the soil is rated good, fair, poor, or very poor.

Good - means that the element of wildlife habitat or the kind of habitat is easily created, improved, or maintained. Few or no limitations affect management, and satisfactory results can be expected if the soil is used for the designated purpose.

Fair - means that the element of wildlife habitat or kind of habitat can be created, improved, or maintained in most places. Moderately intensive management is required for satisfactory results.

Poor - means that limitations are severe for the designated element or kind of wildlife habitat. Habitat can be created, improved, or maintained in most places, but management is difficult and requires intensive effort.

Very Poor - means that limitations are very severe for the designated element or kind of wildlife habitat. Habitat is difficult to create, improve, or maintain in most places, and management is difficult and requires intensive effort.

Description of Wildlife Habitat Elements

Openland habitat consists of croplands, pastures, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. The kind of wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, killdeer, cottontail rabbit, red fox, and coyote.

Woodland habitat consists of hardwood or conifers, or a mixture of these and associated grasses, legumes and wild herbaceous plants. Examples of wildlife attracted to this habitat are wild turkey, thrushes, woodpeckers, owl, tree squirrels, raccoon, and deer.

Wetland habitat consists of water-tolerant plants in open, marshy or swampy, shallow water areas. Examples of wildlife attracted to this habitat are ducks, geese, herons, bitterns, rails, kingfishers, shorebirds, muskrat, mink, and beaver.

The elements of wildlife habitat are described in the following paragraphs.

Grain and seed crops are domestic grains and seed-producing herbaceous plants. Soil properties and features that affect the growth of grain and seed crops are depth of the root zone, texture of the surface layer, available water capacity, wetness, slope, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of grain and seed crops are corn, wheat, oats, and barley.

Grasses and legumes are domestic perennial grasses and herbaceous legumes. Soil properties and features that affect the growth of grasses and legumes are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, flooding, and slope. Soil temperature and soil moisture also are considerations. Examples of grasses and legumes are fescue, lovegrass, bromegrass, clover, and alfalfa.

Wild herbaceous plants are native or naturally established grasses and forbs, including weeds. Soil properties and features that affect the growth of these plants are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of wild herbaceous plants are bluestem, goldenrod, beggarweed, wheatgrass, and grama.

Hardwood trees and woody understory produce nuts or other fruit, buds, catkins, twigs, bark, and foliage. Soil properties and features that affect the growth of hardwood trees and shrubs are depth of the root zone, available water capacity, and wetness. Examples of these plants are oak, poplar, cherry, sweetgum, apple, hawthorn, dogwood, hickory, blackberry, and blueberry. Examples of fruit-producing shrubs that are suitable for planting on soils rated good are Russian-olive, autumn-olive, and crabapple.

Coniferous plants furnish browse and seeds. Soil properties and features that affect the growth of coniferous trees, shrubs, and ground cover are depth of the root zone, available water capacity, and wetness. Examples of coniferous plants are pine, spruce, fir, cedar, and juniper.

Shrubs are bushy woody plants that produce fruit, buds, twigs, bark, and foliage. Soil properties and features that affect the growth of shrubs are depth of the root zone, available water capacity, salinity, and soil moisture. Examples of shrubs are fragrant sumac, chokecherry, American plum, sand plum, and gorden currant.

Wetland plants are annual and perennial wild herbaceous plants that grow on moist or wet sites. Submerged or floating aquatic plants are excluded. Soil properties and features affecting wetland plants are texture of the surface layer, wetness, reaction, salinity, slope, and surface stoniness. Examples of wetland plants are smartweed, wild millet, saltgrass, cordgrass, rushes, sedges, and cattails.

Shallow water areas have an average depth of less than 5 feet. Some are naturally wet areas. Others are created by dams, levees, or other water-control structures. Soil properties and features affecting shallow water areas are depth to bedrock, wetness, surface stoniness, slope, and permeability. Examples of shallow water areas are marshes, waterfowl feeding areas, and ponds.

The habitat for various kinds of wildlife is described in the following paragraphs.

Habitat for openland wildlife consists of cropland, pasture, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. Wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, cottontail, red fox and coyote.

Habitat for woodland wildlife consists of areas of deciduous and/or coniferous plants and associated grasses, legumes, and wild herbaceous plants. Wildlife attracted to these areas include wild turkey, thrushes, woodpeckers, squirrels, gray fox, raccoon, and deer.

Habitat for wetland wildlife consists of open, marshy or swampy shallow water areas. Some of the wildlife attracted to such areas are ducks, geese, herons, shore birds, muskrat, mink, and beaver.

Habitat for rangeland wildlife consists of areas of shrubs and wild herbaceous plants. Wildlife attracted to rangeland include antelope, deer, cottontail rabbit, prairie chicken, meadowlark, quail, and pheasant.

WILDLIFE INTERPRETATIONS
Boyd County, Nebraska

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
Ab: ALBATON-----	Fair	Fair	Fair	Poor	Very poor	---	Good	Good	Fair	Poor	Good	---
AnC: ANSELMO-----	Fair	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
AnD: ANSELMO-----	Fair	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
AnF: ANSELMO-----	Poor	Fair	Good	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor	Good
ArF: ANSELMO-----	Poor	Fair	Good	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor	Good
ROCK OUTCROP----	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor
Ba: BARNEY-----	Very poor	Poor	Fair	Poor	Poor	Poor	Good	Good	Poor	Fair	Good	Fair
Bd: BLAKE-----	Good	Good	Good	Good	Good	---	Good	Good	Good	Good	Good	---
Be: BLENDON-----	Fair	Fair	Good	Fair	Very poor	---	Very poor	Very poor	Fair	Very poor	Very poor	Good
BeC: BLENDON-----	Fair	Fair	Good	Fair	Very poor	---	Very poor	Very poor	Fair	Very poor	Very poor	Good
BoD: BOYD-----	Poor	Fair	Good	Fair	Very poor	---	Very poor	Very poor	Poor	Very poor	Very poor	Good
BrG: BRISTOW-----	Very poor	Very poor	Poor	---	---	Poor	Very poor	Very poor	Very poor	---	Very poor	Poor
Bs: BROCKSBURG-----	Good	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
Bt: BROCKSBURG-----	Good	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
Cb: CASS-----	Good	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
CrE2: CROFTON-----	Fair	Good	Good	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor	Good
DuB: DUNDAY-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Good
DuC: DUNDAY-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Good
DuD: DUNDAY-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Good
DxB: DUNDAY-----	Fair	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor	Good
Et: ELTREE-----	Good	Good	Fair	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
Go: GRIGSTON-----	Good	Good	Good	---	---	Fair	Poor	Fair	Good	---	Poor	Fair

WILDLIFE INTERPRETATIONS--Continued
Boyd County, Nebraska

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
GrB: AOWA-----	Poor	Fair	Poor	---	---	Fair	Poor	Very poor	Poor	---	Very poor	Poor
Ha: HALL-----	Good	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
He: HAYNIE-----	Good	Good	Good	Good	Good	---	Poor	Poor	Good	Good	Poor	---
IfD: INVALE-----	Poor	Fair	Good	Fair	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor	Fair
IgB: INVALE-----	Very poor	Poor	Fair	Fair	Fair	Fair	Very poor	Very poor	Poor	Fair	Very poor	Fair
IhB: INVALE-----	Fair	Fair	Good	Fair	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor	Good
In: INVALE-----	Fair	Fair	Good	Fair	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor	Good
Jn: JANSEN-----	Good	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
JnC: JANSEN-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
JnD: JANSEN-----	Poor	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
LaD: LABU-----	Fair	Good	Fair	Fair	Fair	Fair	Poor	Very poor	Fair	Fair	Very poor	Fair
LcF: LABU-----	Poor	Fair	Fair	Fair	Fair	Fair	Very poor	Very poor	Poor	Fair	Very poor	Fair
SANSARC-----	Very poor	Very poor	Fair	Poor	Very poor	Fair	Very poor	Very poor	Very poor	Very poor	Very poor	Fair
Le: LESHARA-----	Good	Good	Good	Good	Good	Good	Fair	Fair	Good	Good	Fair	Good
LsC: LYNCH-----	Fair	Fair	Poor	---	Good	Good	Very poor	Very poor	Fair	---	Very poor	Fair
LsD: LYNCH-----	Fair	Fair	Poor	---	Good	Good	Very poor	Very poor	Fair	---	Very poor	Fair
LyD: LYNCH-----	Fair	Fair	Poor	---	Good	Good	Very poor	Very poor	Fair	---	Very poor	Fair
BRISTOW-----	Very poor	Very poor	Poor	---	---	Poor	Very poor	Very poor	Very poor	---	Very poor	Poor
LyF: LYNCH-----	Poor	Fair	Poor	---	Good	Good	Very poor	Very poor	Poor	---	Very poor	Fair
BRISTOW-----	Very poor	Very poor	Poor	---	---	Poor	Very poor	Very poor	Very poor	---	Very poor	Poor
MaG: MARIAVILLE-----	Poor	Poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Poor	Fair	Very poor	Poor
PAKA-----	Poor	Fair	Good	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor	Good

WILDLIFE INTERPRETATIONS--Continued
Boyd County, Nebraska

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
MeE: MEADIN-----	Very poor	Poor	Fair	Poor	Poor	Fair	Very poor	Very poor	Poor	Poor	Very poor	Fair
NoC: NORA-----	Good	Good	Good	Good	Very poor	---	Very poor	Very poor	Good	Very poor	Very poor	Good
NoD: NORA-----	Fair	Good	Good	Good	Very poor	---	Very poor	Very poor	Good	Very poor	Very poor	Good
Oa: ONAWA-----	Fair	Fair	Fair	Poor	Very poor	---	Good	Good	Fair	Poor	Good	---
Oe: O'NEILL-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
OeC: O'NEILL-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
OfD: O'NEILL-----	Poor	Fair	Good	Fair	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor	Fair
MEADIN-----	Very poor	Poor	Fair	Poor	Poor	Fair	Very poor	Very poor	Poor	Poor	Very poor	Fair
On: ONITA-----	Good	Good	Good	Good	Very poor	Very poor	Very poor	Very poor	Good	Very poor	Very poor	Good
POORLY DRAINED--	---	---	---	---	---	---	---	---	---	---	---	---
Or: ORD-----	Good	Good	Good	Good	Good	Good	Fair	Fair	Good	Good	Fair	Good
PaC: PAKA-----	Fair	Good	Good	Good	Good	Good	Poor	Very poor	Fair	Good	Very poor	Good
Ph: PAKA-----	Good	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor	Good
PhC: PAKA-----	Fair	Good	Good	Good	Good	Good	Poor	Very poor	Fair	Good	Very poor	Good
PhD: PAKA-----	Fair	Good	Good	Good	Good	Good	Poor	Very poor	Fair	Good	Very poor	Good
PoC: PROMISE-----	Fair	Fair	Good	Fair	Very poor	Very poor	Very poor	Very poor	Fair	Very poor	Very poor	Good
RaC: REE-----	Good	Good	Good	Good	Very poor	Very poor	Very poor	Very poor	Good	Very poor	Very poor	Good
RaD: REE-----	Fair	Good	Good	Fair	Very poor	Very poor	Very poor	Very poor	Fair	Very poor	Very poor	Good
RaE: REE-----	Poor	Good	Good	Poor	Very poor	Very poor	Very poor	Very poor	Poor	Very poor	Very poor	Good
ReC: RELIANCE-----	Good	Good	Good	Good	Very poor	Very poor	Very poor	Very poor	Good	Very poor	Very poor	Good
ReD: RELIANCE-----	Fair	Good	Good	Fair	Very poor	Very poor	Very poor	Very poor	Fair	Very poor	Very poor	Good
RfC: RELIANCE-----	Good	Good	Good	Good	Very poor	Very poor	Very poor	Very poor	Good	Very poor	Very poor	Good

WILDLIFE INTERPRETATIONS--Continued
Boyd County, Nebraska

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
Rw: RIVERWASH-----	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Poor	Very poor	Very poor	Poor	Very poor
SaG: SANSARC-----	Very poor	Very poor	Fair	Poor	Very poor	Fair	Very poor	Very poor	Very poor	Very poor	Very poor	Fair
Sc: SCOTT-----	Poor	Fair	Fair	Fair	Fair	Poor	Good	Good	Fair	Fair	Good	Fair
Sm: SIMEON-----	Poor	Poor	Fair	Poor	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor	Fair
SuC: SIMEON-----	Poor	Poor	Fair	Poor	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor	Fair
VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
SvF2: SIMEON-----	Poor	Poor	Fair	Poor	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor	Fair
VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
VaE: VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
VbB: VALENTINE-----	Fair	Good	Fair	Poor	Fair	Fair	Very poor	Very poor	Fair	Poor	Very poor	Fair
Ve: VERDEL-----	Good	Good	Fair	Good	Good	Fair	Poor	Very poor	Good	Good	Very poor	Fair
WeC: WEWELA-----	Fair	Fair	Good	Fair	Very poor	---	Very poor	Very poor	Fair	Very poor	Very poor	Good
zwa: WATER-----	---	---	---	---	---	---	---	---	---	---	---	---

YIELDS PER ACRE OF PASTURE AND HAYLAND
Boyd County, Nebraska

Use and Explanation of Pastureland and Hayland Interpretations

This subsection provides information concerning the suitability of soils for the production of pasture and hayland. This subsection may contain pasture and hayland suitability groupings, land capability and yield estimates, yield estimates for individual grasses or legumes, or other information pertaining to the production of forage.

Pasture and Hayland Suitability Groupings

Soils are placed in pasture and hayland groups according to their suitability for the production of forage. The soils in each group are enough alike to be suited to the same grasses or legumes, to have similar limitations and hazards, to require similar management, and to have similar productivity and other responses to management. Thus, the pasture and hayland suitability group is a convenient way of grouping the soils for their management. If used, these groupings are identified and described in other reports in the subsection.

Yield Estimates

The average yields per acre that can be expected of the principal pasture or hayland crops, under a high level of management, are presented in this subsection. In any given year, yields may be higher or lower than those indicated in the tables because of variations in rainfall or other climatic factors. The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations are also considered.

Under good management, proper grazing is essential for the production of high quality forage, stand survival, and erosion control. Proper grazing helps plants maintain sufficient and generally vigorous top growth during the growing season. Brush control is essential in many areas, and weed control generally is needed. Rotation grazing and renovation are also important management practices.

The Pasture and Hayland table show yield estimates in tons per acre and animal unit months for pasture and hayland groups. An animal unit month is the amount of forage required by one animal unit (AU) for 30 days. On animal unit (AU) is one (1000 pound) mature cow and a calf up to weaning age (usually six months of age) or their equivalent. The Natural Resources Conservation Service uses 900 pounds of air dry forage as the amount needed to meet this requirement. To maintain a healthy and vigorous plant community, the degree of use should never be greater than 50 percent. Therefore only 25 percent of the total biomass grown is considered consumed by the grazing animal. Animal Unit Months can be converted to air dry pounds per acre production by multiplying the AUM by 30 days, then by 30 pounds per day, and then by four. This figure is the amount of total forage production.

Planners of management systems for individual fields or farms should consider the detailed information given in the description of each soil in the Nontechnical Description section. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
Ab: Albaton-----	3w	3w	3.00	4.50
AnC: Anselmo-----	3e	3e	2.00	5.00
AnD: Anselmo-----	4e	4e	1.30	4.00
AnF: Anselmo-----	6e	6e	1.20	3.00
ArF: Anselmo-----	6e	6e	1.00	2.50
Rock Outcrop-----	8s	---	1.00	2.50
Ba: Barney-----	5w	---	---	---
Bd: Blake-----	2w	2w	3.00	6.00
Be: Blendon-----	2e	2e	2.00	5.50
BeC: Blendon-----	3e	3e	1.90	5.00
BoD: Boyd-----	4e	---	1.00	---
BrG: Bristow-----	7s	---	---	---
Bs: Brocksburg-----	2e	2e	1.30	4.50
Bt: Brocksburg-----	2s	2s	1.50	5.00
Cb: Cass-----	2w	2w	1.90	5.50
CrE2: Crofton-----	4e	---	1.00	---
DuB: Dunday-----	4e	3e	1.20	4.50
DuC: Dunday-----	4e	4e	1.20	4.00
DuD: Dunday-----	6e	4e	1.00	2.00
DxB: Dunday-----	4e	3e	1.60	5.00
Et: Eltree-----	2c	1	2.00	6.00
Go: Grigston-----	2w	2w	2.20	6.00
GrB: Aowa-----	6w	---	---	---
Ha: Hall-----	2c	1	2.20	6.00
He: Haynie-----	1	1	3.00	5.50
IfD: Inavale-----	6e	4e	---	---
IgB: Inavale-----	6w	---	---	---

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
IhB: Inavale-----	4e	3e	1.50	4.00
In: Inavale-----	3e	3e	1.50	4.50
Jn: Jansen-----	2s	2s	1.50	5.00
JnC: Jansen-----	3e	3e	1.30	4.50
JnD: Jansen-----	4e	4e	0.60	4.00
LaD: Labu-----	4e	4e	1.00	---
LcF: Labu-----	6e	---	---	---
Sansarc-----	6s	---	---	---
Le: Leshara-----	2w	2w	2.50	5.00
LsC: Lynch-----	3e	3e	1.00	---
LsD: Lynch-----	4e	4e	0.80	---
LyD: Lynch-----	4e	4e	0.80	---
Bristow-----	6s	---	0.80	---
LyF: Lynch-----	6e	---	---	---
Bristow-----	6s	---	---	---
MaG: Mariaville-----	6s	---	---	---
Paka-----	6e	---	---	---
MeE: Meadin-----	6s	---	---	---
NoC: Nora-----	2e	3e	1.80	5.00
NoD: Nora-----	3e	4e	1.50	4.00
Oa: Onawa-----	2w	2w	3.00	4.50
Oe: O'Neill-----	3e	3e	1.00	4.00
OeC: O'Neill-----	4e	4e	0.80	3.50
OfD: O'Neill-----	6e	4e	0.80	3.00
Meadin-----	6s	---	0.80	3.00
On: Onita-----	2c	1	2.00	5.50
Poorly Drained-----	---	---	---	---
Or: Ord-----	2w	2w	2.20	5.00
PaC: Paka-----	3e	3e	1.80	5.00

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
Ph: Paka-----	2c	1	2.00	5.50
PhC: Paka-----	3e	3e	1.80	5.00
PhD: Paka-----	4e	4e	1.50	---
PoC: Promise-----	3e	---	1.20	---
RaC: Ree-----	3e	3e	1.60	4.50
RaD: Ree-----	4e	4e	1.40	4.00
RaE: Ree-----	6e	---	1.00	---
ReC: Reliance-----	3e	3e	1.80	5.00
ReD: Reliance-----	4e	4e	1.40	4.00
RfC: Reliance-----	3e	3e	1.60	4.80
Rw: Riverwash-----	8w	---	---	---
SaG: Sansarc-----	7s	---	---	---
Sc: Scott-----	4w	---	---	---
Sm: Simeon-----	6s	4s	---	3.00
SuC: Simeon-----	6s	4s	---	3.10
Valentine-----	6e	4e	---	3.10
SvF2: Simeon-----	6s	---	---	---
Valentine-----	6e	---	---	---
VaE: Valentine-----	6e	---	---	---
VbB: Valentine-----	4e	4e	---	3.40
Ve: Verdel-----	2s	2s	1.60	5.00
WeC: Wewela-----	3e	3e	1.20	---
zwa: Water-----	---	---	---	---

CONSERVATION TREE AND SHRUB MANAGEMENT
Boyd County, Nebraska

A Conservation Tree/Shrub Suitability Group (CTSG), formerly Windbreak Suitability Group, is a physiographic unit or area having similar climatic and edaphic characteristics that control the selection and height growth of trees and shrubs.

In this table, the Conservation Tree and Shrub Grouping is expressed as a group index number. The group index for Conservation Tree and Shrub groups (CTSG) are a guide for species best suited for different kinds of soil and for prediction height, growth, and effectiveness. The groupings can be used when selection woody plants for windbreaks, wildlife plantings riparian buffers, reforestation, other environmental plantings, recreation, landscaping, wetland restoration or enhancement and critical area plantings. CTSG's are developed to assure satisfactory species selection and adaptation to specific conditions of soil, climate and physiography. CTSG's are a guide for selection species best suited for different kinds of soil and prediction height growth and effectiveness.

All soil series mapped in the state have been placed in 10 groups of similar soil characteristics. Groups 1, 2, 3, 4, 6, and 9 are further divided into subgroups. In addition, all groups provide information by Major Land Resource Areas.

Each tree or shrub species has certain climatic and physiographic limits. Within these parameters a tree or shrub may be well or poorly suited because of soil characteristics. Each tree or shrub also has definable potentials of height growth depending on the factors just mentioned. Accurate definitions of potential heights are necessary for proper windbreak planning and design.

Windbreaks protect livestock, buildings, roads and yards from wind and snow. They also protect fruit trees and gardens, and they furnish habitat for wildlife. Several rows of low-growing and high-growing broadleaf and coniferous trees and shrubs provide the most protection.

Field windbreaks are narrow plantings made at right angles to the prevailing wind and at specific intervals across the field. The interval depends on the erodibility of the soil. Field windbreaks protect cropland and crops from wind, help to keep snow on the fields, and provide food and cover for wildlife.

Environmental plantings help to beautify and screen houses and other buildings and to abate noise. The plants, mostly evergreen shrubs and trees, are closely spaced. To ensure plant survival, a healthy planting stock of suitable species should be planted properly on a well prepared site and maintained in good condition.

Windbreaks are often planted on land that did not grow trees originally. Knowledge of how trees perform on such land can be gained only by observing and recording their performance where trees have been planted and survived. The problem is compounded by the fact that many favorite windbreak species are not indigenous to the areas in which they are planted.

The Kansas Field Office Technical Guide Notice KS-230, Conservation Tree and Shrub Plantings Suitability Groups shows the adapted species listing for each group index number. Showing the height that locally grown trees and shrubs are expected to reach in 20 years on various soils. The estimates are based on measurements and observation of established plantings that have been given adequate care. This information should be used to determine the placement of a windbreak, the area protected and the arrangement of species.

A number of attributes are included in the CTSG species tables for each group number found in this section of the Field Office Technical Guide. These attributes were rated subjectively and assigned a relative value to further assist those unfamiliar with individual species characteristics or desirability for the intended use. Definitions and explanations can be found. Additional information on planning windbreaks and screens and planting and caring for trees and shrubs can be obtained from the local office of the Natural Resources Conservation Service or of the Cooperative Extension Service or from a commercial nursery. See part 537 of the National Forestry Manual for additional information.

In the Tree and Shrub Management table interpretive ratings are given for various aspects of forest and conservation tree and shrub management. Some rating class terms indicate the degree to which the soils are suited to a specified forest management practice. Well suited indicates that the soil has features that are favorable for the specified practice and has no limitations. Good performance can be expected, and little or no maintenance is needed. Moderately well suited indicates that the soil has features that are moderately favorable for the specified practice. One or more soil properties are less than desirable and fair performance can be expected. Some maintenance is needed. Poorly suited indicates that the soil has one or more properties that are unfavorable for the specified practice. Overcoming the unfavorable properties requires special design, extra maintenance, and costly alteration. Unsited indicates that the expected performance of the soil is unacceptable for the specified practice or that extreme measures are needed to overcome the undesirable soil properties.

The paragraphs that follow indicate the soil properties considered in rating the soils for forest and conservation tree and shrub management practices. More detailed information about the criteria used in the ratings is available in the "National Forestry Manual," which is available in local offices of the Natural Resources Conservation Service or on the Internet. Also, in the Kansas Field Office Technical Guide Notice KS-230, Conservation Tree and Shrub Plantings Suitability Groups.

Ratings in the columns suitability for hand planting and suitability for mechanical planting are based on slope, depth to a restrictive layer, content of sand, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, moderately well suited, poorly suited, or unsited to these methods of planting. It is assumed that necessary site preparation is completed before seedlings are planted.

Ratings in the column suitability for mechanical site preparation (surface) are based on slope, depth to a restrictive layer, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsited to this management activity. The part of the soil from the surface to a depth of about 1-foot is considered in the ratings.

Ratings in the column suitability for mechanical site preparation (deep) are based on slope, depth to a restrictive layer, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsited to this management activity. The part of the soil from the surface to a depth of about 3 feet is considered in the ratings.

Ratings in the column potential for seedling mortality are based on flooding, ponding, depth to a water table, content of lime, reaction, salinity, available water capacity, soil moisture regime, soil temperature regime, aspect, and slope. The soils are described as having a low, moderate, or high potential for seedling mortality. See the National Forestry Manual, Subpart B for criteria used in rating management concerns. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

CONSERVATION TREE AND SHRUB MANAGEMENT
Boyd County,
Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
Ab: Albaton-----		Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	High Wetness Soil reaction
AnC: Anselmo-----		Well suited	Well suited	Well suited	Well suited	Low
AnD: Anselmo-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
AnF: Anselmo-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
ArF: Anselmo-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
Rock Outcrop-----		Not rated	Not rated	Not rated	Not rated	Not rated
Ba: Barney-----		Well suited	Well suited	Well suited	Well suited	High Wetness
Bd: Blake-----		Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Moderate Wetness Lime Soil reaction
Be: Blendon-----		Well suited	Well suited	Well suited	Well suited	Low
BeC: Blendon-----		Well suited	Well suited	Well suited	Well suited	Low
BoD: Boyd-----		Poorly suited Stickiness	Poorly suited Stickiness Slope	Poorly suited Stickiness	Well suited	Low
BrG: Bristow-----		Poorly suited Stickiness	Unsuited Slope Stickiness	Poorly suited Slope Stickiness	Poorly suited Slope	Moderate Lime Soil reaction
Bs: Brocksburg-----		Well suited	Well suited	Well suited	Well suited	Low
Bt: Brocksburg-----		Well suited	Well suited	Well suited	Well suited	Low
Cb: Cass-----		Well suited	Well suited	Well suited	Well suited	Low
CrE2: Crofton-----		Well suited	Moderately suited Slope	Well suited	Well suited	Moderate Soil reaction
DuB: Dunday-----		Well suited	Well suited	Well suited	Well suited	Low
DuC: Dunday-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
DuD: Dunday-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
DxB: Dunday-----		Well suited	Well suited	Well suited	Well suited	Low
Et: Eltree-----		Well suited	Well suited	Well suited	Well suited	Low
Go: Grigston-----		Well suited	Well suited	Well suited	Well suited	Low
GrB: Aowa-----		Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
Ha: Hall-----		Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
He: Haynie-----		Well suited	Well suited	Well suited	Well suited	Moderate Lime

CONSERVATION TREE AND SHRUB MANAGEMENT
Boyd County,
Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
IfD: Inavale-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
IgB: Inavale-----		Well suited	Well suited	Well suited	Well suited	Low
IhB: Inavale-----		Well suited	Well suited	Well suited	Well suited	Low
In: Inavale-----		Well suited	Well suited	Well suited	Well suited	Low
Jn: Jansen-----		Well suited	Well suited	Well suited	Well suited	Low
JnC: Jansen-----		Well suited	Well suited	Well suited	Well suited	Low
JnD: Jansen-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
LaD: Labu-----		Poorly suited Stickiness	Poorly suited Stickiness Slope	Poorly suited Stickiness	Well suited	Low
LcF: Labu-----		Poorly suited Stickiness	Poorly suited Slope Stickiness	Poorly suited Slope Stickiness	Poorly suited Slope	Low
Sansarc-----		Poorly suited Stickiness	Poorly suited Slope Stickiness	Poorly suited Slope Stickiness	Poorly suited Slope	Moderate Soil reaction
Le: Leshara-----		Well suited	Well suited	Well suited	Well suited	Low
LsC: Lynch-----		Moderately suited Stickiness	Moderately suited Stickiness	Poorly suited Stickiness	Well suited	Moderate Lime Soil reaction
LsD: Lynch-----		Moderately suited Stickiness	Moderately suited Stickiness Slope	Poorly suited Stickiness	Well suited	Moderate Lime Soil reaction
LyD: Lynch-----		Moderately suited Stickiness	Moderately suited Stickiness Slope	Poorly suited Stickiness	Well suited	Moderate Lime Soil reaction
Bristow-----		Poorly suited Stickiness	Poorly suited Stickiness Slope	Poorly suited Stickiness	Well suited	Moderate Lime Soil reaction
LyF: Lynch-----		Moderately suited Stickiness	Poorly suited Slope Stickiness	Poorly suited Slope Stickiness	Poorly suited Slope	Moderate Lime Soil reaction
Bristow-----		Poorly suited Stickiness	Poorly suited Slope Stickiness	Poorly suited Slope Stickiness	Poorly suited Slope	Moderate Lime Soil reaction
MaG: Mariaville-----		Well suited	Unsuited Slope	Poorly suited Slope	Poorly suited Slope	Moderate Soil reaction
Paka-----		Well suited	Unsuited Slope	Poorly suited Slope	Poorly suited Slope	Low
MeE: Meadin-----		Well suited	Moderately suited Slope Rock fragments	Well suited	Well suited	Low
NoC: Nora-----		Well suited	Well suited	Well suited	Well suited	Low
NoD: Nora-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
Oa: Onawa-----		Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Moderate Lime Soil reaction
Oe: O'Neill-----		Well suited	Well suited	Well suited	Well suited	Low

CONSERVATION TREE AND SHRUB MANAGEMENT
Boyd County,
Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
OeC: O'Neill-----		Well suited	Well suited	Well suited	Well suited	Low
OFD: O'Neill-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
Meadin-----		Well suited	Moderately suited Slope Rock fragments	Well suited	Well suited	Low
On: Onita----- Poorly Drained-----		Well suited Unsuited Horizon table contains no data	Well suited Unsuited Horizon table contains no data	Well suited Unsuited Horizon table contains no data	Well suited Unsuited Horizon table contains no data	Low High Horizon table contains no data
Or: Ord-----		Well suited	Well suited	Well suited	Well suited	Moderate Lime
PaC: Paka-----		Well suited	Well suited	Well suited	Well suited	Low
Ph: Paka-----		Well suited	Well suited	Well suited	Well suited	Low
PhC: Paka-----		Well suited	Well suited	Well suited	Well suited	Low
PhD: Paka-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
PoC: Promise-----		Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
RaC: Ree-----		Well suited	Well suited	Well suited	Well suited	Low
RaD: Ree-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
RaE: Ree-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
ReC: Reliance-----		Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
ReD: Reliance-----		Moderately suited Stickiness	Moderately suited Slope Stickiness	Well suited	Well suited	Low
RfC: Reliance-----		Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Rw: Riverwash-----		Moderately suited Sandiness	Moderately suited Rock fragments Sandiness	Well suited	Well suited	High Wetness Soil reaction
SaG: Sansarc-----		Poorly suited Stickiness Slope	Unsuited Slope Stickiness	Poorly suited Slope Stickiness	Poorly suited Slope	Moderate Soil reaction
Sc: Scott-----		Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	High Wetness
Sm: Simeon-----		Moderately suited Sandiness	Moderately suited Sandiness	Well suited	Well suited	Low
SuC: Simeon----- Valentine-----		Well suited Well suited	Well suited Well suited	Well suited Well suited	Well suited Well suited	Low Low
SvF2: Simeon-----		Well suited	Poorly suited Slope	Poorly suited Slope	Poorly suited Slope	Low

CONSERVATION TREE AND SHRUB MANAGEMENT
Boyd County,
Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
Valentine-----		Moderately suited Sandiness	Poorly suited Slope Sandiness	Poorly suited Slope	Poorly suited Slope	Low
VaE: Valentine-----		Moderately suited Sandiness	Moderately suited Slope Sandiness	Well suited	Well suited	Low
VbB: Valentine-----		Well suited	Well suited	Well suited	Well suited	Low
Ve: Verdel-----		Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
WeC: Wewela-----		Well suited	Well suited	Well suited	Well suited	Low
zwa: Water-----		Not rated	Not rated	Not rated	Not rated	Not rated

ENGINEERING INDEX PROPERTIES
Boyd County, Nebraska

Engineering Index Properties table gives the engineering classifications and the range of index properties for the layers of each soil in the survey area. Depth to the upper and lower boundaries of each layer is indicated. Texture is given in the standard terms used by the U.S. Department of Agriculture. These terms are defined according to percentages of sand, silt, and clay in the fraction of the soil that is less than 2 millimeters in diameter. Loam, for example, is soil that is 7 to 27 percent clay, 28 to 50 percent silt, and less than 52 percent sand. If the content of particles coarser than sand is 15 percent or more, an appropriate modifier is added, for example, gravelly. Textural terms are defined in the Glossary.

Classification of the soils is determined according to the Unified soil classification system (ASTM, 1998) and the system adopted by the American Association of State Highway and Transportation Officials (AASHTO, 1998). The Unified system classifies soils according to properties that affect their use as construction material. Soils are classified according to particle-size distribution of the fraction less than 3 inches in diameter and according to plasticity index, liquid limit, and organic matter content. Sandy and gravelly soils are identified as GW, GP, GM, GC, SW, SP, SM, and SC; silty and clayey soils as ML, CL, OL, MH, CH, and OH; and highly organic soils as PT. Soils exhibiting engineering properties of two groups can have a dual classification, for example, CL-ML.

The AASHTO system classifies soils according to those properties that affect roadway construction and maintenance. In this system, the fraction of a mineral soil that is less than 3 inches in diameter is classified in one of seven groups from A-1 through A-7 on the basis of particle-size distribution, liquid limit, and plasticity index. Soils in group A-1 are coarse grained and low in content of fines (silt and clay). At the other extreme, soils in group A-7 are fine grained. Highly organic soils are classified in group A-8 on the basis of visual inspection. If laboratory data are available, the A-1, A-2, and A-7 groups are further classified as A-1-a, A-1-b, A-2-4, A-2-5, A-2-6, A-2-7, A-7-5, or A-7-6. As an additional refinement, the suitability of a soil as subgrade material can be indicated by a group index number. Group index numbers range from 0 for the best subgrade material to 20 or higher for the poorest. The AASHTO classification for soils tested, with group index numbers in parentheses, is given in Engineering Index Properties table.

Rock fragments larger than 10 inches in diameter and 3 to 10 inches in diameter are indicated as a percentage of the total soil on a dry-weight basis. The percentages are estimates determined mainly by converting volume percentage in the field to weight percentage. Percentage (of soil particles) passing designated sieves is the percentage of the soil fraction less than 3 inches in diameter based on an oven-dry weight. The sieves, numbers 4, 10, 40, and 200 (USA Standard Series), have openings of 4.76, 2.00, 0.420, and 0.074 millimeters, respectively. Estimates are based on laboratory tests of soils sampled in the survey area and in nearby areas and on estimates made in the field.

Liquid limit and plasticity index (Atterberg limits) indicate the plasticity characteristics of a soil. The estimates are based on test data from the survey area or from nearby areas and on field examination. The estimates of particle-size distribution, liquid limit, and plasticity index are generally rounded to the nearest 5 percent. Thus, if the ranges of gradation and Atterberg limits extend a marginal amount (1 or 2 percentage points) across classification boundaries, the classification in the marginal zone is generally omitted in the table.

ENGINEERING INDEX PROPERTIES--Continued
Boyd County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
Ab: Albaton-----	0-7	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	40-60
	7-54	Stratified silty clay to silt loam	CH	A-7	0	0	100	100	95-100	95-100	60-85	40-60
	54-60	Silty clay			---	---	---	---	---	---	60-85	40-60
AnC: Anselmo-----	0-16	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	100	60-100	30-65	15-25	NP-7
	16-21	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	100	70-95	35-65	15-25	NP-7
	21-34	Fine sandy loam	SC-SM, SM	A-4	0	0	100	100	70-85	40-50	15-25	NP-7
	34-60	Loamy fine sand	SC-SM, SM	A-2	0	0	100	100	50-80	15-35	15-20	NP-5
AnD: Anselmo-----	0-16	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	100	60-100	30-65	15-25	NP-7
	16-21	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	100	70-95	35-65	15-25	NP-7
	21-34	Fine sandy loam	SC-SM, SM	A-4	0	0	100	100	70-85	40-50	15-25	NP-7
	34-60	Loamy fine sand	SC-SM, SM	A-2	0	0	100	100	50-80	15-35	15-20	NP-5
AnF: Anselmo-----	0-16	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	100	60-100	30-65	15-25	NP-7
	16-21	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	100	70-95	35-65	15-25	NP-7
	21-34	Fine sandy loam	SC-SM, SM	A-4	0	0	100	100	70-85	40-50	15-25	NP-7
	34-60	Loamy fine sand	SC-SM, SM	A-2	0	0	100	100	50-80	15-35	15-20	NP-5
ArF: Anselmo-----	0-16	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	100	60-100	30-65	15-25	NP-7
	16-21	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	100	70-95	35-65	15-25	NP-7
	21-34	Fine sandy loam	SC-SM, SM	A-4	0	0	100	100	70-85	40-50	15-25	NP-7
	34-60	Loamy fine sand	SC-SM, SM	A-2	0	0	100	100	50-80	15-35	15-20	NP-5
Rock Outcrop---	0-16	Fine sandy loam	SC-SM, SM, CL-ML, ML	A-2, A-4	0	0	100	100	60-100	30-65	15-25	NP-7
	16-21	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	100	70-95	35-65	15-25	NP-7
	21-34	Fine sandy loam	SC-SM, SM	A-4	0	0	100	100	70-85	40-50	15-25	NP-7
	34-60	Loamy fine sand	SC-SM, SM	A-2	0	0	100	100	50-80	15-35	15-20	NP-5
	0-60	Unweathered bedrock			---	---	---	---	---	---	0-14	---
Ba: Barney-----	0-5	Silt loam	CL	A-6	0	0	100	95-100	90-100	70-95	25-40	12-20
	5-10	Stratified loam to sand	ML, SM	A-2, A-4	0	0	95-100	90-100	55-80	20-60	10-20	NP-5
	10-60	Fine sand	SC-SM, SM, SP, SP-SM	A-1, A-2, A-3	0	0	95-100	75-100	30-80	3-15	10-15	NP-5
Bd: Blake-----	0-8	Silty clay loam	CL	A-6, A-7	0	0	100	100	90-100	85-95	35-50	15-30
	8-25	Stratified silty clay loam to silty clay	CL	A-6, A-7	0	0	100	100	90-100	85-95	30-50	10-30
	25-60	Stratified very fine sandy loam to silty clay loam	CL, ML	A-4, A-6	0	0	100	100	80-90	75-90	30-40	5-15
Be: Blendon-----	0-16	Fine sandy loam	SM	A-4	0	0	100	90-100	60-100	35-50	20-30	NP-5
	16-42	Sandy loam	SC, SC-SM, SM	A-2, A-4	0	0	100	85-100	60-100	20-45	15-30	NP-10
	42-60	Loamy sand	SC-SM, SM, SP-SM	A-2, A-4	0	0	85-100	65-100	50-100	10-45	15-25	NP-5
BeC: Blendon-----	0-16	Fine sandy loam	SM	A-4	0	0	100	90-100	60-100	35-50	20-30	NP-5
	16-42	Sandy loam	SC, SC-SM, SM	A-2, A-4	0	0	100	85-100	60-100	20-45	15-30	NP-10
	42-60	Loamy sand	SC-SM, SM, SP-SM	A-2, A-4	0	0	85-100	65-100	50-100	10-45	15-25	NP-5
BoD: Boyd-----	0-6	Silty clay	CH, MH	A-7	0	0	100	95-100	95-100	90-100	65-100	30-60
	6-22	Clay	CH, MH	A-7	0	0	100	95-100	95-100	80-100	65-100	30-60
	22-32	Clay	CH, MH	A-7	0	0	95-100	85-100	75-100	60-100	65-100	30-60
	32-60	Weathered bedrock			---	---	---	---	---	---	---	---
BrG: Bristow-----	0-7	Silty clay	CH, CL	A-7	0	0	95-100	95-100	90-100	90-100	45-65	25-40
	7-17	Clay	CH, CL	A-7	0	0	100	100	100	85-100	45-65	25-45
	17-80	Weathered bedrock			---	---	---	---	---	---	---	---
Bs: Brocksburg----	0-18	Fine sandy loam	SC-SM, SM	A-4	0	0	100	100	70-85	35-50	15-20	NP-5
	18-32	Clay loam	CL	A-6, A-7	0	0	100	100	90-100	70-80	35-45	11-20
	32-60	Gravelly sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	85-95	50-90	20-60	3-15	---	NP
Bt: Brocksburg----	0-14	Loam	CL, ML	A-4, A-6	0	0	100	100	90-100	70-90	25-40	3-15
	14-34	Clay loam	CL	A-6, A-7	0	0	100	100	90-100	70-80	35-45	11-20
	34-60	Gravelly coarse sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	85-95	50-90	20-60	3-15	---	NP

ENGINEERING INDEX PROPERTIES--Continued
Boyd County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct					Pct	
Cb: Cass-----	In											
	0-12	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	85-95	20-40	15-20	NP-5
	12-26	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	85-95	20-50	15-20	NP-5
	26-60	Stratified coarse sand to fine sandy loam	SM, SP-SM	A-2, A-3	0	0	95-100	95-100	50-75	5-30	10-15	NP-4
CrE2: Crofton-----	0-6	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	6-60	Silt loam	CL	A-6, A-7	0	0	100	95-100	95-100	95-100	30-50	5-25
DuB: Dunday-----	0-15	Loamy fine sand	SC-SM, SM	A-2	0	0	100	100	90-100	13-35	15-25	NP-5
	15-60	Fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	50-95	5-35	15-25	NP-5
DuC: Dunday-----	0-15	Loamy fine sand	SC-SM, SM	A-2	0	0	100	100	90-100	13-35	15-25	NP-5
	15-60	Fine sand	SC-SM, SM, SP-SM	A-3, A-2	0	0	100	100	50-95	5-35	15-25	NP-5
DuD: Dunday-----	0-15	Loamy fine sand	SC-SM, SM	A-2	0	0	100	100	90-100	13-35	15-25	NP-5
	15-60	Fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	50-95	5-35	15-25	NP-5
DxB: Dunday-----	0-14	Loamy fine sand	SC-SM, SM	A-2, A-4	0	0	100	100	75-85	30-50	15-25	NP-5
	14-34	Loamy sand	SC-SM, SM	A-2, A-4	0	0	100	100	50-85	15-40	15-25	NP-5
	34-48	Sandy loam	SC, SC-SM, SM	A-2, A-4	0	0	100	100	60-70	30-40	20-30	3-10
	48-60	Coarse sand	SP-SM, SC-SM, SM, SP	A-1	0	0	85-90	50-90	30-50	3-20	5-10	NP-5
Et: Eltree-----	0-25	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	85-100	65-100	20-40	3-15
	25-38	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	85-100	85-100	25-40	5-20
	38-60	Silt loam	CL	A-4, A-6, A- 7-6	0	0	100	95-100	90-100	65-100	25-45	7-22
Go: Grigston-----	0-15	Silt loam	CL	A-6	0	0	100	100	95-100	80-100	30-35	10-15
	15-31	Silt loam	CL	A-6	0	0	100	100	95-100	85-100	30-40	10-20
	31-60	Stratified loam to fine sandy loam	CL	A-6	0	0	100	100	85-100	70-100	30-40	10-20
GrB: Aowa-----	0-15	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	90-100	80-100	20-40	5-20
	15-60	Stratified silt loam to fine sand	CL, CL-ML, ML	A-4, A-6, A-7	0	0	100	100	85-100	80-100	22-46	3-20
Ha: Hall-----	0-5	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	95-100	95-100	25-40	3-18
	5-36	Silty clay loam	CL	A-6, A-7	0	0	100	100	95-100	95-100	35-50	15-30
	36-42	Silty clay loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	90-100	25-40	5-20
	42-60	Stratified silty clay loam to fine sandy loam	CL, CL-ML, ML	A-4	0	0	100	100	85-100	75-100	20-40	5-20
He: Haynie-----	0-7	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	85-100	70-100	25-40	5-15
	7-60	Stratified silt loam to fine sand	CL, CL-ML	A-4, A-6	0	0	100	100	85-100	85-100	25-35	5-15
IfD: Inavale-----	0-6	Fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	90-100	65-85	5-30	15-25	NP-5
	6-12	Fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	90-100	65-85	5-30	15-25	NP-5
	12-60	Fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	70-90	5-30	15-25	NP-5
IgB: Inavale-----	0-6	Fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	90-100	65-85	5-30	15-25	NP-5
	6-12	Fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	90-100	65-85	5-30	15-25	NP-5
	12-60	Fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	70-90	5-30	15-25	NP-5
IhB: Inavale-----	0-9	Loamy fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	85-95	5-35	15-25	NP-5
	9-19	Fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	90-100	65-85	5-30	15-25	NP-5
	19-60	Loamy fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	70-90	5-30	15-25	NP-5
In: Inavale-----	0-16	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	95-100	65-85	35-55	15-20	NP-5
	16-26	Fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	90-100	65-85	5-30	15-20	NP-5
	26-60	Fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	70-90	5-30	15-20	NP-5

ENGINEERING INDEX PROPERTIES--Continued
Boyd County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct						
Jn: Jansen-----	In 0-13 13-32 32-60	Loam Clay loam Gravelly coarse sand	CL, ML CL SP, SP-SM, SW, SW-SM	A-4, A-6 A-6, A-7 A-1, A-2, A-3	0 0 0	0 0 0	100 95-100 85-100	100 90-100 45-100	90-100 80-100 35-65	50-95 50-80 3-10	25-40 30-45 0-10	3-15 10-25 NP-5
JnC: Jansen-----	0-13 13-32 32-60	Loam Clay loam Gravelly coarse sand	CL, ML CL SP, SP-SM, SW, SW-SM	A-4, A-6 A-6, A-7 A-1, A-2, A-3	0 0 0	0 0 0	100 95-100 85-100	100 90-100 45-100	90-100 80-100 35-65	50-95 50-80 3-10	25-40 30-45 0-10	3-15 10-25 NP-5
JnD: Jansen-----	0-13 13-32 32-60	Loam Clay loam Gravelly coarse sand	CL, ML CL SP, SP-SM, SW, SW-SM	A-4, A-6 A-6, A-7 A-1, A-2, A-3	0 0 0	0 0 0	100 95-100 85-100	100 90-100 45-100	90-100 80-100 35-65	50-95 50-80 3-10	25-40 30-45 0-10	3-15 10-25 NP-5
LaD: Labu-----	0-6 6-34 34-60	Silty clay Clay Weathered bedrock	CH, MH CH, MH	A-7 A-7	0 0	0 0	100 95-100	95-100 95-100	90-100 90-100	85-100 85-100	50-90 50-90	20-55 20-55 ---
LcF: Labu-----	0-6 6-34 34-60	Silty clay Clay Weathered bedrock	MH, CH CH, MH	A-7 A-7	0 0	0 0	100 95-100	95-100 95-100	90-100 90-100	85-100 85-100	50-90 50-90	20-55 20-55 ---
Sansarc-----	0-14 14-60	Silty clay Weathered bedrock	CH, MH	A-7	0 ---	0 ---	95-100 ---	95-100 ---	95-100 ---	85-100 ---	60-90 ---	25-55 ---
Le: Leshara-----	0-21 21-32 32-60	Silt loam Silt loam Stratified silt loam to fine sand	CL, CL-ML, ML CL, CL-ML, ML CL, CL-ML, ML	A-4, A-6 A-4, A-6 A-4, A-6	0 0 0	0 0 0	100 100 100	100 100 100	90-100 90-100 90-100	60-90 60-90 60-90	20-35 20-35 20-35	3-15 3-15 3-15
LsC: Lynch-----	0-8 8-28 28-36 36-60	Silty clay Silty clay Silty clay Weathered bedrock	CH, MH CH, MH CH, CL	A-7 A-7 A-7	0 0 0	0 0 0	100 100 100	100 100 100	95-100 95-100 95-100	90-100 75-100 85-100	50-70 50-70 45-70	20-40 25-35 25-40 ---
LsD: Lynch-----	0-8 8-28 28-36 36-60	Silty clay Silty clay Silty clay Weathered bedrock	CH, MH CH, MH CH, CL	A-7 A-7 A-7	0 0 0	0 0 0	100 100 100	100 100 100	95-100 95-100 95-100	90-100 75-100 85-100	50-70 50-70 45-70	20-40 25-35 25-40 ---
LyD: Lynch-----	0-8 8-28 28-36 36-60	Silty clay Silty clay Silty clay Weathered bedrock	CH, MH CH, MH CH, CL	A-7 A-7 A-7	0 0 0	0 0 0	100 100 100	100 100 100	95-100 95-100 95-100	90-100 75-100 85-100	50-70 50-70 45-70	20-40 25-35 25-40 ---
Bristow-----	0-7 7-17 17-60	Silty clay Clay Weathered bedrock	CH, CL CH, CL	A-7 A-7	0 0	0 0	95-100 100	95-100 100	90-100 100	90-100 85-100	45-65 45-65	25-40 25-45 ---
LyF: Lynch-----	0-8 8-28 28-36 36-60	Silty clay Silty clay Silty clay Weathered bedrock	CH, MH CH, MH CH, CL	A-7 A-7 A-7	0 0 0	0 0 0	100 100 100	100 100 100	95-100 95-100 95-100	90-100 75-100 85-100	50-70 50-70 45-70	20-40 25-35 25-40 ---
Bristow-----	0-7 7-17 17-60	Silty clay Clay Weathered bedrock	CH, CL CH, CL	A-7 A-7	0 0	0 0	95-100 100	95-100 100	90-100 100	90-100 85-100	45-65 45-65	25-40 25-45 ---
MaG: Mariaville----	0-5 5-17 17-60	Loam Silty clay loam Weathered bedrock	CL, CL-ML CL, CL-ML	A-4, A-6 A-4, A-6	0 0	0 0	95-100 95-100	95-100 95-100	85-100 85-100	50-95 50-95	20-35 20-45	5-15 5-25 ---
Paka-----	0-8 8-26 26-48 48-60	Loam Clay loam Silt loam Weathered bedrock	CL CL CL	A-4, A-6 A-6, A-7 A-4, A-6, A-7	0 0 0	0 0 0	100 100 100	100 100 100	95-100 95-100 95-100	60-90 80-95 75-95	20-40 30-50 15-50	5-20 10-25 5-25 ---
MeE: Meadin-----	0-8 8-13 13-60	Sandy loam Gravelly sandy loam Very gravelly coarse sand	CL-ML, ML, SC-SM, SM GM, GP-GM, SM, SP-SM GP, GP-GM, SP, SP-SM	A-2, A-4 A-2, A-3 A-1	0 0 0	0 0 0	85-100 50-90 40-80	75-95 50-90 30-70	45-80 50-65 15-50	25-55 5-35 1-10	15-20 --- ---	NP-5 NP NP
NoC: Nora-----	0-7 7-34 34-60	Silt loam Silt loam Silt loam	CL, ML CL, ML CL, CL-ML, ML	A-4, A-6, A-7 A-6, A-7 A-4, A-6, A-7	0 0 0	0 0 0	100 95-100 95-100	100 95-100 95-100	95-100 95-100 95-100	85-100 85-100 85-100	30-46 35-50 27-50	7-23 11-20 6-20

ENGINEERING INDEX PROPERTIES--Continued
Boyd County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct						
NoD: Nora-----	0-7 7-34 34-60	Silt loam Silt loam Silt loam	CL, ML CL, ML CL, CL-ML, ML	A-4, A-6, A-7 A-6, A-7 A-4, A-6, A-7	0 0 0	0 0 0	100 95-100 95-100	100 95-100 95-100	95-100 95-100 95-100	85-100 85-100 85-100	30-46 35-50 27-50	7-23 11-20 6-20
Oa: Onawa-----	0-7 7-28 28-60	Silty clay Silty clay Stratified silt loam to clay	CH CH CL, CL-ML	A-7 A-7 A-4, A-6	0 0 0	0 0 0	100 100 100	100 100 100	95-100 95-100 95-100	95-100 95-100 85-100	60-85 60-85 25-40	40-60 40-60 5-20
Oe: O'neill-----	0-9 9-34 34-60	Fine sandy loam Sandy loam Coarse sand	CL, ML, SC- SM, SM SC, SC-SM SP, SP-SM	A-4 A-2, A-4 A-1, A-2, A-3	0 0 0	0 0 0	95-100 95-100 70-100	95-100 95-100 50-90	70-85 60-75 25-60	35-55 30-50 0-5	15-25 15-30 10-20	NP-10 NP-10 NP-5
OeC: O'neill-----	0-9 9-34 34-60	Fine sandy loam Fine sandy loam Gravelly sand	CL, ML, SC- SM, SM SC, SC-SM SP, SP-SM	A-4 A-2, A-4 A-1, A-2, A-3	0 0 0	0 0 0	95-100 95-100 70-100	95-100 95-100 50-90	70-85 60-75 25-60	35-55 30-50 0-5	15-25 15-30 10-20	NP-10 NP-10 NP-5
OfD: O'neill-----	0-9 9-34 34-60	Fine sandy loam Fine sandy loam Gravelly sand	CL, ML, SC- SM, SM SC, SC-SM SP, SP-SM	A-4 A-2, A-4 A-1, A-2, A-3	0 0 0	0 0 0	95-100 95-100 70-100	95-100 95-100 50-90	70-85 60-75 25-60	35-55 30-50 0-5	15-25 15-30 10-20	NP-10 NP-10 NP-5
Meadin-----	0-8 8-13 13-60	Fine sandy loam Gravelly sandy loam Very gravelly coarse sand	CL-ML, ML, SC-SM, SM GM, GP-GM, SM, SP-SM GP, GP-GM, SP, SP-SM	A-2, A-4 A-2, A-4 A-2, A-3 A-1	0 0 0 0	0 0 0 0	85-100 50-90 50-90 40-80	75-95 50-90 50-90 30-70	45-80 50-65 50-65 15-50	25-55 5-35 5-35 1-10	15-20 --- --- ---	NP-5 NP NP NP
On: Onita-----	0-18 18-32 32-60	Silt loam Silty clay loam Silt loam	CL, ML CH, CL, MH, ML CH, CL	A-4, A-6, A- 7-6 A-7 A-6, A-7	0 0 0	0 0 0-5	100 100 95-100	95-100 95-100 95-100	90-100 90-100 85-100	70-100 75-100 65-100	30-45 40-60 30-55	5-20 10-30 10-30
Poorly Drained- Or-----	0-11 11-30 30-60	Fine sandy loam Fine sandy loam Stratified sand to loamy fine sand	ML, SM ML, SM SC-SM, SM, SP-SM	A-2, A-4 A-2, A-4 A-2, A-3	0 0 0	0 0 0	95-100 95-100 95-100	95-100 95-100 95-100	70-98 70-100 50-100	30-90 30-85 5-30	20-35 20-35 15-20	NP-10 NP-10 NP-5
PaC: Paka-----	0-9 9-27 27-46 46-60	Fine sandy loam Clay loam Silt loam Weathered bedrock	SC-SM, SM CL CL CL	A-4 A-7, A-6 A-4, A-6, A-7	0 0 0	0 0 0	100 100 100	100 100 100	70-90 95-100 95-100	40-70 80-95 75-95	15-35 30-50 15-50	NP-10 10-25 5-25
Ph: Paka-----	0-8 8-26 26-48 48-60	Loam Silty clay loam Silt loam Weathered bedrock	CL CL CL CL	A-4, A-6 A-6, A-7 A-4, A-6, A-7	0 0 0	0 0 0	100 100 100	100 100 100	95-100 95-100 95-100	60-90 80-95 75-95	20-40 30-50 15-50	5-20 10-25 5-25
PhC: Paka-----	0-8 8-26 26-48 48-60	Loam Silty clay loam Silt loam Weathered bedrock	CL CL CL CL	A-4, A-6 A-6, A-7 A-4, A-6, A-7	0 0 0	0 0 0	100 100 100	100 100 100	95-100 95-100 95-100	60-90 80-95 75-95	20-40 30-50 15-50	5-20 10-25 5-25
PhD: Paka-----	0-8 8-26 26-48 48-60	Loam Clay loam Silt loam Weathered bedrock	CL CL CL CL	A-4, A-6 A-6, A-7 A-4, A-6, A-7	0 0 0	0 0 0	100 100 100	100 100 100	95-100 95-100 95-100	60-90 80-95 75-95	20-40 30-50 15-50	5-20 10-25 5-25
PoC: Promise-----	0-18 18-33 33-60	Silty clay Clay Clay	CH, MH CH, MH CH, MH	A-7 A-7 A-7	0 0 0	0 0 0	100 100 100	100 100 100	90-100 90-100 90-100	80-100 85-100 85-100	55-90 60-85 60-90	25-55 25-50 25-55
RaC: Ree-----	0-7 7-22 22-60	Silt loam Clay loam Sandy loam	CL, CL-ML, ML CL CL, CL-ML, SC, SC-SM	A-4, A-6, A-7 A-6, A-7 A-4, A-6, A-7	0 0 0	0 0 0	95-100 95-100 95-100	90-100 90-100 85-100	80-100 70-100 70-100	70-95 65-85 35-85	24-45 30-45 25-45	3-20 10-20 5-22
RaD: Ree-----	0-7 7-22 22-60	Silt loam Silty clay loam Sandy loam	CL, CL-ML, ML CL CL, CL-ML, SC, SC-SM	A-4, A-6, A-7 A-6, A-7 A-4, A-6, A-7	0 0 0	0 0 0	95-100 95-100 95-100	90-100 90-100 85-100	80-100 70-100 70-100	70-95 65-85 35-85	24-45 30-45 25-45	3-20 10-20 5-22
RaE: Ree-----	0-7 7-22 22-60	Silt loam Sandy clay loam Fine sandy loam	CL, CL-ML, ML CL CL, CL-ML, SC, SC-SM	A-4, A-6, A-7 A-6, A-7 A-4, A-6, A-7	0 0 0	0 0 0	95-100 95-100 95-100	90-100 90-100 85-100	80-100 70-100 70-100	70-95 65-85 35-85	24-45 30-45 25-45	3-20 10-20 5-22

ENGINEERING INDEX PROPERTIES--Continued
Boyd County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct						
	In											
ReC: Reliance-----	0-10	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	95-100	70-100	20-40	3-15
	10-28	Silty clay loam	CH, CL	A-6, A-7	0	0	100	100	95-100	85-100	35-60	15-30
	28-60	Silt loam	CH, CL	A-6, A-7	0	0	100	100	90-100	70-100	30-55	10-30
ReD: Reliance-----	0-10	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	95-100	70-100	20-40	3-15
	10-28	Silty clay loam	CH, CL	A-6, A-7	0	0	100	100	95-100	85-100	35-60	15-30
	28-60	Silt loam	CH, CL	A-6, A-7	0	0	100	100	90-100	70-100	30-55	10-30
RfC: Reliance-----	0-10	Silty clay loam	CH, CL, MH, ML	A-6, A-7	0	0	100	100	95-100	80-100	35-55	15-25
	10-28	Silty clay loam	CH, CL	A-6, A-7	0	0	100	100	95-100	85-100	35-60	15-30
	28-60	Silty clay loam	CH, CL	A-6, A-7	0	0	100	100	90-100	70-100	30-55	10-30
Rw: Riverwash-----	0-6	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	90-100	65-80	5-35	15-20	NP
	6-60	Stratified sand to very gravelly coarse sand	GP, GW, SP, SW	A-1	0	0-10	50-100	30-90	20-80	3-15	---	NP
SaG: Sansarc-----	0-16	Silty clay	CH, MH	A-7	0	0	95-100	95-100	95-100	85-100	60-90	25-55
	16-60	Weathered bedrock			---	---	---	---	---	---	---	---
Sc: Scott-----	0-8	Silt loam	CL, CL-ML, ML	A-4, A-6, A-7	0	0	100	100	100	95-100	20-45	2-20
	8-48	Silty clay	CH, CL	A-7	0	0	100	100	100	95-100	41-75	20-45
	48-56	Silty clay loam	CH, CL	A-6, A-7	0	0	100	100	100	95-100	35-60	20-40
	56-64	Clay loam	CL	A-4, A-6, A-7	0	0	100	100	90-100	90-100	25-50	8-24
Sm: Simeon-----	0-5	Loamy sand	SM, SP-SM	A-2, A-3	0	0	95-100	90-100	51-95	5-35	10-20	NP-5
	5-60	Sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	90-100	80-100	35-95	0-30	10-20	NP-5
SuC: Simeon-----	0-15	Loamy sand	SM, SP-SM	A-2, A-3	0	0	95-100	90-100	51-95	5-35	10-20	NP-5
	15-60	Coarse sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	90-100	80-100	35-95	0-30	10-20	NP-5
Valentine-----	0-5	Loamy sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	95-100	2-35	15-20	NP-5
	5-11	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	90-100	2-35	15-20	NP-5
	11-60	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
SvF2: Simeon-----	0-15	Loamy sand	SM, SP-SM	A-2, A-3	0	0	95-100	90-100	51-95	5-35	10-20	NP-5
	15-60	Coarse sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	90-100	80-100	35-95	0-30	10-20	NP-5
Valentine-----	0-5	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
	5-11	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	90-100	2-35	15-20	NP-5
	11-60	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
VaE: Valentine-----	0-5	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
	5-11	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	90-100	2-35	15-20	NP-5
	11-60	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
VbB: Valentine-----	0-5	Loamy sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	95-100	2-35	15-20	NP-5
	5-11	Loamy sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	90-100	2-35	15-20	NP-5
	11-60	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
Ve: Verdel-----	0-14	Silty clay	CH	A-7	0	0	100	95-100	95-100	85-100	50-70	27-45
	14-60	Silty clay	CH	A-7	0	0	100	95-100	95-100	85-100	50-70	27-45
WeC: Wewela-----	0-8	Fine sandy loam	ML, SC-SM, SM, CL-ML	A-4	0	0	100	100	70-85	40-55	15-30	NP-7
	8-15	Sandy clay loam	CL, SC	A-6, A-7	0	0	100	100	60-100	35-55	30-45	10-20
	15-22	Clay	CH, MH	A-7	0	0	100	95-100	90-100	85-100	55-85	20-50
	22-60	Weathered bedrock			---	---	---	---	---	---	---	---
zwa: Water-----	---	---	---	---	---	---	---	---	---	---	---	---

Physical Properties

This table shows estimates of some physical characteristics and features that affect soil behavior. These estimates are given for the layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils.

Depth to the upper and lower boundaries of each layer is indicated.

Particle size is the effective diameter of a soil particle as measured by sedimentation, sieving, or micrometric methods. Particle sizes are expressed as classes with specific effective diameter class limits. The broad classes are sand, silt, and clay, ranging from the larger to the smaller.

Sand as a soil separate consists of mineral soil particles that are 0.05 millimeter to 2 millimeters in diameter. In this table, the estimated sand content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Silt as a soil separate consists of mineral soil particles that are 0.002 to 0.05 millimeter in diameter. In this table, the estimated silt content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Clay as a soil separate consists of mineral soil particles that are less than 0.002 millimeter in diameter. In this table, the estimated clay content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of sand, silt, and clay affects the physical behavior of a soil. Particle size is important for engineering and agronomic interpretations, for determination of soil hydrologic qualities, and for soil classification.

The amount and kind of clay affect the fertility and physical condition of the soil and the ability of the soil to adsorb cations and to retain moisture. They influence shrink-swell potential, permeability, plasticity, the ease of soil dispersion, and other soil properties. The amount and kind of clay in a soil also affect tillage and earthmoving operations.

Moist bulk density is the weight of soil (ovendry) per unit volume. Volume is measured when the soil is at field moisture capacity, that is, the moisture content at 1/3- or 1/10-bar (33kPa or 10kPa) moisture tension. Weight is determined after the soil is dried at 105 degrees C. In the table, the estimated moist bulk density of each soil horizon is expressed in grams per cubic centimeter of soil material that is less than 2 millimeters in diameter. Bulk density data are used to compute shrink-swell potential, available water capacity, total pore space, and other soil properties. The moist bulk density of a soil indicates the pore space available for water and roots. Depending on soil texture, a bulk density of more than 1.4 can restrict water storage and root penetration. Moist bulk density is influenced by texture, kind of clay, content of organic matter, and soil structure.

Saturated hydraulic conductivity refers to the ability of a soil to transmit water or air. The term "permeability," as used in soil surveys, indicates saturated hydraulic conductivity (K-sat). The estimates in the table indicate the rate of water movement, in micrometers per second (um/sec), when the soil is saturated. They are based on soil characteristics observed in the field, particularly structure, porosity, and texture. Permeability is considered in the design of soil drainage systems and septic tank absorption fields.

Available water capacity refers to the quantity of water that the soil is capable of storing for use by plants. The capacity for water storage is given in inches of water per inch of soil for each soil layer. The capacity varies, depending on soil properties that affect retention of water. The most important properties are the content of organic matter, soil texture, bulk density, and soil structure. Available water capacity is an important factor in the choice of plants or crops to be grown and in the design and management of irrigation systems. Available water capacity is not an estimate of the quantity of water actually available to plants at any given time.

Linear extensibility refers to the change in length of an unconfined clod as moisture content is decreased from a moist to a dry state. It is an expression of the volume change between the water content of the clod at 1/3- or 1/10-bar tension (33kPa or 10kPa tension) and oven dryness. The volume change is reported in the table as percent change for the whole soil. Volume change is influenced by the amount and type of clay minerals in the soil.

Linear extensibility is used to determine the shrink-swell potential of soils. The shrink-swell potential is low if the soil has a linear extensibility of less than 3 percent; moderate if 3 to 6 percent; high if 6 to 9 percent; and very high if more than 9 percent. If the linear extensibility is more than 3, shrinking and swelling can cause damage to buildings, roads, and other structures and to plant roots. Special design commonly is needed.

Organic matter is the plant and animal residue in the soil at various stages of decomposition. In this table, the estimated content of organic matter is expressed as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of organic matter in a soil can be maintained by returning crop residue to the soil. Organic matter has a positive effect on available water capacity, water infiltration, soil organism activity, and tilth. It is a source of nitrogen and other nutrients for crops and soil organisms.

Erosion factors are shown in this table as the K factor (Kw and Kf) and the T factor. Erosion factor K indicates the susceptibility of a soil to sheet and rill erosion by water. Factor K is one of six factors used in the Universal Soil Loss Equation (USLE) and the Revised Universal Soil Loss Equation (RUSLE) to predict the average annual rate of soil loss by sheet and rill erosion in tons per acre per year. The estimates are based primarily on percentage of silt, sand, and organic matter and on soil structure and permeability. Values of K range from 0.02 to 0.69. Other factors being equal, the higher the value, the more susceptible the soil is to sheet and rill erosion by water.

Erosion factor Kw indicates the erodibility of the whole soil. The estimates are modified by the presence of rock fragments.

Erosion factor Kf indicates the erodibility of the fine-earth fraction, or the material less than 2 millimeters in size.

Erosion factor T is an estimate of the maximum average annual rate of soil erosion by wind or water that can occur without affecting crop productivity over a sustained period. The rate is in tons per acre per year.

Wind erodibility groups are made up of soils that have similar properties affecting their susceptibility to wind erosion in cultivated areas. The soils assigned to group 1 are the most susceptible to wind erosion, and those assigned to group 8 are the least susceptible. The groups are as follows:

1. Coarse sands, sands, fine sands, and very fine sands.

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					

2. Loamy coarse sands, loamy sands, loamy fine sands, loamy very fine sands, ash material, and sapric soil material.

3. Coarse sandy loams, sandy loams, fine sandy loams, and very fine sandy loams.

4L. Calcareous loams, silt loams, clay loams, and silty clay loams.

4. Clays, silty clays, noncalcareous clay loams, and silty clay loams that are more than 35 percent clay.

5. Noncalcareous loams and silt loams that are less than 20 percent clay and sandy clay loams, sandy clays, and hemic soil material.

6. Noncalcareous loams and silt loams that are more than 20 percent clay and noncalcareous clay loams that are less than 35 percent clay.

7. Silts, noncalcareous silty clay loams that are less than 35 percent clay, and fibric soil material.

8. Soils that are not subject to wind erosion because of coarse fragments on the surface or because of surface wetness.

Wind erodibility index is a numerical value indicating the susceptibility of soil to wind erosion, or the tons per acre per year that can be expected to be lost to wind erosion. There is a close correlation between wind erosion and the texture of the surface layer, the size and durability of surface clods, rock fragments, organic matter, and a calcareous reaction. Soil moisture and frozen soil layers also influence wind erosion.

PHYSICAL PROPERTIES OF THE SOILS
Boyd County, Nebraska: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
Ab:														
Albaton-----	0-7	2	40	42-60	1.35-1.40	0.00-0.06	0.11-0.13	9.0-25.0	2.0-3.0	.28	.28	5	4	86
	7-54	3	37	50-73	1.35-1.45	0.00-0.06	0.11-0.13	9.0-25.0	0.9-1.2	.28	.28			
	54-60	10	48	23-54	---	0.00-0.06	0.11-0.13	9.0-25.0	0.9-1.2	.28	.28			
AnC:														
Anselmo-----	0-16	66	20	10-18	1.30-1.60	1.98-5.95	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	16-21	66	20	10-18	1.40-1.60	2.00-6.00	0.15-0.19	0.0-2.9	0.5-1.0	.20	.20			
	21-34	66	20	10-18	1.40-1.60	2.00-6.00	0.12-0.16	0.0-2.9	0.5-1.0	.24	.24			
	34-60	86	7	5-10	1.50-1.75	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.17	.17			
AnD:														
Anselmo-----	0-16	66	20	10-18	1.30-1.60	1.98-5.95	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	16-21	66	20	10-18	1.40-1.60	2.00-6.00	0.15-0.19	0.0-2.9	0.5-1.0	.20	.20			
	21-34	66	20	10-18	1.40-1.60	2.00-6.00	0.12-0.16	0.0-2.9	0.5-1.0	.24	.24			
	34-60	86	7	5-10	1.50-1.75	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.17	.17			
AnF:														
Anselmo-----	0-16	66	20	10-18	1.30-1.60	1.98-5.95	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	16-21	66	20	10-18	1.40-1.60	2.00-6.00	0.15-0.19	0.0-2.9	0.5-1.0	.20	.20			
	21-34	66	20	10-18	1.40-1.60	2.00-6.00	0.12-0.16	0.0-2.9	0.5-1.0	.24	.24			
	34-60	86	7	5-10	1.50-1.75	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.17	.17			
ArF:														
Anselmo-----	0-16	66	20	10-18	1.30-1.60	1.98-5.95	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	16-21	66	20	10-18	1.40-1.60	2.00-6.00	0.15-0.19	0.0-2.9	0.5-1.0	.20	.20			
	21-34	66	20	10-18	1.40-1.60	2.00-6.00	0.12-0.16	0.0-2.9	0.5-1.0	.24	.24			
	34-60	86	7	5-10	1.50-1.75	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.17	.17			
Rock Outcrop-														
Ba:	0-60				---	---	0.00-0.00			---	---		8	0
Barney-----	0-5	27	54	12-27	1.30-1.50	0.57-1.98	0.20-0.23	0.0-2.9	2.0-4.0	.32	.32	5	8	0
	5-10	55	35	5-15	1.60-1.80	1.98-19.98	0.09-0.14	0.0-2.9	0.0-0.5	.17	.17			
	10-60	91	6	0-5	1.70-1.90	5.95-19.98	0.04-0.07	0.0-2.9	0.0-0.5	.10	.10			
Bd:														
Blake-----	0-8	7	61	27-35	1.25-1.30	0.06-0.60	0.20-0.22	3.0-5.9	1.0-3.0	.32	.32	5	4L	86
	8-25	7	64	22-35	1.25-1.30	0.20-2.00	0.20-0.22	3.0-5.9	0.0-1.0	.43	.43			
	25-60	59	26	10-20	1.30-1.35	0.60-2.00	0.20-0.22	0.0-2.9	0.0-0.5	.43	.43			
Be:														
Blendon-----	0-16	66	20	10-18	1.25-1.35	2.00-6.00	0.11-0.17	0.0-2.9	2.0-4.0	.20	.20	5	3	86
	16-42	67	20	10-15	1.25-1.35	2.00-6.00	0.09-0.15	0.0-2.9	0.5-1.0	.20	.20			
	42-60	62	26	5-18	1.30-1.45	1.98-19.98	0.08-0.15	0.0-2.9	0.0-0.5	.17	.17			
BeC:														
Blendon-----	0-16	66	20	10-18	1.25-1.35	2.00-6.00	0.11-0.17	0.0-2.9	2.0-4.0	.20	.20	5	3	86
	16-42	67	20	10-15	1.25-1.35	2.00-6.00	0.09-0.15	0.0-2.9	0.5-1.0	.20	.20			
	42-60	62	26	5-18	1.30-1.45	1.98-19.98	0.08-0.15	0.0-2.9	0.0-0.5	.17	.17			
BoD:														
Boyd-----	0-6	3	45	45-60	1.15-1.30	0.00-0.06	0.10-0.14	9.0-25.0	2.0-4.0	.37	.37	3	4	86
	6-22	17	28	50-60	1.20-1.35	0.00-0.06	0.08-0.12	9.0-25.0	1.0-4.0	.37	.37			
	22-32	18	29	45-60	1.25-1.40	0.01-0.06	0.08-0.12	9.0-25.0	1.0-2.0	.37	.37			
	32-60			---	---	0.00-0.00	---	---	0.0-0.5	---	---			
BrG:														
Bristow-----	0-7	3	45	40-65	1.20-1.40	0.00-0.06	0.10-0.12	6.0-8.9	1.0-3.0	.43	.43	2	4	86
	7-17	18	29	40-65	1.20-1.40	0.00-0.06	0.08-0.10	6.0-8.9	0.0-0.5	.43	.43			
	17-80			---	---	0.06-0.20	---	---	---	---	---			
Bs:														
Brocksburg---	0-18	64	26	7-15	1.40-1.60	2.00-6.00	0.16-0.18	0.0-2.9	1.0-2.0	.20	.20	4	3	86
	18-32	34	38	20-30	1.30-1.50	0.20-2.00	0.15-0.19	3.0-5.9	1.0-2.0	.28	.28			
	32-60	96	2	0-5	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.10	.15			
Bt:														
Brocksburg---	0-14	44	41	10-20	1.30-1.50	0.60-2.00	0.20-0.24	0.0-2.9	1.0-3.0	.28	.28	4	5	56
	14-34	34	38	20-30	1.30-1.50	0.20-2.00	0.15-0.19	3.0-5.9	1.0-2.0	.28	.28			
	34-60	96	2	0-5	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.05	.10			
Cb:														
Cass-----	0-12	68	20	7-17	1.40-1.60	2.00-6.00	0.16-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	12-26	64	26	5-15	1.40-1.60	2.00-6.00	0.15-0.17	0.0-2.9	0.5-1.0	.20	.20			
	26-60	91	6	2-10	1.50-1.70	5.95-19.98	0.08-0.10	0.0-2.9	0.0-0.5	.10	.10			
CrE2:														
Crofton-----	0-6	9	67	20-27	1.20-1.30	0.60-2.00	0.21-0.24	0.0-2.9	0.5-2.0	.43	.43	5	4L	86
	6-60	11	68	15-27	1.10-1.20	0.60-2.00	0.18-0.22	0.0-2.9	0.0-0.5	.43	.43			
DuB:														
Dunday-----	0-15	87	7	3-10	1.40-1.60	5.95-19.98	0.10-0.12	0.0-2.9	1.0-2.0	.15	.15	5	2	134
	15-60	95	1	2-7	1.50-1.70	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.17	.17			
DuC:														
Dunday-----	0-15	87	7	3-10	1.40-1.60	5.95-19.98	0.10-0.12	0.0-2.9	1.0-2.0	.17	.17	5	2	134
	15-60	95	1	2-7	1.50-1.70	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
DuD:														
Dunday-----	0-15	87	7	3-10	1.40-1.60	5.95-19.98	0.10-0.12	0.0-2.9	1.0-2.0	.17	.17	5	2	134
	15-60	95	1	2-7	1.50-1.70	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
DxB:														
Dunday-----	0-14	78	16	3-8	1.50-1.70	5.95-19.98	0.10-0.12	0.0-2.9	1.0-3.0	.17	.17	5	2	134
	14-34	85	9	3-8	1.50-1.70	5.95-19.98	0.05-0.10	0.0-2.9	0.5-1.0	.17	.17			
	34-48	67	23	5-15	1.40-1.60	2.00-6.00	0.11-0.13	0.0-2.9	0.0-0.5	.20	.20			
	48-60	90	6	2-5	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.5-1.0	.10	.10			
Et:														
Eltree-----	0-25	12	69	12-27	1.25-1.35	0.60-2.00	0.20-0.24	0.0-2.9	1.0-3.0	.32	.32	5	6	48
	25-38	10	68	18-27	1.30-1.45	0.60-2.00	0.17-0.22	0.0-2.9	---	.43	.43			
	38-60	9	64	18-27	1.35-1.45	0.60-2.00	0.17-0.20	0.0-2.9	---	.43	.43			

PHYSICAL PROPERTIES OF THE SOILS
Boyd County, Nebraska: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
Go: Grigston-----	0-15	9	67	21-26	1.30-1.40	0.60-2.00	0.21-0.24	0.0-2.9	2.0-4.0	.32	.32	5	6	48
	15-31	9	65	21-30	1.35-1.45	0.60-2.00	0.18-0.22	0.0-2.9	1.0-2.0	.32	.32			
	31-60	32	42	21-30	1.30-1.45	0.60-2.00	0.16-0.22	0.0-2.9	0.0-0.5	.32	.32			
GrB: Aowa-----	0-15	11	67	16-27	1.20-1.40	0.60-2.00	0.22-0.24	0.0-2.9	2.0-4.0	.32	.32	5	4L	86
	15-60	9	66	20-35	1.15-1.40	0.60-2.00	0.17-0.22	0.0-2.9	0.0-0.5	.43	.43			
Ha: Hall-----	0-5	11	68	15-27	1.30-1.40	0.60-2.00	0.20-0.24	3.0-5.9	2.0-4.0	.32	.32	5	6	48
	5-36	7	65	20-35	1.30-1.50	0.20-2.00	0.18-0.20	3.0-5.9	1.0-2.0	.43	.43			
	36-42	7	65	15-30	1.30-1.40	0.20-2.00	0.18-0.22	3.0-5.9	0.5-1.0	.43	.43			
	42-60	12	60	12-30	1.40-1.60	0.20-5.95	0.14-0.20	0.0-2.9	0.0-0.5	.43	.43			
He: Haynie-----	0-7	11	69	15-25	1.20-1.35	0.60-2.00	0.18-0.23	0.0-2.9	1.0-3.0	.32	.32	5	4L	86
	7-60	14	70	15-18	1.20-1.35	0.60-2.00	0.18-0.23	0.0-2.9	0.0-1.0	.43	.43			
IfD: Inavale-----	0-6	96	1	1-5	1.50-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	220
	6-12	92	1	3-10	1.50-1.60	5.95-19.98	0.06-0.11	0.0-2.9	0.0-0.5	.15	.15			
	12-60	92	1	3-10	1.50-1.60	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
IgB: Inavale-----	0-6	96	1	1-5	1.50-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	220
	6-12	92	1	3-10	1.50-1.60	5.95-19.98	0.06-0.11	0.0-2.9	0.0-0.5	.15	.15			
	12-60	92	1	3-10	1.50-1.60	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
IhB: Inavale-----	0-9	87	7	2-10	1.50-1.60	5.95-19.98	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	9-19	92	1	3-10	1.50-1.60	5.95-19.98	0.06-0.11	0.0-2.9	0.0-0.5	.15	.15			
	19-60	92	1	3-10	1.50-1.60	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.17	.17			
In: Inavale-----	0-16	63	26	7-15	1.40-1.50	2.00-6.00	0.13-0.18	0.0-2.9	0.5-1.0	.24	.24	5	3	86
	16-26	92	1	3-10	1.50-1.60	5.95-19.98	0.06-0.11	0.0-2.9	0.0-0.5	.15	.15			
	26-60	92	1	3-10	1.50-1.60	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
Jn: Jansen-----	0-13	43	40	12-22	1.20-1.30	0.60-2.00	0.20-0.24	0.0-2.9	2.0-4.0	.28	.28	4	5	56
	13-32	34	38	18-32	1.30-1.50	0.20-2.00	0.15-0.19	3.0-5.9	0.5-2.0	.37	.37			
	32-60	92	7	0-3	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.05	.10			
JnC: Jansen-----	0-13	43	40	12-22	1.20-1.30	0.60-2.00	0.20-0.24	0.0-2.9	2.0-4.0	.28	.28	4	5	56
	13-32	34	38	18-32	1.30-1.50	0.20-2.00	0.15-0.19	3.0-5.9	0.5-2.0	.37	.37			
	32-60	92	7	0-3	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.05	.10			
JnD: Jansen-----	0-13	43	40	12-22	1.20-1.30	0.60-2.00	0.20-0.24	0.0-2.9	2.0-4.0	.28	.28	4	5	56
	13-32	34	38	18-32	1.30-1.50	0.20-2.00	0.15-0.19	3.0-5.9	0.5-2.0	.37	.37			
	32-60	92	7	0-3	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.05	.10			
LaD: Labu-----	0-6	3	45	45-60	1.20-1.30	0.00-0.06	0.08-0.14	6.0-8.9	1.0-3.0	.32	.32	3	4	86
	6-34	18	29	45-60	1.20-1.30	0.00-0.06	0.08-0.14	6.0-8.9	0.5-1.0	.32	.32			
	34-60			---	---	0.06-0.20	---	---	---	---	---			
LcF: Labu-----	0-6	3	45	45-60	1.20-1.30	0.00-0.06	0.08-0.14	6.0-8.9	1.0-3.0	.32	.32	3	4	86
	6-34	18	29	45-60	1.20-1.30	0.00-0.06	0.08-0.14	6.0-8.9	0.5-1.0	.32	.32			
	34-60			---	---	0.06-0.20	---	---	---	---	---			
Sansarc-----	0-14	5	41	50-65	1.10-1.20	0.06-0.20	0.06-0.12	9.0-25.0	0.1-1.0	.37	.37	2	4	86
	14-60			---	---	---	---	---	---	---	---			
Le: Leshara-----	0-21	11	68	15-27	1.30-1.50	0.60-2.00	0.20-0.24	0.0-2.9	1.0-3.0	.32	.32	5	6	48
	21-32	12	69	12-27	1.30-1.50	0.60-2.00	0.20-0.22	0.0-2.9	0.5-1.0	.43	.43			
	32-60	9	66	18-32	1.30-1.50	0.60-2.00	0.20-0.22	0.0-2.9	0.0-0.5	.43	.43			
LsC: Lynch-----	0-8	5	45	40-60	1.10-1.30	0.00-0.06	0.12-0.14	6.0-8.9	1.0-3.0	.32	.32	3	4	86
	8-28	2	42	45-65	1.20-1.40	0.00-0.06	0.09-0.13	6.0-8.9	0.5-1.0	.32	.32			
	28-36	3	45	45-68	1.20-1.40	0.00-0.06	0.09-0.17	6.0-8.9	0.5-1.0	.32	.32			
	36-60			---	---	0.06-0.20	---	---	---	---	---			
LsD: Lynch-----	0-8	5	45	40-60	1.10-1.30	0.00-0.06	0.12-0.14	6.0-8.9	1.0-3.0	.32	.32	3	4	86
	8-28	2	42	45-65	1.20-1.40	0.00-0.06	0.09-0.13	6.0-8.9	0.5-1.0	.32	.32			
	28-36	3	45	45-68	1.20-1.40	0.00-0.06	0.09-0.17	6.0-8.9	0.5-1.0	.32	.32			
	36-60			---	---	0.06-0.20	---	---	---	---	---			
LyD: Lynch-----	0-8	5	45	40-60	1.10-1.30	0.00-0.06	0.12-0.14	6.0-8.9	1.0-3.0	.32	.32	3	4	86
	8-28	2	42	45-65	1.20-1.40	0.00-0.06	0.09-0.13	6.0-8.9	0.5-1.0	.32	.32			
	28-36	3	45	45-68	1.20-1.40	0.00-0.06	0.09-0.17	6.0-8.9	0.5-1.0	.32	.32			
	36-60			---	---	0.06-0.20	---	---	---	---	---			
Bristow-----	0-7	3	45	40-65	1.20-1.40	0.00-0.06	0.10-0.12	6.0-8.9	1.0-3.0	.43	.43	2	4	86
	7-17	18	29	40-65	1.20-1.40	0.00-0.06	0.08-0.10	6.0-8.9	0.0-0.5	.43	.43			
	17-60			---	---	0.06-0.20	---	---	---	---	---			
LyF: Lynch-----	0-8	5	45	40-60	1.10-1.30	0.00-0.06	0.12-0.14	6.0-8.9	1.0-3.0	.32	.32	3	4	86
	8-28	2	42	45-65	1.20-1.40	0.00-0.06	0.09-0.13	6.0-8.9	0.5-1.0	.32	.32			
	28-36	3	45	45-68	1.20-1.40	0.00-0.06	0.09-0.17	6.0-8.9	0.5-1.0	.32	.32			
	36-60			---	---	0.06-0.20	---	---	---	---	---			
Bristow-----	0-7	3	45	40-65	1.20-1.40	0.00-0.06	0.10-0.12	6.0-8.9	1.0-3.0	.43	.43	2	4	86
	7-17	18	29	40-65	1.20-1.40	0.00-0.06	0.08-0.10	6.0-8.9	0.0-0.5	.43	.43			
	17-60			---	---	0.06-0.20	---	---	---	---	---			

PHYSICAL PROPERTIES OF THE SOILS
Boyd County, Nebraska: Published

PAGE 5 OF 6

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct	K	Kf	T		
MaG: Mariaville----	0-5 5-17 17-60	43 20 20	38 52 52	12-26 12-30 ---	1.20-1.40 1.10-1.30 ---	0.60-2.00 0.20-2.00 0.06-0.20	0.20-0.22 0.17-0.22 ---	0.0-2.9 0.0-2.9 ---	1.0-3.0 0.5-1.0 0.5-1.0	.37 .43 .43	.37 .43 .43	2	4L	86
Paka-----	0-8 8-26 26-48 48-60	38 27 10 68	43 44 68 68	14-25 24-35 10-35 ---	1.20-1.40 1.40-1.60 1.20-1.40 ---	0.60-2.00 0.20-2.00 0.60-2.00 0.06-0.20	0.20-0.22 0.18-0.20 0.20-0.22 ---	3.0-5.9 3.0-5.9 3.0-5.9 ---	2.0-4.0 1.0-2.0 0.5-1.0 ---	.28 .43 .43 ---	.28 .43 .43 ---	4	5	56
MeE: Meadin-----	0-8 8-13 13-60	68 65 90	24 23 6	5-12 5-18 2-5	1.50-1.60 1.50-1.60 1.50-1.70	1.98-5.95 1.98-19.98 19.98-19.98	0.13-0.18 0.09-0.11 0.02-0.05	0.0-2.9 0.0-2.9 0.0-2.9	1.0-2.0 0.5-1.0 0.0-0.5	.20 .17 .05	.20 .28 .10	5	3	86
NoC: Nora-----	0-7 7-34 34-60	9 9 9	67 64 67	20-27 20-35 18-30	1.20-1.30 1.25-1.35 1.30-1.45	0.60-2.00 0.20-2.00 0.20-2.00	0.19-0.22 0.17-0.20 0.17-0.20	0.0-2.9 3.0-5.9 3.0-5.9	2.0-4.0 0.5-1.0 0.0-1.0	.32 .43 .43	.32 .43 .43	5	6	48
NoD: Nora-----	0-7 7-34 34-60	9 9 9	67 64 67	20-27 20-35 18-30	1.20-1.30 1.25-1.35 1.30-1.45	0.60-2.00 0.20-2.00 0.20-2.00	0.19-0.22 0.17-0.20 0.17-0.20	0.0-2.9 3.0-5.9 3.0-5.9	2.0-4.0 0.5-1.0 0.0-1.0	.32 .43 .43	.32 .43 .43	5	6	48
Oa: Onawa-----	0-7 7-28 28-60	6 2 30	47 42 55	40-55 50-60 12-18	1.30-1.35 1.30-1.40 1.40-1.50	0.00-0.06 0.00-0.06 0.57-5.95	0.12-0.14 0.12-0.14 0.20-0.22	6.0-8.9 6.0-8.9 0.0-2.9	2.0-3.0 0.0-1.0 0.0-0.5	.32 .32 .43	.32 .32 .43	5	4	86
Oe: O'Neill-----	0-9 9-34 34-60	64 68 97	27 20 2	7-12 7-18 0-3	1.60-1.80 1.60-1.80 1.50-1.70	1.98-5.95 2.00-6.00 19.98-19.98	0.10-0.15 0.10-0.15 0.02-0.04	0.0-2.9 0.0-2.9 0.0-2.9	1.0-3.0 0.5-1.0 0.0-0.5	.20 .20 .10	.20 .20 .10	4	3	86
OeC: O'Neill-----	0-9 9-34 34-60	64 68 97	27 20 2	7-12 7-18 0-3	1.60-1.80 1.60-1.80 1.50-1.70	1.98-5.95 2.00-6.00 19.98-19.98	0.10-0.15 0.10-0.15 0.02-0.04	0.0-2.9 0.0-2.9 0.0-2.9	1.0-3.0 0.5-1.0 0.0-0.5	.20 .20 .10	.20 .20 .15	4	3	86
OFD: O'Neill-----	0-9 9-34 34-60	64 68 97	27 20 2	7-12 7-18 0-3	1.60-1.80 1.60-1.80 1.50-1.70	1.98-5.95 2.00-6.00 19.98-19.98	0.10-0.15 0.10-0.15 0.02-0.04	0.0-2.9 0.0-2.9 0.0-2.9	1.0-3.0 0.5-1.0 0.0-0.5	.20 .20 .10	.20 .20 .15	4	3	86
Meadin-----	0-8 8-13 13-60	65 65 90	27 23 6	7-12 7-18 2-5	1.50-1.60 1.50-1.60 1.50-1.70	2.00-5.95 2.00-19.98 19.98-19.98	0.13-0.18 0.09-0.11 0.02-0.05	0.0-2.9 0.0-2.9 0.0-2.9	1.0-2.0 0.5-1.0 0.0-0.5	.20 .17 .05	.20 .28 .10	5	3	86
On: Onita-----	0-18 18-32 32-60	25 8 20	52 54 48	20-26 35-50 25-40	1.15-1.30 1.20-1.40 1.25-1.40	0.60-2.00 0.00-0.20 0.06-2.00	0.19-0.22 0.11-0.17 0.17-0.20	0.0-2.9 6.0-8.9 3.0-5.9	4.0-6.0 0.0-2.0 0.0-0.5	.28 .28 .43	.28 .28 .43	5	6	48
Poorly Drained-----	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Or: Ord-----	0-11 11-30 30-60	62 62 92	26 26 3	8-15 8-15 3-12	1.40-1.60 1.50-1.70 1.60-1.70	2.00-6.00 2.00-6.00 5.95-19.98	0.16-0.24 0.15-0.17 0.02-0.04	0.0-2.9 0.0-2.9 0.0-2.9	1.0-2.0 0.5-1.0 0.0-0.5	.20 .20 .15	.20 .20 .15	4	3	86
PaC: Paka-----	0-9 9-27 27-46 46-60	65 27 10 68	20 44 68 68	10-20 24-35 10-35 ---	1.30-1.50 1.40-1.60 1.20-1.40 ---	2.00-6.00 0.20-2.00 0.20-2.00 0.06-0.20	0.13-0.18 0.18-0.20 0.20-0.22 ---	0.0-2.9 3.0-5.9 3.0-5.9 ---	1.0-3.0 1.0-2.0 0.5-1.0 ---	.20 .43 .43 ---	.20 .43 .43 ---	4	3	86
Ph: Paka-----	0-8 8-26 26-48 48-60	38 7 10 68	43 64 68 68	14-25 24-35 10-35 ---	1.20-1.40 1.40-1.60 1.20-1.40 ---	0.60-2.00 0.20-0.60 0.20-2.00 0.06-0.20	0.20-0.22 0.18-0.20 0.20-0.22 ---	3.0-5.9 3.0-5.9 3.0-5.9 ---	2.0-4.0 1.0-2.0 0.5-1.0 ---	.28 .43 .43 ---	.28 .43 .43 ---	4	5	56
PhC: Paka-----	0-8 8-26 26-48 48-60	38 7 10 68	43 64 68 68	14-25 24-35 10-35 ---	1.20-1.40 1.40-1.60 1.20-1.40 ---	0.60-2.00 0.20-0.60 0.20-2.00 0.06-0.20	0.20-0.22 0.18-0.20 0.20-0.22 ---	3.0-5.9 3.0-5.9 3.0-5.9 ---	2.0-4.0 1.0-2.0 0.5-1.0 ---	.28 .43 .43 ---	.28 .43 .43 ---	4	5	56
PhD: Paka-----	0-8 8-26 26-48 48-60	38 7 10 68	43 64 68 68	14-25 24-35 10-35 ---	1.20-1.40 1.40-1.60 1.20-1.40 ---	0.60-2.00 0.20-0.60 0.20-2.00 0.06-0.20	0.20-0.22 0.18-0.20 0.20-0.22 ---	3.0-5.9 3.0-5.9 3.0-5.9 ---	2.0-4.0 1.0-2.0 0.5-1.0 ---	.28 .43 .43 ---	.28 .43 .43 ---	4	5	56
PoC: Promise-----	0-18 18-33 33-60	2 11 16	42 26 26	50-60 60-65 50-65	1.10-1.25 1.10-1.25 1.10-1.25	0.00-0.06 0.00-0.06 0.00-0.06	0.10-0.14 0.08-0.14 0.10-0.12	9.0-25.0 9.0-25.0 9.0-25.0	2.0-4.0 1.0-3.0 0.0-0.5	.37 .37 .37	.37 .37 .37	5	4	86
RaC: Ree-----	0-7 7-22 22-60	24 35 67	52 34 15	22-26 27-35 5-25	1.15-1.30 1.20-1.35 1.30-1.50	0.60-2.00 0.20-0.60 0.60-19.98	0.18-0.22 0.17-0.22 0.09-0.20	0.0-2.9 3.0-5.9 0.0-2.9	2.0-4.0 0.5-2.0 0.0-0.5	.32 .28 .28	.32 .28 .28	5	6	48
RaD: Ree-----	0-7 7-22 22-60	24 35 67	52 34 15	22-26 27-35 10-25	1.15-1.30 1.20-1.35 1.30-1.50	0.60-2.00 0.20-2.00 0.60-5.95	0.18-0.22 0.17-0.22 0.09-0.20	0.0-2.9 3.0-5.9 0.0-2.9	2.0-4.0 0.5-2.0 0.0-0.5	.32 .28 .28	.32 .28 .28	5	6	48
RaE: Ree-----	0-7 7-22 22-60	24 55 57	52 14 18	22-26 27-35 10-25	1.15-1.30 1.20-1.35 1.30-1.50	0.60-2.00 0.20-2.00 0.60-5.95	0.18-0.22 0.17-0.22 0.09-0.20	0.0-2.9 3.0-5.9 0.0-2.9	2.0-4.0 0.5-2.0 0.0-0.5	.32 .28 .28	.32 .28 .28	5	6	48

PHYSICAL PROPERTIES OF THE SOILS
Boyd County, Nebraska: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
ReC: Reliance-----	0-10	24	52	22-26	1.10-1.25	0.60-2.00	0.19-0.22	0.0-2.9	2.0-4.0	.32	.32	4	6	48
	10-28	7	53	35-45	1.20-1.40	0.06-0.20	0.11-0.19	6.0-8.9	0.5-2.0	.32	.32			
	28-60	20	48	25-40	1.20-1.35	0.20-2.00	0.14-0.20	3.0-5.9	0.0-1.0	.43	.43			
ReD: Reliance-----	0-10	24	52	22-26	1.10-1.25	0.60-2.00	0.19-0.22	0.0-2.9	2.0-4.0	.32	.32	4	6	48
	10-28	7	53	35-45	1.20-1.40	0.06-0.20	0.11-0.19	6.0-8.9	0.5-2.0	.32	.32			
	28-60	20	48	25-40	1.20-1.35	0.20-2.00	0.14-0.20	3.0-5.9	0.0-1.0	.43	.43			
RfC: Reliance-----	0-10	20	49	27-35	1.15-1.25	0.20-0.60	0.19-0.22	3.0-5.9	2.0-4.0	.32	.32	4	7	38
	10-28	7	53	35-45	1.20-1.40	0.20-0.60	0.11-0.19	6.0-8.9	0.5-2.0	.32	.32			
	28-60	20	48	25-40	1.20-1.35	0.20-2.00	0.14-0.20	3.0-5.9	0.0-1.0	.43	.43			
Rw: Riverwash----	0-6	95	1	1-8	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.0-0.1	.15	.15	5	8	0
	6-60	97	0	0-5	1.65-1.85	5.95-19.98	0.02-0.03	0.0-2.9	0.0-0.5	.15	.15			
SaG: Sansarc-----	0-16	5	35	55-65	1.10-1.20	0.06-0.20	0.06-0.12	9.0-25.0	0.1-1.0	.37	.37	2	4	86
	16-60			---	---	---	---	---	---	---	---			
Sc: Scott-----	0-8	26	53	15-27	1.25-1.40	0.60-2.00	0.21-0.24	0.0-2.9	2.0-4.0	.37	.37	3	6	48
	8-48	6	47	40-55	1.20-1.40	0.00-0.06	0.08-0.16	6.0-8.9	1.0-2.0	.37	.37			
	48-56	19	48	27-40	1.15-1.40	0.06-0.60	0.18-0.20	6.0-8.9	0.5-1.0	.43	.43			
	56-64	24	50	18-35	1.30-1.50	0.20-2.00	0.14-0.22	3.0-5.9	0.5-1.0	.43	.43			
Sm: Simeon-----	0-5	84	9	5-10	1.30-1.50	5.95-19.98	0.08-0.14	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	5-60	92	2	2-10	1.50-1.70	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
SuC: Simeon-----	0-15	84	9	5-10	1.30-1.50	5.95-19.98	0.08-0.14	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	15-60	92	2	2-10	1.50-1.70	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.10	.10			
Valentine----	0-5	85	9	2-10	1.35-1.55	5.95-19.98	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	5-11	93	1	2-10	1.55-1.75	5.95-19.98	0.09-0.11	0.0-2.9	0.0-0.5	.15	.15			
	11-60	96	1	0-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
SvF2: Simeon-----	0-15	84	9	5-10	1.30-1.50	5.95-19.98	0.08-0.14	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	15-60	92	2	2-10	1.50-1.70	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.10	.10			
Valentine----	0-5	96	1	0-6	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	5-11	93	1	2-10	1.55-1.75	5.95-19.98	0.09-0.11	0.0-2.9	0.0-0.5	.15	.15			
	11-60	96	1	0-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
VaE: Valentine----	0-5	96	1	0-6	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	5-11	93	1	2-10	1.55-1.75	5.95-19.98	0.09-0.11	0.0-2.9	0.0-0.5	.15	.15			
	11-60	96	1	0-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
VbB: Valentine----	0-5	85	9	2-10	1.35-1.55	5.95-19.98	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	5-11	87	7	2-10	1.55-1.75	5.95-19.98	0.09-0.11	0.0-2.9	0.0-0.5	.17	.17			
	11-60	96	1	0-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
Ve: Verdel-----	0-14	5	45	40-60	1.20-1.40	0.00-0.06	0.12-0.15	6.0-8.9	1.0-3.0	.28	.28	5	4	86
	14-60	5	45	40-60	1.30-1.50	0.00-0.06	0.09-0.14	6.0-8.9	0.5-1.0	.32	.32			
WeC: Wewela-----	0-8	66	20	10-18	1.25-1.40	2.00-6.00	0.14-0.17	0.0-2.9	2.0-4.0	.24	.24	3	3	86
	8-15	56	14	25-35	1.30-1.45	0.20-0.60	0.16-0.18	3.0-5.9	1.0-3.0	.32	.32			
	15-22	22	28	40-60	1.25-1.40	0.00-0.06	0.08-0.12	6.0-8.9	0.0-1.0	.37	.37			
	22-60			---	---	0.06-0.20	---	---	0.0-0.5	---	---			
zwa: Water-----	---			---	---	---	---	---	---	---	---	-	---	---

CHEMICAL PROPERTIES OF THE SOILS
Boyd County, Nebraska

The Chemical Properties table shows estimates of some characteristics and features that affect soil behavior. These estimates are given for the major layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils. Depth to the upper and lower boundaries of each layer is indicated.

Cation-exchange capacity is the total amount of extractable bases that can be held by the soil, expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. Soils having a low cation-exchange capacity hold fewer cations and may require more frequent applications of fertilizer than soils having a high cation-exchange capacity. Soils having a high cation-exchange capacity can retain cations. The ability to retain cations helps to prevent the pollution of ground water.

Soil reaction is a measure of acidity or alkalinity and is expressed as a range in pH values. The range in pH of each major horizon is based on many field tests. For many soils, values have been verified by laboratory analyses. Soil reaction is important in selecting crops and other plants, in evaluating soil amendments for fertility and stabilization, and in determining the risk of corrosion.

Calcium carbonate equivalent is the percent of carbonates, by weight, in the fraction of the soil less than 2 millimeters in size. The availability of plant nutrients is influenced by the amount of carbonates in the soil. Incorporating nitrogen fertilizer into calcareous soils helps to prevent nitrite accumulation and ammonium-N volatilization.

Gypsum is expressed as a percent, by weight, of hydrated calcium sulfates in the fraction of the soil less than 20 millimeters in size. Gypsum is partially soluble in water and can be dissolved and removed by water. Soils that have a high content of gypsum may collapse if the gypsum is removed by percolating water.

Salinity is a measure of soluble salts in the soil at saturation. It is expressed as the electrical conductivity of the saturation extract, in millimhos per centimeter at 25 degrees C. Estimates are based on field and laboratory measurements at representative sites of nonirrigated soils. The salinity of irrigated soils is affected by the quality of the irrigation water and by the frequency of water application. Hence, the salinity of soils in individual fields can differ greatly from the value given in the table. Salinity affects the suitability of a soil for crop production, the stability of soil if used as construction material, and the potential of the soil to corrode metal and concrete.

Sodium adsorption ratio (SAR) is a measure of the amount of sodium (Na) relative to calcium (Ca) and magnesium (Mg) in the water extract from saturated soil paste. It is the ratio of the Na concentration divided by the square root of one-half of the Ca + Mg concentration. Soils that have SAR values of 13 or more may be characterized by an increased dispersion of organic matter and clay particles, reduced permeability and aeration, and a general degradation of soil structure.

CHEMICAL PROPERTIES OF THE SOILS--Continued
Boyd County, Nebraska

PAGE 2 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
Ab:							
Albaton-----	0-7	36-41	7.4-8.4	5-30	0	0	0
	7-54	41-50	7.4-8.4	5-30	0	0	0
	54-60	---	---	5-30	---	---	---
AnC:							
Anselmo-----	0-16	10-20	5.6-7.3	0	0	0	0
	16-21	5.0-15	6.1-7.8	0	0	0	0
	21-34	5.0-15	6.6-7.8	0-3	0	0	0
	34-60	0.0-10	6.6-7.8	0-3	0	0	0
AnD:							
Anselmo-----	0-16	10-20	5.6-7.3	0	0	0	0
	16-21	5.0-15	6.1-7.8	0	0	0	0
	21-34	5.0-15	6.6-7.8	0-3	0	0	0
	34-60	0.0-10	6.6-7.8	0-3	0	0	0
AnF:							
Anselmo-----	0-16	10-20	5.6-7.3	0	0	0	0
	16-21	5.0-15	6.1-7.8	0	0	0	0
	21-34	5.0-15	6.6-7.8	0-3	0	0	0
	34-60	0.0-10	6.6-7.8	0-3	0	0	0
ArF:							
Anselmo-----	0-16	10-20	5.6-7.3	0	0	0	0
	16-21	5.0-15	6.1-7.8	0	0	0	0
	21-34	5.0-15	6.6-7.8	0-3	0	0	0
	34-60	0.0-10	6.6-7.8	0-3	0	0	0
Rock Outcrop----	0-60	---	---	---	---	0	---
Ba:							
Barney-----	0-5	15-28	6.6-8.4	0-5	0	0	0
	5-10	2.0-7.0	6.6-8.4	0-5	0	0	0
	10-60	0.0-4.0	6.6-7.8	0	0	0	0
Bd:							
Blake-----	0-8	25-35	7.4-8.4	5-30	0	0	0
	8-25	20-30	7.4-8.4	5-30	0	0	0
	25-60	10-20	7.4-8.4	5-30	0	0	0
Be:							
Blendon-----	0-16	10-20	5.6-7.3	0	0	0.0-2.0	0
	16-42	5.0-10	6.1-8.4	0-5	0	0.0-2.0	0
	42-60	5.0-10	6.6-8.4	0-5	0	0.0-2.0	0
BeC:							
Blendon-----	0-16	10-20	5.6-7.3	0	0	0.0-2.0	0
	16-42	5.0-10	6.1-8.4	0-5	0	0.0-2.0	0
	42-60	5.0-10	6.6-8.4	0-5	0	0.0-2.0	0
BoD:							
Boyd-----	0-6	33-46	6.6-8.4	0-5	0-5	0.0-2.0	0
	6-22	37-46	6.6-8.4	0-5	0-5	0.0-2.0	0
	22-32	30-45	6.6-8.4	0-10	0-5	0.0-2.0	0
	32-60	---	---	---	---	---	---
BrG:							
Bristow-----	0-7	29-48	7.4-8.4	15-25	0-10	0	0
	7-17	29-48	7.4-8.4	15-40	5-15	0	0
	17-80	---	---	---	---	---	---
Bs:							
Brocksburg-----	0-18	5.0-15	6.1-7.3	0	0	0	0
	18-32	15-25	6.6-7.8	0	0	0	0
	32-60	0.0-5.0	6.6-7.8	0	0	0	0
Bt:							
Brocksburg-----	0-14	10-20	6.1-7.3	0	0	0	0
	14-34	15-25	6.6-7.8	0	0	0	0
	34-60	0.0-5.0	6.6-7.8	0	0	0	0
Cb:							
Cass-----	0-12	6.0-14	5.6-7.3	0	0	0	0
	12-26	4.0-12	6.1-8.4	0	0	0	0
	26-60	1.0-8.0	6.1-8.4	0	0	0	0
CrE2:							
Crofton-----	0-6	15-25	7.4-8.4	1-10	0	0	0
	6-60	15-25	7.4-8.4	1-15	0	0	0
DuB:							
Dunday-----	0-15	5.0-10	6.1-7.3	0	0	0	0
	15-60	1.0-5.0	6.1-7.8	0	0	0	0
DuC:							
Dunday-----	0-15	5.0-10	6.1-7.3	0	0	0	0
	15-60	1.0-5.0	6.1-7.8	0	0	0	0

CHEMICAL PROPERTIES OF THE SOILS--Continued
Boyd County, Nebraska

PAGE 3 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
DuD:							
Dunday-----	0-15	5.0-10	6.1-7.3	0	0	0	0
	15-60	1.0-5.0	6.1-7.8	0	0	0	0
DxB:							
Dunday-----	0-14	2.0-10	6.1-7.3	0	0	0	0
	14-34	2.0-10	6.1-7.3	0	0	0	0
	34-48	3.0-15	7.4-7.8	0	0	0	0
	48-60	1.0-5.0	7.4-7.8	0	0	0	0
Et:							
Eltree-----	0-25	---	6.6-8.4	---	---	0	---
	25-38	---	7.4-8.4	---	---	0	---
	38-60	---	7.4-8.4	---	---	0	---
Go:							
Grigston-----	0-15	20-30	6.6-7.8	0	0	0	0
	15-31	15-30	7.4-8.4	5-10	0	0	0
	31-60	10-15	7.4-8.4	5-10	0	0	0
GrB:							
Aowa-----	0-15	15-28	7.4-8.4	0-5	0	0	0
	15-60	18-32	7.4-8.4	0-5	0	0	0
Ha:							
Hall-----	0-5	13-23	6.1-7.3	0	0	0	0
	5-36	15-27	6.1-7.8	0	0	0	0
	36-42	11-22	6.6-7.8	0-5	0	0	0
	42-60	8.0-22	6.6-8.4	1-5	0	0	0
He:							
Haynie-----	0-7	15-20	6.6-8.4	0-25	0	0	0
	7-60	15-20	7.4-8.4	5-30	0	0	0
IfD:							
Inavale-----	0-6	1.0-5.0	5.6-7.8	0	0	0	0
	6-12	1.0-10	5.6-7.8	0	0	0	0
	12-60	1.0-10	5.6-7.8	0	0	0	0
IgB:							
Inavale-----	0-6	1.0-5.0	5.6-7.8	0	0	0	0
	6-12	1.0-10	5.6-7.8	0	0	0	0
	12-60	1.0-10	5.6-7.8	0	0	0	0
IhB:							
Inavale-----	0-9	2.0-10	5.6-7.8	0	0	0	0
	9-19	1.0-10	5.6-7.8	0	0	0	0
	19-60	1.0-10	5.6-7.8	0	0	0	0
In:							
Inavale-----	0-16	5.0-15	5.6-7.8	0	0	0	0
	16-26	0.0-10	5.6-7.8	0	0	0	0
	26-60	0.0-10	5.6-7.8	0	0	0	0
Jn:							
Jansen-----	0-13	10-20	5.1-7.3	0	0	0	0
	13-32	12-25	5.1-7.3	0	0	0	0
	32-60	0.0-2.0	5.1-7.3	0	0	0	0
JnC:							
Jansen-----	0-13	10-20	5.1-7.3	0	0	0	0
	13-32	12-25	5.1-7.3	0	0	0	0
	32-60	0.0-2.0	5.1-7.3	0	0	0	0
JnD:							
Jansen-----	0-13	10-20	5.1-7.3	0	0	0	0
	13-32	12-25	5.1-7.3	0	0	0	0
	32-60	0.0-2.0	5.1-7.3	0	0	0	0
LaD:							
Labu-----	0-6	30-45	6.6-8.4	5-15	0	0	0
	6-34	30-45	7.4-8.4	5-15	0	0	0
	34-60	---	---	---	---	---	---
LcF:							
Labu-----	0-6	30-45	6.6-8.4	5-15	0	0	0
	6-34	30-45	7.4-8.4	5-15	0	0	0
	34-60	---	---	---	---	---	---
Sansarc-----	0-14	35-45	7.4-8.4	1-10	0-2	0.0-2.0	0-1
	14-60	---	---	---	---	---	---
Le:							
Leshara-----	0-21	10-22	6.1-8.4	0-10	0	0.0-2.0	0
	21-32	5.0-20	6.6-8.4	0-10	0	0.0-2.0	0
	32-60	3.0-15	6.6-8.4	0-10	0	0.0-2.0	0

CHEMICAL PROPERTIES OF THE SOILS--Continued
Boyd County, Nebraska

PAGE 4 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
LsC:							
Lynch-----	0-8	20-40	7.4-8.4	30-40	0-10	0	0
	8-28	10-20	7.4-8.4	30-40	0-15	0	0
	28-36	10-20	7.4-9.0	30-40	0-15	0	0
	36-60	---	---	---	---	---	---
LsD:							
Lynch-----	0-8	20-40	7.4-8.4	30-40	0-10	0	0
	8-28	10-20	7.4-8.4	30-40	0-15	0	0
	28-36	10-20	7.4-9.0	30-40	0-15	0	0
	36-60	---	---	---	---	---	---
LyD:							
Lynch-----	0-8	20-40	7.4-8.4	30-40	0-10	0	0
	8-28	10-20	7.4-8.4	30-40	0-15	0	0
	28-36	10-20	7.4-9.0	30-40	0-15	0	0
	36-60	---	---	---	---	---	---
Bristow-----	0-7	29-48	7.4-8.4	15-25	0-10	0	0
	7-17	29-48	7.4-8.4	15-40	5-15	0	0
	17-60	---	---	---	---	---	---
LyF:							
Lynch-----	0-8	20-40	7.4-8.4	30-40	0-10	0	0
	8-28	10-20	7.4-8.4	30-40	0-15	0	0
	28-36	10-20	7.4-9.0	30-40	0-15	0	0
	36-60	---	---	---	---	---	---
Bristow-----	0-7	29-48	7.4-8.4	15-25	0-10	0	0
	7-17	29-48	7.4-8.4	15-40	5-15	0	0
	17-60	---	---	---	---	---	---
MaG:							
Mariaville-----	0-5	10-20	7.4-8.4	3-10	0	0.0-2.0	0
	5-17	10-20	7.4-8.4	8-14	0	0.0-2.0	0
	17-60	---	---	---	---	---	---
Paka-----	0-8	10-45	6.6-7.3	0	0	0	0
	8-26	15-25	6.6-7.8	0	0	0	0
	26-48	10-25	7.4-8.4	0-5	0	0	0
	48-60	---	---	---	---	---	---
MeE:							
Meadin-----	0-8	10-20	5.1-7.3	0	0	0	0
	8-13	5.0-15	5.1-7.3	0	0	0	0
	13-60	0.0-5.0	6.1-7.3	0	0	0	0
NoC:							
Nora-----	0-7	15-25	6.1-7.3	0	0	0.0-2.0	0
	7-34	15-25	6.1-7.8	0-5	0	0.0-2.0	0
	34-60	10-22	6.6-8.4	5-15	0	0.0-2.0	0
NoD:							
Nora-----	0-7	15-25	6.1-7.3	0	0	0.0-2.0	0
	7-34	15-25	6.1-7.8	0-5	0	0.0-2.0	0
	34-60	10-22	6.6-8.4	5-15	0	0.0-2.0	0
Oa:							
Onawa-----	0-7	36-41	7.4-8.4	5-30	0	0	0
	7-28	41-50	7.4-8.4	5-30	0	0	0
	28-60	15-20	7.4-8.4	5-30	0	0	0
Oe:							
O'Neill-----	0-9	10-20	5.1-6.5	0	0	0	0
	9-34	5.0-15	5.6-7.3	0	0	0	0
	34-60	0.0-5.0	5.6-7.3	0	0	0	0
OeC:							
O'Neill-----	0-9	10-20	5.1-6.5	0	0	0	0
	9-34	5.0-15	5.6-7.3	0	0	0	0
	34-60	0.0-5.0	5.6-7.3	0	0	0	0
OfD:							
O'Neill-----	0-9	10-20	5.1-6.5	0	0	0	0
	9-34	5.0-15	5.6-7.3	0	0	0	0
	34-60	0.0-5.0	5.6-7.3	0	0	0	0
Meadin-----	0-8	10-20	5.1-7.3	0	0	0	0
	8-13	5.0-15	5.1-7.3	0	0	0	0
	13-60	0.0-5.0	6.1-7.3	0	0	0	0
On:							
Onita-----	0-18	20-30	5.6-7.3	0	0	0.0-2.0	0
	18-32	25-35	6.1-7.3	0	0	0.0-2.0	0
	32-60	20-30	7.4-8.4	5-15	0-1	0.0-2.0	0-1
Poorly Drained--	---	---	---	---	---	---	---

CHEMICAL PROPERTIES OF THE SOILS--Continued
Boyd County, Nebraska

PAGE 5 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
Or:							
Ord-----	0-11	10-20	7.4-8.4	5-40	0	0.0-2.0	0-5
	11-30	5.0-15	6.6-8.4	0-5	0	0.0-2.0	0-2
	30-60	0.0-10	6.6-8.4	0-5	0	0	0
PaC:							
Paka-----	0-9	10-25	6.6-7.3	0	0	0	0
	9-27	15-25	6.6-7.8	0	0	0	0
	27-46	10-25	7.4-8.4	0-5	0	0	0
	46-60	---	---	---	---	---	---
Ph:							
Paka-----	0-8	10-45	6.6-7.3	0	0	0	0
	8-26	15-25	6.6-7.8	0	0	0	0
	26-48	10-25	7.4-8.4	0-5	0	0	0
	48-60	---	---	---	---	---	---
PhC:							
Paka-----	0-8	10-45	6.6-7.3	0	0	0	0
	8-26	15-25	6.6-7.8	0	0	0	0
	26-48	10-25	7.4-8.4	0-5	0	0	0
	48-60	---	---	---	---	---	---
PhD:							
Paka-----	0-8	10-45	6.6-7.3	0	0	0	0
	8-26	15-25	6.6-7.8	0	0	0	0
	26-48	10-25	7.4-8.4	0-5	0	0	0
	48-60	---	---	---	---	---	---
PoC:							
Promise-----	0-18	40-45	6.1-7.8	0-3	---	0.0-2.0	0-1
	18-33	35-45	7.4-9.0	2-10	0-3	0.0-2.0	1-6
	33-60	35-40	7.4-9.0	2-10	0-10	2.0-4.0	1-6
RaC:							
Ree-----	0-7	15-25	6.1-7.3	0	0	0.0-2.0	0
	7-22	18-27	6.6-8.4	0-10	0	0.0-2.0	0
	22-60	10-25	7.4-8.4	5-15	0	0.0-2.0	0
RaD:							
Ree-----	0-7	15-25	6.1-7.3	0	0	0.0-2.0	0
	7-22	18-27	6.6-8.4	0-10	0	0.0-2.0	0
	22-60	10-25	7.4-8.4	5-15	0	0.0-2.0	0
RaE:							
Ree-----	0-7	15-25	6.1-7.3	0	0	0.0-2.0	0
	7-22	18-27	6.6-8.4	0-10	0	0.0-2.0	0
	22-60	10-25	7.4-8.4	5-15	0	0.0-2.0	0
ReC:							
Reliance-----	0-10	20-25	6.1-7.3	0-1	0	0.0-2.0	0
	10-28	25-35	6.1-7.8	1-10	0	0.0-2.0	0
	28-60	20-30	7.4-8.4	5-20	0	0.0-2.0	0
ReD:							
Reliance-----	0-10	20-25	6.1-7.3	0-1	0	0.0-2.0	0
	10-28	25-35	6.1-7.8	1-10	0	0.0-2.0	0
	28-60	20-30	7.4-8.4	5-20	0	0.0-2.0	0
RfC:							
Reliance-----	0-10	20-30	6.1-7.3	0-1	0	0.0-2.0	0
	10-28	25-35	6.1-7.8	1-10	0	0.0-2.0	0
	28-60	20-30	7.4-8.4	5-20	0	0.0-2.0	0
Rw:							
Riverwash-----	0-6	1.0-10	7.4-8.4	0-5	0	0	0
	6-60	0.0-5.0	7.4-8.4	0-5	0	0	0
SaG:							
Sansarc-----	0-16	35-45	7.4-8.4	1-10	0-2	0.0-2.0	0-1
	16-60	---	---	---	---	---	---
Sc:							
Scott-----	0-8	20-30	5.1-6.5	0	0	0	0
	8-48	20-35	5.6-7.8	0	0	0	0
	48-56	20-35	6.6-8.4	0-5	0	0	0
	56-64	25-35	6.6-8.4	0-5	0	0	0
Sm:							
Simeon-----	0-5	0.0-5.0	6.1-7.8	0	0	0	0
	5-60	0.0-5.0	6.1-7.8	0	0	0	0

CHEMICAL PROPERTIES OF THE SOILS--Continued
Boyd County, Nebraska

PAGE 6 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
SuC:							
Simeon-----	0-15	0.0-5.0	6.1-7.8	0	0	0	0
	15-60	0.0-5.0	6.1-7.8	0	0	0	0
Valentine-----	0-5	2.0-8.0	5.6-7.3	0	0	0	0
	5-11	1.0-8.0	5.6-7.3	0	0	0	0
	11-60	0.0-5.0	5.6-7.3	0	0	0	0
SvF2:							
Simeon-----	0-15	0.0-5.0	6.1-7.8	0	0	0	0
	15-60	0.0-5.0	6.1-7.8	0	0	0	0
Valentine-----	0-5	0.0-6.0	5.6-7.3	0	0	0	0
	5-11	1.0-8.0	5.6-7.3	0	0	0	0
	11-60	0.0-5.0	5.6-7.3	0	0	0	0
VaE:							
Valentine-----	0-5	0.0-6.0	5.6-7.3	0	0	0	0
	5-11	1.0-8.0	5.6-7.3	0	0	0	0
	11-60	0.0-5.0	5.6-7.3	0	0	0	0
VbB:							
Valentine-----	0-5	2.0-8.0	5.6-7.3	0	0	0	0
	5-11	1.0-8.0	5.6-7.3	0	0	0	0
	11-60	0.0-5.0	5.6-7.3	0	0	0	0
Ve:							
Verdel-----	0-14	25-35	6.1-7.8	0	0	0	0-2
	14-60	20-35	6.6-9.0	0-5	0	0	0-2
WeC:							
Wewela-----	0-8	9.0-17	6.1-7.3	0	0	0.0-2.0	0
	8-15	15-26	6.1-7.3	0-10	0	0.0-2.0	0
	15-22	28-43	6.1-8.4	0-10	0	0.0-2.0	0
	22-60	---	---	---	---	---	---
zwa:							
Water-----	---	---	---	---	---	---	---

WATER FEATURES Boyd County, Nebraska

The Water Features table gives estimates of various water features. The estimates are used in land use planning that involves engineering considerations. Hydrologic soil groups are based on estimates of runoff potential. Soils are assigned to one of four groups according to the rate of water infiltration when the soils are not protected by vegetation, are thoroughly wet, and receive precipitation from long-duration storms.

The four hydrologic soil groups are:

Group A. Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravelly sands. These soils have a high rate of water transmission.

Group B. Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C. Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D. Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink-swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

The months in the table indicate the portion of the year in which the feature is most likely to be a concern.

Water table refers to a saturated zone in the soil. The Water Features table indicates, by month, depth to the top (upper limit) and base (lower limit) of the saturated zone in most years. Estimates of the upper and lower limits are based mainly on observations of the water table at selected sites and on evidence of a saturated zone, namely grayish colors or mottles (redoximorphic features) in the soil. A saturated zone that lasts for less than a month is not considered a water table. Ponding is standing water in a closed depression. Unless a drainage system is installed, the water is removed only by percolation, transpiration, or evaporation. The Water Features table indicates surface water depth and the duration and frequency of ponding. Duration is expressed as very brief if less than 2 days, brief if 2 to 7 days, long if 7 to 30 days, and very long if more than 30 days. Frequency is expressed as none, rare, occasional, and frequent. None means that ponding is not probable; rare that it is unlikely but possible under unusual weather conditions (the chance of ponding is nearly 0 percent to 5 percent in any year); occasional that it occurs, on the average, once or less in 2 years (the chance of ponding is 5 to 50 percent in any year); and frequent that it occurs, on the average, more than once in 2 years (the chance of ponding is more than 50 percent in any year).

Flooding, the temporary inundation of an area, is caused by overflowing streams, by runoff from adjacent slopes, or by tides. Water standing for short periods after rainfall or snowmelt is not considered flooding, and water standing in swamps and marshes is considered ponding rather than flooding.

Duration and frequency are estimated. Duration is expressed as extremely brief if 0.1 hour to 4 hours, very brief if 4 hours to 2 days, brief if 2 to 7 days, long if 7 to 30 days, and very long if more than 30 days. Frequency is expressed as none, very rare, rare, occasional, frequent, and very frequent. None means that flooding is not probable; very rare that it is very unlikely but possible under extremely unusual weather conditions (the chance of flooding is less than 1 percent in any year); rare that it is unlikely but possible under unusual weather conditions (the chance of flooding is 1 to 5 percent in any year); occasional that it occurs infrequently under normal weather conditions (the chance of flooding is 5 to 50 percent in any year); frequent that it is likely to occur often under normal weather conditions (the chance of flooding is more than 50 percent in any year but is less than 50 percent in all months in any year); and very frequent that it is likely to occur very often under normal weather conditions (the chance of flooding is more than 50 percent in all months of any year).

The information is based on evidence in the soil profile, namely thin strata of gravel, sand, silt, or clay deposited by floodwater; irregular decrease in organic matter content with increasing depth; and little or no horizon development.

Also considered are local information about the extent and levels of flooding and the relation of each soil on the landscape to historic floods. Information on the extent of flooding based on soil data is less specific than that provided by detailed engineering surveys that delineate flood-prone areas at specific flood frequency levels.

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
Ab: Albaton-----	D	January	0.0-1.0	>6.0	---	---	---	---	None
		February	0.0-1.0	>6.0	---	---	---	Brief	Occasional
		March	0.0-1.0	>6.0	---	---	---	Brief	Occasional
		April	0.0-1.0	>6.0	---	---	---	Brief	Occasional
		May	0.0-1.0	>6.0	---	---	---	Brief	Occasional
		June	0.0-1.0	>6.0	---	---	---	Brief	Occasional
		July	0.0-1.0	>6.0	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		September	---	---	---	---	---	Brief	Occasional
		October	---	---	---	---	---	Brief	Occasional
		November	0.0-1.0	>6.0	---	---	---	Brief	Occasional
		December	0.0-1.0	>6.0	---	---	---	---	None
AnC: Anselmo-----	B		---	---	---	---	---	---	---
AnD: Anselmo-----	B		---	---	---	---	---	---	---
AnF: Anselmo-----	B		---	---	---	---	---	---	---
ArF: Anselmo-----	B		---	---	---	---	---	---	---
Rock Outcrop-----	D		---	---	---	---	---	---	---
Ba: Barney-----	D	January	0.0-1.0	>6.0	---	---	---	---	None
		February	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		March	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		April	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		May	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		June	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		July	---	---	---	---	---	Brief	Frequent
		November	0.0-1.0	>6.0	---	---	---	---	None
		December	0.0-1.0	>6.0	---	---	---	---	None
Bd: Blake-----	B	January	2.0-4.0	>6.0	---	---	---	---	None
		February	2.0-4.0	>6.0	---	---	---	Long	Occasional
		March	2.0-4.0	>6.0	---	---	---	Long	Occasional
		April	2.0-4.0	>6.0	---	---	---	Long	Occasional
		May	2.0-4.0	>6.0	---	---	---	Long	Occasional
		June	2.0-4.0	>6.0	---	---	---	Long	Occasional
		July	2.0-4.0	>6.0	---	---	---	Long	Occasional
		August	---	---	---	---	---	Long	Occasional
		September	---	---	---	---	---	Long	Occasional
		October	---	---	---	---	---	Long	Occasional
		November	2.0-4.0	>6.0	---	---	---	Long	Occasional
		December	2.0-4.0	>6.0	---	---	---	---	None
Be: Blendon-----	B		---	---	---	---	---	---	---
BeC: Blendon-----	B		---	---	---	---	---	---	---
BoD: Boyd-----	D		---	---	---	---	---	---	---
BrG: Bristow-----	D		---	---	---	---	---	---	---
Bs: Brocksburg-----	B		---	---	---	---	---	---	---
Bt: Brocksburg-----	B		---	---	---	---	---	---	---
Cb: Cass-----	B	March	---	---	---	---	---	Brief	Occasional
		April	---	---	---	---	---	Brief	Occasional
		May	---	---	---	---	---	Brief	Occasional
		June	---	---	---	---	---	Brief	Occasional
CrE2: Crofton-----	B		---	---	---	---	---	---	---
DuB: Dunday-----	A		---	---	---	---	---	---	---
DuC:			---	---	---	---	---	---	---

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Dunday-----	A		Ft	Ft	Ft				
DuD: Dunday-----	A		---	---	---	---	---	---	---
DxB: Dunday-----	A		---	---	---	---	---	---	---
Et: Eltree-----	B		---	---	---	---	---	---	---
Go: Grigston-----	B		---	---	---	---	---	---	---
		April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
GrB: Aowa-----	B		---	---	---	---	---	Very brief	Frequent
		April	---	---	---	---	---	Very brief	Frequent
		May	---	---	---	---	---	Very brief	Frequent
		June	---	---	---	---	---	Very brief	Frequent
		July	---	---	---	---	---	Very brief	Frequent
		August	---	---	---	---	---	Very brief	Frequent
		September	---	---	---	---	---	Very brief	Frequent
Ha: Hall-----	B		---	---	---	---	---	---	---
He: Haynie-----	B		---	---	---	---	---	---	---
		March	---	---	---	---	---	Very brief	Occasional
		April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
IfD: Inavale-----	A		---	---	---	---	---	Very brief	Rare
		January	---	---	---	---	---	Very brief	Rare
		February	---	---	---	---	---	Very brief	Rare
		March	---	---	---	---	---	Very brief	Rare
		April	---	---	---	---	---	Very brief	Rare
		May	---	---	---	---	---	Very brief	Rare
		June	---	---	---	---	---	Very brief	Rare
		July	---	---	---	---	---	Very brief	Rare
IgB: Inavale-----	A		---	---	---	---	---	Very brief	Frequent
		January	---	---	---	---	---	Very brief	Frequent
		February	---	---	---	---	---	Very brief	Frequent
		March	---	---	---	---	---	Very brief	Frequent
		April	---	---	---	---	---	Very brief	Frequent
		May	---	---	---	---	---	Very brief	Frequent
		June	---	---	---	---	---	Very brief	Frequent
		July	---	---	---	---	---	Very brief	Frequent
IhB: Inavale-----	A		---	---	---	---	---	Very brief	Occasional
		January	---	---	---	---	---	Very brief	Occasional
		February	---	---	---	---	---	Very brief	Occasional
		March	---	---	---	---	---	Very brief	Occasional
		April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
In: Inavale-----	A		---	---	---	---	---	Very brief	Occasional
		January	---	---	---	---	---	Very brief	Occasional
		February	---	---	---	---	---	Very brief	Occasional
		March	---	---	---	---	---	Very brief	Occasional
		April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
Jn: Jansen-----	B		---	---	---	---	---	---	---
JnC: Jansen-----	B		---	---	---	---	---	---	---
JnD: Jansen-----	B		---	---	---	---	---	---	---
LaD: Labu-----	D		---	---	---	---	---	---	---

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
LcF: Labu-----	D		---	---	---	---	---	---	---
Sansarc-----	D		---	---	---	---	---	---	---
Le: Leshara-----	B								
		January	1.5-3.0	>6.0	---	---	---	---	None
		March	1.0-3.0	>6.0	---	---	---	Very brief	Occasional
		April	1.0-3.0	>6.0	---	---	---	Very brief	Occasional
		May	1.0-3.0	>6.0	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
LsC: Lynch-----	D		---	---	---	---	---	---	---
LsD: Lynch-----	D		---	---	---	---	---	---	---
LyD: Lynch-----	D		---	---	---	---	---	---	---
Bristow-----	D		---	---	---	---	---	---	---
LyF: Lynch-----	D		---	---	---	---	---	---	---
Bristow-----	D		---	---	---	---	---	---	---
MaG: Mariaville-----	D		---	---	---	---	---	---	---
Paka-----	B		---	---	---	---	---	---	---
MeE: Meadin-----	A		---	---	---	---	---	---	---
NoC: Nora-----	B		---	---	---	---	---	---	---
NoD: Nora-----	B		---	---	---	---	---	---	---
Oa: Onawa-----	D								
		January	2.0-4.0	>6.0	---	---	---	---	None
		February	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		March	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		April	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		May	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		June	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		July	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		September	---	---	---	---	---	Brief	Occasional
		October	---	---	---	---	---	Brief	Occasional
		November	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		December	2.0-4.0	>6.0	---	---	---	---	None
Oe: O'neill-----	B		---	---	---	---	---	---	---
OeC: O'neill-----	B		---	---	---	---	---	---	---
OfD: O'neill-----	B		---	---	---	---	---	---	---
Meadin-----	A		---	---	---	---	---	---	---
On: Onita-----	C		---	---	---	---	---	---	---
Poorly Drained-----	---		---	---	---	---	---	---	---
Or: Ord-----	B								
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		April	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		May	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
PaC: Paka-----	B								

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
Ph: Paka-----	B		---	---	---	---	---	---	---
PhC: Paka-----	B		---	---	---	---	---	---	---
PhD: Paka-----	B		---	---	---	---	---	---	---
PoC: Promise-----	D		---	---	---	---	---	---	---
RaC: Ree-----	B		---	---	---	---	---	---	---
RaD: Ree-----	B		---	---	---	---	---	---	---
RaE: Ree-----	B		---	---	---	---	---	---	---
ReC: Reliance-----	C		---	---	---	---	---	---	---
ReD: Reliance-----	C		---	---	---	---	---	---	---
RfC: Reliance-----	C		---	---	---	---	---	---	---
Rw: Riverwash-----	D		---	---	---	---	---	---	---
		January	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
		February	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
		March	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
		April	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
		May	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
		June	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
		July	---	---	---	---	---	Very long	Frequent
		October	---	---	---	---	---	Very long	Frequent
		November	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
		December	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
SAg: Sansarc-----	D		---	---	---	---	---	---	---
Sc: Scott-----	D		---	---	---	---	---	---	---
		March	0.0	>6.0	0.0-1.0	Long	---	---	None
		April	0.0	>6.0	0.0-1.0	Long	---	---	None
		May	0.0	>6.0	0.0-1.0	Long	---	---	None
		June	0.0	>6.0	0.0-1.0	Long	---	---	None
		July	0.0	>6.0	0.0-1.0	Long	---	---	None
Sm: Simeon-----	A		---	---	---	---	---	---	---
SuC: Simeon-----	A		---	---	---	---	---	---	---
Valentine-----	A		---	---	---	---	---	---	---
SvF2: Simeon-----	A		---	---	---	---	---	---	---
Valentine-----	A		---	---	---	---	---	---	---
VaE: Valentine-----	A		---	---	---	---	---	---	---
VbB: Valentine-----	A		---	---	---	---	---	---	---
Ve: Verdel-----	D		---	---	---	---	---	---	---
WeC: Wewela-----	B		---	---	---	---	---	---	---
zwa: Water-----	---		---	---	---	---	---	---	---

The following table gives estimates of various soil features. The estimates are used in land use planning that involves engineering considerations.

A restrictive layer is a nearly continuous layer that has one or more physical, chemical, or thermal properties that significantly impede the movement of water and air through the soil or that restrict roots or otherwise provide an unfavorable root environment. Examples are bedrock, cemented layers, dense layers, and frozen layers. The table indicates the hardness and thickness of the restrictive layer, both of which significantly affect the ease of excavation. Depth to top is the vertical distance from the soil surface to the upper boundary of the restrictive layer.

Potential for frost action is the likelihood of upward or lateral expansion of the soil caused by the formation of segregated ice lenses (frost heave) and the subsequent collapse of the soil and loss of strength on thawing. Frost action occurs when moisture moves into the freezing zone of the soil. Temperature, texture, density, permeability, content of organic matter, and depth to the water table are the most important factors considered in evaluating the potential for frost action. It is assumed that the soil is not insulated by vegetation or snow and is not artificially drained. Silty and highly structured, clayey soils that have a high water table in winter are the most susceptible to frost action. Well drained, very gravelly, or very sandy soils are the least susceptible. Frost heave and low soil strength during thawing cause damage to pavements and other rigid structures.

Risk of corrosion pertains to potential soil-induced electrochemical or chemical action that corrodes or weakens uncoated steel or concrete. The rate of corrosion of uncoated steel is related to such factors as soil moisture, particle-size distribution, acidity, and electrical conductivity of the soil. The rate of corrosion of concrete is based mainly on the sulfate and sodium content, texture, moisture content, and acidity of the soil. Special site examination and design may be needed if the combination of factors results in a severe hazard of corrosion. The steel or concrete in installations that intersect soil boundaries or soil layers is more susceptible to corrosion than the steel or concrete in installations that are entirely within one kind of soil or within one soil layer.

For uncoated steel, the risk of corrosion, expressed as low, moderate, or high, is based on soil drainage class, total acidity, electrical resistivity near field capacity, and electrical conductivity of the saturation extract.

For concrete, the risk of corrosion also is expressed as low, moderate, or high. It is based on soil texture, acidity, and amount of sulfates in the saturation extract.

Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top In	Thickness In	Hardness		Uncoated Steel	Concrete
Ab:							
Albaton-----	---	---	---	---	Moderate	High	Low
AnC:							
Anselmo-----	---	---	---	---	Moderate	Moderate	Low
AnD:							
Anselmo-----	---	---	---	---	Moderate	Moderate	Low
AnF:							
Anselmo-----	---	---	---	---	Moderate	Moderate	Low
ArF:							
Anselmo-----	---	---	---	---	Moderate	Moderate	Low
Rock Outcrop----	0-0	Bedrock (paralithic)	---	---	None	---	---
Ba:							
Barney-----	---	---	---	---	Moderate	High	Low
Bd:							
Blake-----	---	---	---	---	High	High	Low
Be:							
Blendon-----	---	---	---	---	Moderate	Moderate	Low
BeC:							
Blendon-----	---	---	---	---	Moderate	Moderate	Low
BoD:							
Boyd-----	20-40	Bedrock (paralithic)	---	---	Low	High	Moderate
BrG:							
Bristow-----	10-20	Bedrock (paralithic)	---	---	Moderate	High	Moderate
Bs:							
Brocksburg-----	---	---	---	---	Moderate	Low	Low
Bt:							
Brocksburg-----	---	---	---	---	Moderate	Low	Low
Cb:							
Cass-----	---	---	---	---	Moderate	Moderate	Low
CrE2:							
Crofton-----	---	---	---	---	Moderate	Low	Low
DuB:							
Dunday-----	---	---	---	---	Low	Low	Low
DuC:							
Dunday-----	---	---	---	---	Low	Low	Low
DuD:							
Dunday-----	---	---	---	---	Low	Low	Low
DxB:							
Dunday-----	---	---	---	---	Low	Low	Low
Et:							
Eltree-----	---	---	---	---	Moderate	Low	Low
Go:							
Grigston-----	---	---	---	---	Moderate	Low	Low
GrB:							
Aowa-----	---	---	---	---	High	Moderate	Low
Ha:							
Hall-----	---	---	---	---	Moderate	Moderate	Low
He:							
Haynie-----	---	---	---	---	High	Low	Low
IfD:							
Inavale-----	---	---	---	---	Low	Moderate	Low
IgB:							
Inavale-----	---	---	---	---	Low	Moderate	Low
IhB:							
Inavale-----	---	---	---	---	Low	Moderate	Low
In:							
Inavale-----	---	---	---	---	Low	Moderate	Low
Jn:							
Jansen-----	---	---	---	---	Moderate	Moderate	Low
JnC:							
Jansen-----	---	---	---	---	Moderate	Moderate	Low
JnD:							
Jansen-----	---	---	---	---	Moderate	Moderate	Low
LaD:							
Labu-----	20-40	Bedrock (paralithic)	---	---	Low	High	Moderate
LcF:							
Labu-----	20-40	Bedrock (paralithic)	---	---	Low	High	Moderate
Sansarc-----	4-20	Bedrock (paralithic)	---	---	Low	High	Moderate
Le:							
Leshara-----	---	---	---	---	High	High	Low
LsC:							
Lynch-----	20-40	Bedrock (paralithic)	---	---	Low	High	High
LsD:							
Lynch-----	20-40	Bedrock (paralithic)	---	---	Low	High	High
LyD:							
Lynch-----	20-40	Bedrock (paralithic)	---	---	Low	High	High
Bristow-----	10-20	Bedrock (paralithic)	---	---	Moderate	High	Moderate

Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top In	Thickness In	Hardness		Uncoated Steel	Concrete
LyF: Lynch-----	20-40	Bedrock (paralithic)	---	---	Low	High	High
Bristow-----	10-20	Bedrock (paralithic)	---	---	Moderate	High	Moderate
MaG: Mariaville-----	10-20	Bedrock (paralithic)	---	---	Moderate	Moderate	Low
Paka-----	40-60	Bedrock (paralithic)	---	---	Moderate	High	Low
MeE: Meadin-----	---	---	---	---	Low	Low	Moderate
NoC: Nora-----	---	---	---	---	High	Moderate	Low
NoD: Nora-----	---	---	---	---	High	Moderate	Low
Oa: Onawa-----	---	---	---	---	High	High	Low
Oe: O'neill-----	---	---	---	---	Moderate	Moderate	Low
OeC: O'neill-----	---	---	---	---	Moderate	Moderate	Low
OfD: O'neill-----	---	---	---	---	Moderate	Moderate	Low
Meadin-----	---	---	---	---	Low	Low	Moderate
On: Onita-----	---	---	---	---	Moderate	High	Low
Poorly Drained--	---	---	---	---	---	---	---
Or: Ord-----	---	---	---	---	High	High	Low
PaC: Paka-----	40-60	Bedrock (paralithic)	---	---	Moderate	High	Low
Ph: Paka-----	40-60	Bedrock (paralithic)	---	---	Moderate	High	Low
PhC: Paka-----	40-60	Bedrock (paralithic)	---	---	Moderate	High	Low
PhD: Paka-----	40-60	Bedrock (paralithic)	---	---	Moderate	High	Low
PoC: Promise-----	---	---	---	---	Low	High	Low
RaC: Ree-----	---	---	---	---	Moderate	High	Low
RaD: Ree-----	---	---	---	---	Moderate	High	Low
RaE: Ree-----	---	---	---	---	Moderate	High	Low
ReC: Reliance-----	---	---	---	---	Low	High	Low
ReD: Reliance-----	---	---	---	---	Low	High	Low
RfC: Reliance-----	---	---	---	---	Low	High	Low
Rw: Riverwash-----	---	---	---	---	Low	Moderate	Moderate
SaG: Sansarc-----	4-20	Bedrock (paralithic)	---	---	Low	High	Moderate
Sc: Scott-----	---	---	---	---	High	High	Low
Sm: Simeon-----	---	---	---	---	Low	Low	Low
SuC: Simeon-----	---	---	---	---	Low	Low	Low
Valentine-----	---	---	---	---	Low	Low	Low
SvF2: Simeon-----	---	---	---	---	Low	Low	Low
Valentine-----	---	---	---	---	Low	Low	Low
VaE: Valentine-----	---	---	---	---	Low	Low	Low
VbB: Valentine-----	---	---	---	---	Low	Low	Low
Ve: Verdel-----	---	---	---	---	Moderate	High	Low
WeC: Wewela-----	20-40	Bedrock (paralithic)	---	---	Moderate	High	Moderate
zwa: Water-----	---	---	---	---	---	---	---

WATER MANAGEMENT
Boyd County, Nebraska

The soils of the survey area are rated in the Water Management table according to limitations that affect their suitability for water management. Soils are rated for pond reservoir areas, drainage, irrigation, terraces and diversions, and grassed waterways. Restrictive features that affect each soil for the specified use is also provided in the table.

The ratings in the table are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the specified use. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Moderately limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Limited indicates that the soil has one or more features that are significant limitations for the specified use. The limitations can be overcome, but generally require special design, soil reclamation, or installation procedures that may result in additional expense. Fair performance and moderate to high maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Limitation class terms, such as very limited or limited, etc., limitation ratings, and numerical ratings are shown for each soil feature listed. As many as three soil features may be listed for each soil component if applicable. The overall limitation rating for the soil component is based on the most severe limitation.

Pond reservoir areas hold water behind a dam or embankment. Soils best suited to this use have low seepage potential in the upper 60 inches. The seepage potential is determined by the permeability of the soil and the depth to fractured bedrock or other permeable material. Excessive slope can affect the storage capacity of the reservoir area.

Embankments, dikes, and levees are raised structures of soil material, generally less than 20 feet high, constructed to impound water or to protect land against overflow. In this table, the soils are rated as a source of material for embankment fill. The ratings apply to the soil material below the surface layer to a depth of about 5 feet. It is assumed that soil layers will be uniformly mixed and compacted during construction.

The ratings do not indicate the ability of the natural soil to support an embankment. Soil properties to a depth even greater than the height of the embankment can affect performance and safety of the embankment. Generally, deeper onsite investigation is needed to determine these properties.

Soil material in embankments must be resistant to seepage, piping, and erosion and have favorable compaction characteristics. Unfavorable features include less than 5 feet of suitable material and a high content of stones or boulders, organic matter, or salts or sodium. A high water table affects the amount of usable material. It also affects traffic ability.

Aquifer-fed excavated ponds are pits or dugouts that extend to a ground-water aquifer or to a depth below a permanent water table. Excluded are ponds that are fed only by surface runoff and embankment ponds that impound water 3 feet or more above the original surface. Excavated ponds are affected by depth to a permanent water table, permeability of the aquifer, and quality of the water as inferred from the salinity of the soil. Depth to bedrock and the content of large stones affect the ease of excavation.

Drainage is the removal of excess surface and subsurface water from the soil. How easily and effectively the soil is drained depends on the depth to bedrock, to a cemented pan, or to other layers that affect the rate of water movement; permeability; depth to a high water table or depth of standing water if the soil is subject to ponding; slope; susceptibility to flooding; subsidence of organic layers; and the potential for frost action. Excavating and grading and the stability of ditch banks are affected by depth to bedrock or to a cemented pan, large stones, slope, and the hazard of cutbanks caving. The productivity of the soil after drainage is adversely affected by extreme acidity or by toxic substances in the root zone, such as salts, sodium, and sulfur. Availability of drainage outlets is not considered in the ratings.

Irrigation is the controlled application of water to supplement rainfall and support plant growth. The design and management of an irrigation system are affected by depth to the water table, the need for drainage, flooding, available water capacity, intake rate, permeability, erosion hazard, and slope. The construction of a system is affected by large stones and depth to bedrock or to a cemented pan. The performance of a system is affected by the depth of the root zone, the amount of salts or sodium, and soil reaction.

Terraces and diversions are embankments or a combination of channels and ridges constructed across a slope to control erosion and conserve moisture by intercepting runoff. Slope, wetness, large stones, and depth to bedrock or to a cemented pan affect the construction of terraces and diversions. A restricted rooting depth, a very limited hazard of wind erosion or water erosion, an excessively coarse texture, and restricted permeability adversely affect maintenance.

Grassed waterways are natural or constructed channels, generally broad and shallow, which conduct surface water to outlets at a non-erosive velocity. Large stones, wetness, slope, and depth to bedrock or to a cemented pan affect the construction of grassed waterways. A hazard of wind erosion, low available water capacity, restricted rooting depth, toxic substances such as salts and sodium, and restricted permeability adversely affect the growth and maintenance of the grass after construction.

WATER MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
Ab: Albaton-----	Limitation: flooding percs slowly	Limitation: percs slowly slow intake wetness	Limitation: percs slowly wetness	Limitation: percs slowly wetness
AnC: Anselmo-----	Limitation: deep to water	Limitation: slope	Limitation: soil blowing	Favorable
AnD: Anselmo-----	Limitation: deep to water	Limitation: slope	Limitation: slope soil blowing	Limitation: slope
AnF: Anselmo-----	Limitation: deep to water	Limitation: slope	Limitation: slope soil blowing	Limitation: slope
ArF: Anselmo-----	Limitation: deep to water	Limitation: slope	Limitation: slope soil blowing	Limitation: slope
Rock Outcrop---	Limitation: deep to water	Limitation: slope depth to rock	Limitation: slope depth to rock	Limitation: slope depth to rock
Ba: Barney-----	Limitation: flooding cutbanks cave	Limitation: rooting depth wetness droughty	Limitation: too sandy wetness	Limitation: rooting depth wetness droughty
Bd: Blake-----	Limitation: flooding frost action	Limitation: flooding wetness	Limitation: erodes easily wetness	Limitation: erodes easily
Be: Blendon-----	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable
BeC: Blendon-----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: soil blowing	Favorable
BoD: Boyd-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: area reclaim erodes easily slope	Limitation: erodes easily slope droughty
BrG: Bristow-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope droughty
Bs: Brocksburg-----	Limitation: deep to water	Limitation: soil blowing	Limitation: too sandy soil blowing	Favorable
Bt: Brocksburg-----	Limitation: deep to water	Favorable	Limitation: too sandy	Favorable
Cb: Cass-----	Limitation: deep to water	Limitation: flooding soil blowing	Limitation: soil blowing	Favorable
CrE2: Crofton-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily slope	Limitation: erodes easily slope
DuB: Dunday-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
DuC: Dunday-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
DuD: Dunday-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
DxB: Dunday-----	Limitation: deep to water	Limitation: fast intake soil blowing droughty	Limitation: too sandy soil blowing	Limitation: droughty

WATER MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
Et: Eltree-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
Go: Grigston-----	Limitation: deep to water	Limitation: flooding	Favorable	Favorable
GrB: Aowa-----	Limitation: deep to water	Limitation: flooding	Limitation: erodes easily	Limitation: erodes easily
Ha: Hall-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
He: Haynie-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
IfD: Inavale-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
IgB: Inavale-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
IhB: Inavale-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
In: Inavale-----	Limitation: deep to water	Limitation: flooding soil blowing droughty	Limitation: too sandy soil blowing	Limitation: droughty
Jn: Jansen-----	Limitation: deep to water	Favorable	Limitation: erodes easily too sandy	Limitation: erodes easily
JnC: Jansen-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily too sandy	Limitation: erodes easily
JnD: Jansen-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope too sandy	Limitation: erodes easily slope
LaD: Labu-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: percs slowly slope depth to rock	Limitation: slope depth to rock droughty
LcF: Labu-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: percs slowly slope depth to rock	Limitation: slope depth to rock droughty
Sansarc-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope droughty
Le: Leshara-----	Limitation: flooding frost action	Limitation: flooding wetness	Limitation: erodes easily wetness	Limitation: erodes easily wetness
LsC: Lynch-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: percs slowly depth to rock	Limitation: depth to rock droughty
LsD: Lynch-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: percs slowly slope depth to rock	Limitation: slope depth to rock droughty
LyD: Lynch-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: percs slowly slope depth to rock	Limitation: slope depth to rock droughty
Bristow-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope droughty

WATER MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
LyF: Lynch-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: percs slowly slope depth to rock	Limitation: slope depth to rock droughty
Bristow-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope droughty
MaG: Mariaville-----	Limitation: deep to water	Limitation: slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock
Paka-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
MeE: Meadin-----	Limitation: deep to water	Limitation: slope droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
NoC: Nora-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily
NoD: Nora-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
Oa: Onawa-----	Limitation: flooding frost action percs slowly	Limitation: percs slowly slow intake wetness	Limitation: erodes easily wetness	Limitation: erodes easily percs slowly
Oe: O'Neill-----	Limitation: deep to water	Limitation: droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
OeC: O'Neill-----	Limitation: deep to water	Limitation: slope droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
OfD: O'Neill-----	Limitation: deep to water	Limitation: slope droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
Meadin-----	Limitation: deep to water	Limitation: slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
On: Onita-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
Poorly Drained-- Or: Ord-----	Limitation: flooding frost action cutbanks cave	Limitation: flooding wetness soil blowing	Limitation: too sandy wetness soil blowing	Favorable
PaC: Paka-----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: erodes easily soil blowing	Limitation: erodes easily
Ph: Paka-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
PhC: Paka-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily
PhD: Paka-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
PoC: Promise-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: erodes easily percs slowly	Limitation: erodes easily droughty
RaC: Ree-----	Limitation: deep to water	Limitation: slope	Favorable	Favorable
RaD: Ree-----	Limitation: deep to water	Limitation: slope	Limitation: slope	Limitation: slope
RaE: Ree-----	Limitation: deep to water	Limitation: slope	Limitation: slope	Limitation: slope

WATER MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
ReC: Reliance-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily
ReD: Reliance-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
RfC: Reliance-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily
Rw: Riverwash-----	Limitation: flooding ponding cutbanks cave	Limitation: fast intake ponding droughty	Limitation: too sandy ponding	Limitation: rooting depth wetness droughty
SaG: Sansarc-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope droughty
Sc: Scott-----	Limitation: frost action percs slowly ponding	Limitation: erodes easily percs slowly ponding	Limitation: erodes easily percs slowly ponding	Limitation: erodes easily percs slowly wetness
Sm: Simeon-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
SuC: Simeon-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
SvF2: Simeon-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: rooting depth slope droughty
VaE: Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: rooting depth slope droughty
VbB: Valentine-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
Ve: Verdel-----	Limitation: deep to water	Limitation: percs slowly slow intake	Limitation: percs slowly	Limitation: percs slowly
WeC: Wewela-----	Limitation: deep to water	Limitation: percs slowly slope thin layer	Limitation: area reclaim soil blowing	Limitation: area reclaim percs slowly
zwa: Water-----	---	---	---	---

WATER MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Not limited		Very limited Depth to saturated zone Hard to pack	1.00 1.00	Very limited Slow refill Cutbanks cave	1.00 0.10
AnC: Anselmo-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
AnD: Anselmo-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
AnF: Anselmo-----	100	Very limited Seepage Slope	1.00 0.02	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
ArF: Anselmo-----	75	Very limited Seepage Slope	1.00 0.02	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
Ba: Barney-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.95	Very limited Cutbanks cave	1.00
Bd: Blake-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping Depth to saturated zone Seepage	0.81 0.43 0.05	Somewhat limited Slow refill Deep to water Cutbanks cave	0.30 0.25 0.10
Be: Blendon-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
BeC: Blendon-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
BoD: Boyd-----	100	Somewhat limited Depth to bedrock	0.11	Very limited Hard to pack Thin layer	1.00 0.85	Very limited Deep to water	1.00
BrG: Bristow-----	100	Somewhat limited Depth to bedrock Slope	0.66 0.50	Very limited Thin layer Hard to pack	1.00 1.00	Very limited Deep to water	1.00
Bs: Brocksburg-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Bt: Brocksburg-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Cb: Cass-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.95	Very limited Deep to water	1.00
CrE2: Crofton-----	100	Somewhat limited Seepage Slope	0.70 0.01	Somewhat limited Piping	0.50	Very limited Deep to water	1.00
DuB: Dunday-----	100	Very limited		Very limited		Very limited	

WATER MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
		Seepage	1.00	Seepage	1.00	Deep to water	1.00
DuC: Dunday-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
DuD: Dunday-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
DxB: Dunday-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.91	Very limited Deep to water	1.00
Et: Eltree-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.89	Very limited Deep to water	1.00
Go: Grigston-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.55	Very limited Deep to water	1.00
GrB: Aowa-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.95	Very limited Deep to water	1.00
Ha: Hall-----	100	Very limited Seepage	1.00	Somewhat limited Piping	0.11	Very limited Deep to water	1.00
He: Haynie-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
IfD: Inavale-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
IgB: Inavale-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
IhB: Inavale-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.98	Very limited Deep to water	1.00
In: Inavale-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.98	Very limited Deep to water	1.00
Jn: Jansen-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.97	Very limited Deep to water	1.00
JnC: Jansen-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.97	Very limited Deep to water	1.00
JnD: Jansen-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.97	Very limited Deep to water	1.00
LaD: Labu-----	100	Somewhat limited Depth to bedrock	0.11	Somewhat limited Hard to pack Thin layer	0.97 0.85	Very limited Deep to water	1.00
LcF: Labu-----	65	Somewhat limited Slope Depth to bedrock	0.15 0.11	Somewhat limited Hard to pack Thin layer	0.97 0.85	Very limited Deep to water	1.00
Sansarc-----	35	Very limited Seepage Depth to bedrock Slope	1.00 0.76 0.15	Very limited Thin layer Hard to pack	1.00 1.00	Very limited Deep to water	1.00
Le: Leshara-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Cutbanks cave	1.00

WATER MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
LsC: Lynch-----	100	Somewhat limited Depth to bedrock	0.11	Depth to saturated zone	1.00	Slow refill	0.30
						Deep to water	0.00
				Somewhat limited Thin layer Hard to pack	0.85 0.57	Very limited Deep to water	1.00
LsD: Lynch-----	100	Somewhat limited Depth to bedrock	0.11	Somewhat limited Thin layer Hard to pack	0.85 0.57	Very limited Deep to water	1.00
LyD: Lynch-----	65	Somewhat limited Depth to bedrock	0.11	Somewhat limited Thin layer Hard to pack	0.85 0.57	Very limited Deep to water	1.00
Bristow-----	35	Somewhat limited Depth to bedrock	0.66	Very limited Thin layer Hard to pack	1.00 1.00	Very limited Deep to water	1.00
LyF: Lynch-----	55	Somewhat limited Slope Depth to bedrock	0.15 0.11	Somewhat limited Thin layer Hard to pack	0.85 0.57	Very limited Deep to water	1.00
Bristow-----	45	Somewhat limited Depth to bedrock Slope	0.66 0.15	Very limited Thin layer Hard to pack	1.00 1.00	Very limited Deep to water	1.00
MaG: Mariaville-----	55	Somewhat limited Depth to bedrock Slope	0.66 0.40	Very limited Thin layer Piping	1.00 0.50	Very limited Deep to water	1.00
Paka-----	45	Somewhat limited Seepage Slope Depth to bedrock	0.70 0.40 0.00	Somewhat limited Piping Thin layer	0.45 0.11	Very limited Deep to water	1.00
MeE: Meadin-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.91	Very limited Deep to water	1.00
NoC: Nora-----	100	Somewhat limited Seepage	0.55	Somewhat limited Piping	0.65	Very limited Deep to water	1.00
NoD: Nora-----	100	Somewhat limited Seepage	0.55	Somewhat limited Piping	0.65	Very limited Deep to water	1.00
Oa: Onawa-----	100	Very limited Seepage	1.00	Somewhat limited Depth to saturated zone	0.43	Somewhat limited Deep to water	0.25
						Cutbanks cave	0.10
Oe: O'Neill-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
OeC: O'Neill-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
OfD: O'Neill-----	55	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Meadin-----	45	Very limited Seepage	1.00	Somewhat limited Seepage	0.91	Very limited Deep to water	1.00
On: Onita-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.03	Very limited Deep to water	1.00
Poorly Drained-----		Very limited		Very limited		Very limited	

WATER MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Or: Ord-----	100	Seepage Slope	1.00 0.50	Hard to pack	1.00	Deep to water	1.00
		Very limited Seepage	1.00	Somewhat limited Seepage Depth to saturated zone	0.97 0.95	Very limited Cutbanks cave Deep to water	1.00 0.02
PaC: Paka-----	100	Somewhat limited Seepage Depth to bedrock	0.70 0.00	Somewhat limited Piping Thin layer	0.58 0.11	Very limited Deep to water	1.00
Ph: Paka-----	100	Somewhat limited Seepage Depth to bedrock	0.70 0.00	Somewhat limited Piping Thin layer	0.45 0.11	Very limited Deep to water	1.00
PhC: Paka-----	100	Somewhat limited Seepage Depth to bedrock	0.70 0.00	Somewhat limited Piping Thin layer	0.45 0.11	Very limited Deep to water	1.00
PhD: Paka-----	100	Somewhat limited Seepage Depth to bedrock	0.70 0.00	Somewhat limited Piping Thin layer	0.45 0.11	Very limited Deep to water	1.00
PoC: Promise-----	100	Not limited		Somewhat limited Hard to pack	1.00	Very limited Deep to water	1.00
RaC: Ree-----	100	Very limited Seepage	1.00	Somewhat limited Piping Seepage	0.70 0.09	Very limited Deep to water	1.00
RaD: Ree-----	100	Very limited Seepage	1.00	Somewhat limited Piping Seepage	0.70 0.09	Very limited Deep to water	1.00
RaE: Ree-----	100	Very limited Seepage Slope	1.00 0.01	Somewhat limited Piping Seepage	0.70 0.04	Very limited Deep to water	1.00
ReC: Reliance-----	100	Somewhat limited Seepage	0.57	Not limited		Very limited Deep to water	1.00
ReD: Reliance-----	100	Somewhat limited Seepage	0.57	Not limited		Very limited Deep to water	1.00
RfC: Reliance-----	100	Somewhat limited Seepage	0.57	Not limited		Very limited Deep to water	1.00
Rw: Riverwash-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Cutbanks cave	1.00
SaG: Sansarc-----	100	Very limited Seepage Depth to bedrock Slope	1.00 0.76 0.72	Very limited Thin layer Hard to pack	1.00 1.00	Very limited Deep to water	1.00
Sc: Scott-----	100	Somewhat limited Seepage	0.04	Very limited Depth to saturated zone Hard to pack	1.00 0.46	Somewhat limited Slow refill Cutbanks cave	0.30 0.10
Sm: Simeon-----	100	Very limited		Somewhat limited		Very limited	

WATER MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
SuC: Simeon-----	55	Seepage	1.00	Seepage	0.98	Deep to water	1.00
		Very limited Seepage	1.00	Somewhat limited Seepage	0.98	Very limited Deep to water	1.00
		Valentine-----	45	Very limited Seepage	1.00	Very limited Deep to water	1.00
SvF2: Simeon-----	70	Very limited Seepage Slope	1.00 0.06	Somewhat limited Seepage	0.98	Very limited Deep to water	1.00
		Valentine-----	30	Very limited Seepage Slope	1.00 0.06	Very limited Deep to water	1.00
		VaE: Valentine-----	100	Very limited Seepage Slope	1.00 0.00	Very limited Deep to water	1.00
VbB: Valentine-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Ve: Verdel-----	100	Not limited		Somewhat limited Hard to pack	0.92	Very limited Deep to water	1.00
WeC: Wewela-----	100	Somewhat limited Depth to bedrock	0.11	Somewhat limited Thin layer Seepage	0.85 0.09	Very limited Deep to water	1.00
zwa: Water-----	100	Not rated		Not rated		Not rated	

SANITARY FACILITIES
Boyd County, Nebraska

Sanitary Facilities

The following tables show the degree and kind of soil limitations that affect septic tank absorption fields, sewage lagoons, sanitary landfills, and daily cover for landfill. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect these uses. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Septic tank absorption fields are areas in which effluent from a septic tank is distributed into the soil through subsurface tiles or perforated pipe. Only that part of the soil between depths of 24 and 60 inches is evaluated. The ratings are based on the soil properties that affect absorption of the effluent, construction and maintenance of the system, and public health. Permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, and flooding affect absorption of the effluent. Stones and boulders, ice, and bedrock or a cemented pan interfere with installation. Subsidence interferes with installation and maintenance. Excessive slope may cause lateral seepage and surfacing of the effluent in downslope areas.

Some soils are underlain by loose sand and gravel or fractured bedrock at a depth of less than 4 feet below the distribution lines. In these soils the absorption field may not adequately filter the effluent, particularly when the system is new. As a result, the ground water may become contaminated.

Sewage lagoons are shallow ponds constructed to hold sewage while aerobic bacteria decompose the solid and liquid wastes. Lagoons should have a nearly level floor surrounded by cut slopes or embankments of compacted soil. Nearly impervious soil material for the lagoon floor and sides is required to minimize seepage and contamination of ground water. Considered in the ratings are slope, permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, flooding, large stones, and content of organic matter.

Soil permeability is a critical property affecting the suitability for sewage lagoons. Most porous soils eventually become sealed when they are used as sites for sewage lagoons. Until sealing occurs, however, the hazard of pollution is severe. Soils that have a permeability rate of more than 2 inches per hour are too porous for the proper functioning of sewage lagoons. In these soils, seepage of the effluent can result in contamination of the ground water. Ground-water contamination is also a hazard if fractured bedrock is within a depth of 40 inches, if the water table is high enough to raise the level of sewage in the lagoon, or if floodwater overtops the lagoon.

A high content of organic matter is detrimental to proper functioning of the lagoon because it inhibits aerobic activity. Slope, bedrock, and cemented pans can cause construction problems, and large stones can hinder compaction of the lagoon floor. If the lagoon is to be uniformly deep throughout, the slope must be gentle enough and the soil material must be thick enough over bedrock or a cemented pan to make land smoothing practical.

A trench sanitary landfill is an area where solid waste is placed in successive layers in an excavated trench. The waste is spread, compacted, and covered daily with a thin layer of soil excavated at the site. When the trench is full, a final cover of soil material at least 2 feet thick is placed over the landfill. The ratings in the table are based on the soil properties that affect the risk of pollution, the ease of excavation, trafficability, and revegetation. These properties include permeability, depth to bedrock or a cemented pan, depth to a water table, ponding, slope, flooding, texture, stones and boulders, highly organic layers, soil reaction, and content of salts and sodium. Unless otherwise stated, the ratings apply only to that part of the soil within a depth of about 6 feet. For deeper trenches, onsite investigation may be needed.

Hard, nonrippable bedrock, creviced bedrock, or highly permeable strata in or directly below the proposed trench bottom can affect the ease of excavation and the hazard of ground-water pollution. Slope affects construction of the trenches and the movement of surface water around the landfill. It also affects the construction and performance of roads in areas of the landfill.

Soil texture and consistence affect the ease with which the trench is dug and the ease with which the soil can be used as daily or final cover. They determine the workability of the soil when dry and when wet. Soils that are plastic and sticky when wet are difficult to excavate, grade, or compact and are difficult to place as a uniformly thick cover over a layer of refuse.

The soil material used as the final cover for a trench landfill should be suitable for plants. It should not have excess sodium or salts and should not be too acid. The surface layer generally has the best workability, the highest content of organic matter, and the best potential for plants. Material from the surface layer should be stockpiled for use as the final cover.

SANITARY FACILITIES
Boyd County, Nebraska

In an area sanitary landfill, solid waste is placed in successive layers on the surface of the soil. The waste is spread, compacted, and covered daily with a thin layer of soil from a source away from the site. A final cover of soil material at least 2 feet thick is placed over the completed landfill. The ratings in the table are based on the soil properties that affect trafficability and the risk of pollution. These properties include flooding, permeability, depth to a water table, ponding, slope, and depth to bedrock or a cemented pan.

Flooding is a serious problem because it can result in pollution in areas downstream from the landfill. If permeability is too rapid or if fractured bedrock, a fractured cemented pan, or the water table is close to the surface, the leachate can contaminate the water supply. Slope is a consideration because of the extra grading required to maintain roads in the steeper areas of the landfill. Also, leachate may flow along the surface of the soils in the steeper areas and cause difficult seepage problems.

Daily cover for landfill is the soil material that is used to cover compacted solid waste in an area sanitary landfill. The soil material is obtained offsite, transported to the landfill, and spread over the waste. The ratings in the table also apply to the final cover for a landfill. They are based on the soil properties that affect workability, the ease of digging, and the ease of moving and spreading the material over the refuse daily during wet and dry periods. These properties include soil texture, depth to a water table, ponding, rock fragments, slope, depth to bedrock or a cemented pan, reaction, and content of salts, sodium, or lime.

Loamy or silty soils that are free of large stones and excess gravel are the best cover for a landfill. Clayey soils may be sticky and difficult to spread; sandy soils are subject to wind erosion.

Slope affects the ease of excavation and of moving the cover material. Also, it can influence runoff, erosion, and reclamation of the borrow area.

After soil material has been removed, the soil material remaining in the borrow area must be thick enough over bedrock, a cemented pan, or the water table to permit revegetation. The soil material used as the final cover for a landfill should be suitable for plants. It should not have excess sodium, salts, or lime and should not be too acid.

SANITARY FACILITIES--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Very limited Flooding Restricted permeability Depth to saturated zone	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
AnC: Anselmo-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.33
AnD: Anselmo-----	100	Very limited Filtering capacity Slope	1.00 0.04	Very limited Seepage Slope	1.00 1.00
AnF: Anselmo-----	100	Very limited Filtering capacity Slope	1.00 0.96	Very limited Slope Seepage	1.00 1.00
ArF: Anselmo-----	75	Very limited Filtering capacity Slope	1.00 0.96	Very limited Slope Seepage	1.00 1.00
Rock Outcrop-----	25	Not rated		Not rated	
Ba: Barney-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
Bd: Blake-----	100	Very limited Flooding Depth to saturated zone Restricted permeability	1.00 1.00 0.71	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 0.50
Be: Blendon-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
BeC: Blendon-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.33
BoD: Boyd-----	100	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to soft bedrock Slope	1.00 1.00
BrG: Bristow-----	100	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 1.00
Bs: Brocksburg-----	100	Very limited Filtering capacity Restricted permeability	1.00 1.00	Very limited Seepage	1.00
Bt: Brocksburg-----	100	Very limited Filtering capacity Restricted permeability	1.00 1.00	Very limited Seepage	1.00
Cb: Cass-----	100	Very limited Flooding Filtering capacity	1.00 1.00	Very limited Flooding Seepage	1.00 1.00

SANITARY FACILITIES--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
CrE2: Crofton-----	100	Somewhat limited Slope Restricted permeability	0.84 0.50	Very limited Slope Seepage	1.00 0.50
DuB: Dunday-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.00
DuC: Dunday-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.67
DuD: Dunday-----	100	Very limited Filtering capacity Slope	1.00 0.04	Very limited Seepage Slope	1.00 1.00
DxB: Dunday-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.00
Et: Eltree-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
Go: Grigston-----	100	Very limited Flooding Restricted permeability	1.00 0.50	Very limited Flooding Seepage	1.00 0.50
GrB: Aowa-----	100	Very limited Flooding Restricted permeability	1.00 0.50	Very limited Flooding Seepage	1.00 0.50
Ha: Hall-----	100	Very limited Restricted permeability	1.00	Very limited Seepage	1.00
He: Haynie-----	100	Very limited Flooding Restricted permeability	1.00 0.50	Very limited Flooding Seepage	1.00 0.50
IfD: Inavale-----	100	Very limited Filtering capacity Flooding	1.00 0.40	Very limited Seepage Slope Flooding	1.00 1.00 0.40
IgB: Inavale-----	100	Very limited Flooding Filtering capacity	1.00 1.00	Very limited Flooding Seepage Slope	1.00 1.00 0.00
IhB: Inavale-----	100	Very limited Flooding Filtering capacity	1.00 1.00	Very limited Flooding Seepage Slope	1.00 1.00 0.00
In: Inavale-----	100	Very limited Flooding Filtering capacity	1.00 1.00	Very limited Flooding Seepage Slope	1.00 1.00 0.00
Jn: Jansen-----	100	Very limited Filtering capacity Restricted permeability	1.00 1.00	Very limited Seepage	1.00
JnC: Jansen-----	100	Very limited		Very limited	

SANITARY FACILITIES--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
JnD: Jansen-----	100	Filtering capacity	1.00	Seepage	1.00
		Restricted permeability	1.00	Slope	0.33
		Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Restricted permeability	1.00	Slope	1.00
LaD: Labu-----	100	Slope	0.04		
		Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
LcF: Labu-----	65	Slope	0.04	Slope	1.00
		Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
Sansarc-----	35	Slope	1.00	Slope	1.00
		Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
		Slope	1.00	Slope	1.00
Le: Leshara-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Restricted permeability	0.50	Seepage	0.50
LsC: Lynch-----	100	Very limited Restricted permeability	1.00	Very limited Depth to soft bedrock	1.00
		Depth to bedrock	1.00	Slope	0.33
LsD: Lynch-----	100	Very limited Restricted permeability	1.00	Very limited Depth to soft bedrock	1.00
		Depth to bedrock	1.00	Slope	1.00
LyD: Lynch-----	65	Slope	0.04		
		Very limited Restricted permeability	1.00	Very limited Depth to soft bedrock	1.00
		Depth to bedrock	1.00	Slope	1.00
		Slope	0.04		
Bristow-----	35	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
		Slope	0.04	Slope	1.00
LyF: Lynch-----	55	Very limited Restricted permeability	1.00	Very limited Depth to soft bedrock	1.00
		Depth to bedrock	1.00	Slope	1.00
		Slope	1.00		
Bristow-----	45	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
		Slope	1.00	Slope	1.00
MaG: Mariaville-----	55	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
		Slope	1.00	Slope	1.00
Paka-----	45	Very limited Slope	1.00	Very limited Slope	1.00
		Restricted permeability	1.00	Seepage	0.50
		Depth to bedrock	0.78	Depth to soft bedrock	0.42
MeE: Meadin-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Slope	0.16	Slope	1.00

SANITARY FACILITIES--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
NoC: Nora-----	100	Somewhat limited Restricted permeability	0.71	Somewhat limited Slope	0.33
				Seepage	0.29
NoD: Nora-----	100	Somewhat limited Restricted permeability Slope	0.71	Very limited Slope	1.00
			0.04	Seepage	0.29
Oa: Onawa-----	100	Very limited Flooding Restricted permeability Depth to saturated zone	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00
Oe: O'Neill-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
OeC: O'Neill-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
				Slope	0.33
OfD: O'Neill-----	55	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
				Slope	0.91
Meadin-----	45	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
				Slope	0.91
On: Onita-----	100	Very limited Restricted permeability	1.00	Somewhat limited Seepage	0.50
Poorly Drained-----		Very limited Slope	1.00	Very limited Slope	1.00
Or: Ord-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
PaC: Paka-----	100	Very limited Restricted permeability Depth to bedrock	1.00 0.78	Somewhat limited Seepage Depth to soft bedrock Slope	0.50 0.42 0.33
Ph: Paka-----	100	Very limited Restricted permeability Depth to bedrock	1.00 0.78	Somewhat limited Seepage Depth to soft bedrock	0.50 0.42
PhC: Paka-----	100	Very limited Restricted permeability Depth to bedrock	1.00 0.78	Somewhat limited Seepage Depth to soft bedrock Slope	0.50 0.42 0.33
PhD: Paka-----	100	Very limited Restricted permeability Depth to bedrock Slope	1.00 0.78 0.04	Very limited Slope Seepage Depth to soft bedrock	1.00 0.50 0.42
PoC: Promise-----	100	Very limited Restricted permeability	1.00	Somewhat limited Slope	0.33

SANITARY FACILITIES--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
RaC: Ree-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.33
RaD: Ree-----	100	Somewhat limited Slope	0.04	Very limited Slope Seepage	1.00 1.00
RaE: Ree-----	100	Somewhat limited Slope	0.84	Very limited Slope Seepage	1.00 1.00
ReC: Reliance-----	100	Very limited Restricted permeability	1.00	Somewhat limited Slope Seepage	0.33 0.32
ReD: Reliance-----	100	Very limited Restricted permeability Slope	1.00 0.04	Very limited Slope Seepage	1.00 0.32
RfC: Reliance-----	100	Very limited Restricted permeability	1.00	Somewhat limited Slope Seepage	0.33 0.32
Rw: Riverwash-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
SaG: Sansarc-----	100	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 1.00
Sc: Scott-----	100	Very limited Restricted permeability Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone	1.00
Sm: Simeon-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
SuC: Simeon-----	55	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.09
Valentine-----	45	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.09
SvF2: Simeon-----	70	Very limited Filtering capacity Slope	1.00 1.00	Very limited Seepage Slope	1.00 1.00
Valentine-----	30	Very limited Filtering capacity Slope	1.00 1.00	Very limited Seepage Slope	1.00 1.00
VaE: Valentine-----	100	Very limited Filtering capacity Slope	1.00 0.63	Very limited Seepage Slope	1.00 1.00
VbB: Valentine-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.00
Ve: Verdel-----	100	Very limited		Not limited	

SANITARY FACILITIES--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
WeC: Wewela-----	100	Restricted permeability	1.00		
		Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock Slope	1.00 0.33
zwa: Water-----	100	Not rated		Not rated	

SANITARY FACILITIES--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Too clayey	1.00
		Too clayey	1.00			Hard to compact	1.00
AnC: Anselmo-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Seepage	1.00
		Too Sandy	1.00			Too Sandy	0.50
AnD: Anselmo-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Seepage	1.00
		Too Sandy	1.00	Slope	0.04	Too Sandy	0.50
		Slope	0.04			Slope	0.04
AnF: Anselmo-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Seepage	1.00
		Too Sandy	1.00	Slope	0.96	Slope	0.96
		Slope	0.96			Too Sandy	0.50
ArF: Anselmo-----	75	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Seepage	1.00
		Too Sandy	1.00	Slope	0.96	Slope	0.96
		Slope	0.96			Too Sandy	0.50
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
Ba: Barney-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Too Sandy	1.00
		Seepage	1.00	Seepage	1.00	Seepage	1.00
		Too Sandy	1.00				
Bd: Blake-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.09
		Depth to saturated zone	1.00	Depth to saturated zone	1.00		
Be: Blendon-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Somewhat limited Seepage	0.50
BeC: Blendon-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Somewhat limited Seepage	0.50
BoD: Boyd-----	100	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Too clayey	1.00	Slope	0.04	Too clayey	1.00
		Slope	0.04			Hard to compact	1.00
						Slope	0.04
BrG: Bristow-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Depth to bedrock	1.00
		Depth to bedrock	1.00	Depth to bedrock	1.00	Slope	1.00
		Too clayey	1.00			Too clayey	1.00
						Hard to compact	1.00
Bs: Brocksburg-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
Bt: Brocksburg-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
Cb: Cass-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Seepage	1.00
		Seepage	1.00	Seepage	1.00		
CrE2: Crofton-----	100	Somewhat limited Slope	0.84	Somewhat limited Slope	0.84	Somewhat limited Slope	0.84
DuB: Dunday-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00

SANITARY FACILITIES--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
DuC: Dunday-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage	1.00 1.00
DuD: Dunday-----	100	Very limited Seepage Too Sandy Slope	1.00 1.00 0.04	Very limited Seepage Slope	1.00 0.04	Very limited Too Sandy Seepage Slope	1.00 1.00 0.04
DxB: Dunday-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Seepage Too Sandy	1.00 0.50
Et: Eltree-----	100	Not limited		Not limited		Not limited	
Go: Grigston-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Not limited	
GrB: Aowa-----	100	Very limited Flooding Too Sandy	1.00 1.00	Very limited Flooding	1.00	Very limited Too Sandy	1.00
Ha: Hall-----	100	Very limited Seepage Too clayey	1.00 0.50	Not limited		Somewhat limited Too clayey	0.50
He: Haynie-----	100	Very limited Flooding Too Sandy	1.00 1.00	Very limited Flooding	1.00	Very limited Too Sandy	1.00
IfD: Inavale-----	100	Very limited Seepage Too Sandy Flooding	1.00 1.00 0.40	Very limited Seepage Flooding	1.00 0.40	Very limited Too Sandy Seepage	1.00 1.00
IgB: Inavale-----	100	Very limited Flooding Seepage Too Sandy	1.00 1.00 1.00	Very limited Flooding Seepage	1.00 1.00	Very limited Too Sandy Seepage	1.00 1.00
ThB: Inavale-----	100	Very limited Flooding Seepage Too Sandy	1.00 1.00 1.00	Very limited Flooding Seepage	1.00 1.00	Very limited Seepage Too Sandy	1.00 0.50
In: Inavale-----	100	Very limited Flooding Seepage Too Sandy	1.00 1.00 1.00	Very limited Flooding Seepage	1.00 1.00	Very limited Too Sandy Seepage	1.00 1.00
Jn: Jansen-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage	1.00 1.00
JnC: Jansen-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage	1.00 1.00
JnD: Jansen-----	100	Very limited Seepage Too Sandy Slope	1.00 1.00 0.04	Very limited Seepage Slope	1.00 0.04	Very limited Too Sandy Seepage Slope	1.00 1.00 0.04
LaD: Labu-----	100	Very limited Depth to bedrock Too clayey Slope	1.00 1.00 0.04	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to bedrock Too clayey Hard to compact Slope	1.00 1.00 1.00 0.04
LcF: Labu-----	65	Very limited Depth to bedrock Too clayey Slope	1.00 1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Too clayey Hard to compact Slope	1.00 1.00 1.00 1.00
Sansarc-----	35	Very limited Depth to bedrock Too clayey Slope Seepage	1.00 1.00 1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Too clayey Hard to compact Slope	1.00 1.00 1.00 1.00
Le: Leshara-----	100	Very limited		Very limited		Very limited	

SANITARY FACILITIES--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
LsC: Lynch-----	100	Flooding	1.00	Flooding	1.00	Too Sandy	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	0.86
		Too Sandy	1.00				
LsD: Lynch-----	100	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Too clayey	1.00			Too clayey	1.00
						Hard to compact	1.00
LyD: Lynch-----	100	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Too clayey	1.00	Depth to bedrock	1.00	Too clayey	1.00
		Slope	0.04	Slope	0.04	Hard to compact	1.00
LyD: Lynch-----	65	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Too clayey	1.00	Slope	0.04	Too clayey	1.00
		Slope	0.04			Hard to compact	1.00
Bristow-----	35	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Too clayey	1.00	Slope	0.04	Too clayey	1.00
		Slope	0.04			Hard to compact	1.00
LyF: Lynch-----	55	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Too clayey	1.00	Slope	1.00	Too clayey	1.00
		Slope	1.00			Hard to compact	1.00
Bristow-----	45	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Too clayey	1.00	Slope	1.00	Too clayey	1.00
		Slope	1.00			Slope	1.00
MaG: Mariaville-----	55	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Slope	1.00	Slope	1.00	Slope	1.00
		Too clayey	0.50			Too clayey	0.50
Paka-----	45	Very limited Depth to bedrock	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Slope	1.00	Depth to bedrock	0.42	Depth to bedrock	0.42
MeE: Meadin-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00	Slope	0.16	Seepage	1.00
		Slope	0.16			Gravel content	0.87
NoC: Nora-----	100	Not limited		Not limited		Not limited	
NoD: Nora-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04
Oa: Onawa-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Somewhat limited Seepage	0.21
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	0.09
		Seepage	1.00	Seepage	1.00		
Oe: O'Neill-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
OeC: O'Neill-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
OfD: O'Neill-----	55	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
Meadin-----	45	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
						Gravel content	0.87
On: Onita-----	100	Not limited		Not limited		Very limited Hard to compact	1.00
Poorly Drained-----		Very limited		Very limited		Very limited	

SANITARY FACILITIES--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Or:		Slope	1.00	Slope	1.00	Slope	1.00
Ord-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Too Sandy	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Seepage	1.00
		Seepage	1.00	Seepage	1.00	Depth to saturated zone	0.68
		Too Sandy	1.00				
PaC:							
Paka-----	100	Very limited Depth to bedrock	1.00	Somewhat limited Depth to bedrock	0.42	Somewhat limited Depth to bedrock	0.42
Ph:							
Paka-----	100	Very limited Depth to bedrock	1.00	Somewhat limited Depth to bedrock	0.42	Somewhat limited Depth to bedrock	0.42
PhC:							
Paka-----	100	Very limited Depth to bedrock	1.00	Somewhat limited Depth to bedrock	0.42	Somewhat limited Depth to bedrock	0.42
PhD:							
Paka-----	100	Very limited Depth to bedrock	1.00	Somewhat limited Depth to bedrock	0.42	Somewhat limited Depth to bedrock	0.42
		Slope	1.00	Slope	0.42	Slope	0.42
			0.04		0.04		0.04
PoC:							
Promise-----	100	Very limited Too clayey	1.00	Not limited		Very limited Too clayey	1.00
						Hard to compact	1.00
RaC:							
Ree-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Seepage	1.00
RaD:							
Ree-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Somewhat limited Seepage	0.50
		Slope	0.04	Slope	0.04	Slope	0.04
RaE:							
Ree-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Somewhat limited Slope	0.84
		Slope	0.84	Slope	0.84	Seepage	0.50
ReC:							
Reliance-----	100	Not limited		Not limited		Very limited Hard to compact	1.00
ReD:							
Reliance-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Hard to compact	1.00
						Slope	0.04
RfC:							
Reliance-----	100	Somewhat limited Too clayey	0.50	Not limited		Very limited Hard to compact	1.00
						Too clayey	0.50
Rw:							
Riverwash-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Too Sandy	1.00
		Seepage	1.00	Seepage	1.00	Seepage	1.00
		Too Sandy	1.00			Gravel content	0.30
SaG:							
Sansarc-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Depth to bedrock	1.00
		Depth to bedrock	1.00	Depth to bedrock	1.00	Slope	1.00
		Too clayey	1.00			Too clayey	1.00
		Seepage	1.00			Hard to compact	1.00
Sc:							
Scott-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Too clayey	1.00			Too clayey	1.00
						Hard to compact	1.00
Sm:							
Simeon-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
SuC:							
Simeon-----	55	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
Valentine-----	45	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
SvF2:							
Simeon-----	70	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00

SANITARY FACILITIES--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Valentine-----	30	Too Sandy Slope Very limited Seepage Too Sandy Slope	1.00 1.00 1.00 1.00 1.00	Slope Very limited Seepage Slope	1.00 1.00 1.00	Seepage Slope Very limited Too Sandy Seepage Slope	1.00 1.00 1.00 1.00 1.00
VaE: Valentine-----	100	Very limited Seepage Too Sandy Slope	1.00 1.00 0.63	Very limited Seepage Slope	1.00 0.63	Very limited Too Sandy Seepage Slope	1.00 1.00 0.63
VbB: Valentine-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage	1.00 1.00
Ve: Verdel-----	100	Very limited Too clayey	1.00	Not limited		Very limited Too clayey Hard to compact	1.00 1.00
WeC: Wewela-----	100	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Hard to compact	1.00 1.00
zwa: Water-----	100	Not rated		Not rated		Not rated	

AGRICULTURAL WASTE MANAGEMENT
Boyd County, Nebraska

The nature of the soil is also important in the application of organic wastes and wastewater to land as fertilizers and irrigation; it is also important when the soil is used as a medium for treatment and disposal of these wastes. Favorable soil properties are required to prevent environmental damage.

The use of organic wastes and wastewater as production resources will result in energy conservation, prevent the waste of these important resources, and prevent problems associated with their disposal. Where disposal is the goal, and a maximum amount is disposed in a minimum area to hold costs to a minimum, risk of environmental damage is the principal constraint. Where the reuse goal is pursued, and a minimum amount is applied to a maximum area to obtain the greatest benefit, environmental damage is unlikely.

Interpretations developed for waste management may include ratings for (1) manure and food processing wastes; (2) municipal sewage sludge; (3) irrigation use of wastewater; or (4) treatment of wastewater by the slow rate process, overland flow process, or rapid infiltration process. If available, these should be located in this subsection.

Soil properties are important considerations in areas where soils are used as sites for the treatment and disposal of organic waste and wastewater. Selection of soils with properties that favor waste management can help to prevent environmental damage.

The Ag-Waste tables show the degree and kind of soil limitations affecting the treatment of agricultural waste, including municipal and food-processing wastewater and effluent from lagoons or storage ponds. Municipal wastewater is the waste stream from a municipality. It contains domestic waste and may contain industrial waste. It may have received primary or secondary treatment. It is rarely untreated sewage. Food-processing wastewater results from the preparation of fruits, vegetables, milk, cheese, and meats for public consumption. In places it is high in content of sodium and chloride. In the context of these tables, the effluent in lagoons and storage ponds is from facilities used to treat or store food-processing wastewater or domestic or animal waste. Domestic and food-processing wastewater is very dilute, and the effluent from the facilities that treat or store it commonly is very low in content of carbonaceous and nitrogenous material; the content of nitrogen commonly ranges from 10 to 30 milligrams per liter. The wastewater from animal waste treatment lagoons or storage ponds, however, has much higher concentrations of these materials, mainly because the manure has not been diluted as much as the domestic waste. The content of nitrogen in this wastewater generally ranges from 50 to 2,000 milligrams per liter. When wastewater is applied, checks should be made to ensure that nitrogen, phosphorus, heavy metals, and salts are not added in excessive amounts.

The ratings in the tables are for waste management systems that not only dispose of and treat organic waste or wastewater but also are beneficial to crops (application of manure and food-processing waste, application of sewage sludge, and disposal of wastewater by irrigation) and for waste management systems that are designed only for the purpose of wastewater disposal and treatment (overland flow of wastewater, rapid infiltration of wastewater, and slow rate treatment of wastewater).

The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect agricultural waste management. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are generally favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Application of manure and food-processing waste not only disposes of waste material but also can improve crop production by increasing the supply of nutrients in the soils where the material is applied. Manure is the excrement of livestock and poultry, and food-processing waste is damaged fruit and vegetables and the peelings, stems, leaves, pits, and soil particles removed in food preparation. The manure and food-processing waste are either solid, slurry, or liquid. Their nitrogen content varies. A high content of nitrogen limits the application rate. Toxic or otherwise dangerous wastes, such as those mixed with the lye used in food processing, are not considered in the ratings.

The ratings are based on the soil properties that affect absorption, plant growth, microbial activity, erodibility, the rate at which the waste is applied, and the method by which the waste is applied. The properties that affect absorption include permeability, depth to a water table, ponding, the sodium adsorption ratio, depth to bedrock or a cemented pan, and available water capacity. The properties that affect plant growth and microbial activity include reaction, the sodium adsorption ratio, salinity, and bulk density. The wind erodibility group, the soil erodibility factor K, and slope are considered in estimating the likelihood that wind erosion or water erosion will transport the waste material from the application site. Stones, cobbles, a water table, ponding, and flooding can hinder the application of waste. Permanently frozen soils are unsuitable for waste treatment.

Application of sewage sludge not only disposes of waste material but also can improve crop production by increasing the supply of nutrients in the soils where the material is applied. In the context of this table, sewage sludge is the residual product of the treatment of municipal sewage. The solid component consists mainly of cell mass, primarily bacteria cells that developed during secondary treatment and have incorporated soluble organics into their own bodies. The sludge has small amounts of sand, silt, and other solid debris. The content of nitrogen varies. Some sludge has constituents that are toxic to plants or hazardous to the food chain, such as heavy metals and exotic organic compounds, and should be analyzed chemically prior to use.

AGRICULTURAL WASTE MANAGEMENT
Boyd County, Nebraska

The content of water in the sludge ranges from about 98 percent to less than 40 percent. The sludge is considered liquid if it is more than about 90 percent water, slurry if it is about 50 to 90 percent water, and solid if it is less than about 50 percent water.

The ratings in the table are based on the soil properties that affect absorption, plant growth, microbial activity, erodibility, the rate at which the sludge is applied, and the method by which the sludge is applied. The properties that affect absorption, plant growth, and microbial activity include permeability, depth to a water table, ponding, the sodium adsorption ratio, depth to bedrock or a cemented pan, available water capacity, reaction, salinity, and bulk density. The wind erodibility group, the soil erodibility factor K, and slope are considered in estimating the likelihood that wind erosion or water erosion will transport the waste material from the application site. Stones, cobbles, a water table, ponding, and flooding can hinder the application of sludge. Permanently frozen soils are unsuitable for waste treatment.

Disposal of wastewater by irrigation not only disposes of municipal wastewater and wastewater from food-processing plants, lagoons, and storage ponds but also can improve crop production by increasing the amount of water available to crops. The ratings in the table are based on the soil properties that affect the design, construction, management, and performance of the irrigation system. The properties that affect design and management include the sodium adsorption ratio, depth to a water table, ponding, available water capacity, permeability, slope, and flooding. The properties that affect construction include stones, cobbles, depth to bedrock or a cemented pan, depth to a water table, and ponding.

The properties that affect performance include depth to bedrock or a cemented pan, bulk density, the sodium adsorption ratio, salinity, reaction, and the cation-exchange capacity, which is used to estimate the capacity of a soil to adsorb heavy metals. Permanently frozen soils are not suitable for disposal of wastewater by irrigation.

See the National Soil Handbook, September 1992, Part 620, for criteria used in rating soils for sanitary facilities and waste management.

AGRICULTURAL WASTE MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Very limited Restricted permeability Depth to saturated zone Flooding Runoff limitation	1.00 1.00 0.60 0.40	Very limited Restricted permeability Depth to saturated zone Flooding	1.00 1.00 1.00	Very limited Restricted permeability Depth to saturated zone Flooding	1.00 1.00 0.60
AnC: Anselmo-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity Too steep for surface application	1.00 0.08
AnD: Anselmo-----	100	Very limited Filtering capacity Slope	1.00 0.04	Very limited Filtering capacity Slope	1.00 0.04	Very limited Too steep for surface application Filtering capacity Too steep for sprinkler application	1.00 1.00 0.22
AnF: Anselmo-----	100	Very limited Filtering capacity Slope	1.00 0.96	Very limited Filtering capacity Slope	1.00 0.96	Very limited Too steep for surface application Filtering capacity Too steep for sprinkler application	1.00 1.00 0.97
ArF: Anselmo-----	75	Very limited Filtering capacity Slope	1.00 0.96	Very limited Filtering capacity Slope	1.00 0.96	Very limited Too steep for surface application Filtering capacity Too steep for sprinkler application	1.00 1.00 0.97
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
Ba: Barney-----	100	Very limited Depth to saturated zone Flooding Depth to dense layer Filtering capacity Runoff limitation	1.00 1.00 1.00 1.00 0.40	Very limited Depth to saturated zone Flooding Filtering capacity Droughty	1.00 1.00 1.00 0.34	Very limited Depth to saturated zone Flooding Filtering capacity Droughty	1.00 1.00 1.00 0.34
Bd: Blake-----	100	Somewhat limited Flooding Depth to saturated zone Restricted permeability	0.60 0.43 0.43	Very limited Flooding Depth to saturated zone Restricted permeability	1.00 0.43 0.32	Somewhat limited Flooding Depth to saturated zone Restricted permeability	0.60 0.43 0.32
Be: Blendon-----	100	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00
BeC: Blendon-----	100	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00	Somewhat limited Too steep for surface application	0.08

AGRICULTURAL WASTE MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
BoD: Boyd-----	100	Very limited Restricted permeability Droughty	1.00 0.99	Very limited Restricted permeability Droughty	1.00 0.99	Filtering capacity	0.00
						Very limited Restricted permeability	1.00
						Too steep for surface application	1.00
						Droughty	0.99
BrG: Bristow-----	100	Depth to bedrock Runoff limitation Slope	0.42 0.40 0.04	Depth to bedrock Slope	0.42 0.04	Depth to bedrock	0.42
						Too steep for sprinkler application	0.22
						Very limited Droughty	1.00
						Restricted permeability	1.00
Bs: Brocksburg-----	100	Depth to bedrock Droughty	1.00 1.00	Depth to bedrock Slope	1.00 1.00	Depth to bedrock	1.00
						Too steep for surface application	1.00
						Too steep for sprinkler application	1.00
						Runoff limitation	0.40
Bt: Brocksburg-----	100	Very limited Filtering capacity Restricted permeability	1.00 0.43	Very limited Filtering capacity Restricted permeability	1.00 0.32	Very limited Filtering capacity	1.00
						Restricted permeability	0.32
Cb: Cass-----	100	Very limited Filtering capacity Flooding	1.00 0.60	Very limited Flooding Filtering capacity	1.00 1.00	Very limited Filtering capacity	1.00
						Flooding	0.60
CrE2: Crofton-----	100	Somewhat limited Slope	0.84	Somewhat limited Slope	0.84	Very limited Too steep for surface application	1.00
						Too steep for sprinkler application	0.89
DuB: Dunday-----	100	Very limited Filtering capacity Leaching limitation Droughty	1.00 0.45 0.10	Very limited Filtering capacity Droughty	1.00 0.10	Very limited Filtering capacity	1.00
						Droughty	0.10
DuC: Dunday-----	100	Very limited Filtering capacity Leaching limitation Droughty	1.00 0.45 0.10	Very limited Filtering capacity Droughty	1.00 0.10	Very limited Filtering capacity	1.00
						Too steep for surface application	0.31
DuD: Dunday-----	100	Droughty	0.10	Droughty	0.10	Droughty	0.10
						Very limited Too steep for surface application	1.00
						Filtering capacity	1.00
						Too steep for sprinkler application	0.22
DuD: Dunday-----	100	Very limited Filtering capacity Leaching limitation Droughty	1.00 0.45 0.10	Very limited Filtering capacity Droughty	1.00 0.10	Very limited Filtering capacity	1.00
						Too steep for sprinkler application	0.10
DuD: Dunday-----	100	Slope	0.04	Slope	0.04	Droughty	0.10
						Droughty	0.10

AGRICULTURAL WASTE MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
DxB: Dunday-----	100	Very limited Filtering capacity Leaching limitation Droughty	1.00 0.45 0.12	Very limited Filtering capacity Droughty	1.00 0.12	Very limited Filtering capacity Droughty	1.00 0.12
Et: Eltree-----	100	Very limited Low adsorption	1.00	Very limited Low adsorption	1.00	Very limited Low adsorption	1.00
Go: Grigston-----	100	Somewhat limited Flooding	0.60	Very limited Flooding	1.00	Somewhat limited Flooding	0.60
GrB: Aowa-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Ha: Hall-----	100	Somewhat limited Restricted permeability	0.43	Somewhat limited Restricted permeability	0.32	Somewhat limited Restricted permeability	0.32
He: Haynie-----	100	Somewhat limited Flooding	0.60	Very limited Flooding	1.00	Somewhat limited Flooding	0.60
IfD: Inavale-----	100	Very limited Filtering capacity Leaching limitation Droughty	1.00 0.45 0.26	Very limited Filtering capacity Flooding Droughty	1.00 0.40 0.26	Very limited Filtering capacity Too steep for surface application Droughty Too steep for sprinkler application	1.00 0.91 0.26 0.02
IgB: Inavale-----	100	Very limited Flooding Filtering capacity Leaching limitation Droughty	1.00 1.00 0.45 0.26	Very limited Flooding Filtering capacity Droughty	1.00 1.00 0.26	Very limited Flooding Filtering capacity Droughty	1.00 1.00 0.26
IhB: Inavale-----	100	Very limited Filtering capacity Flooding Leaching limitation Droughty	1.00 0.60 0.45 0.13	Very limited Flooding Filtering capacity Droughty	1.00 1.00 0.13	Very limited Filtering capacity Flooding Droughty	1.00 0.60 0.13
In: Inavale-----	100	Very limited Filtering capacity Flooding Leaching limitation	1.00 0.60 0.45	Very limited Flooding Filtering capacity	1.00 1.00	Very limited Filtering capacity Flooding	1.00 0.60
Jn: Jansen-----	100	Very limited Filtering capacity Restricted permeability Too acid	1.00 0.43 0.02	Very limited Filtering capacity Restricted permeability Too acid	1.00 0.32 0.07	Very limited Filtering capacity Restricted permeability Too acid	1.00 0.32 0.07
JnC: Jansen-----	100	Very limited Filtering capacity Restricted permeability Too acid	1.00 0.43 0.02	Very limited Filtering capacity Restricted permeability Too acid	1.00 0.32 0.07	Very limited Filtering capacity Restricted permeability Too steep for surface application Too acid	1.00 0.32 0.08 0.07
JnD: Jansen-----	100	Very limited		Very limited		Very limited	

AGRICULTURAL WASTE MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food-processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
LaD: Labu-----	100	Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Restricted permeability	0.43	Restricted permeability	0.32	Too steep for surface application	1.00
		Slope	0.04	Too acid	0.07	Restricted permeability	0.32
		Too acid	0.02	Slope	0.04	Too steep for sprinkler application	0.22
						Too acid	0.07
		Very limited		Very limited		Very limited	
		Restricted permeability	1.00	Restricted permeability	1.00	Restricted permeability	1.00
		Droughty	0.97	Droughty	0.97	Too steep for surface application	1.00
		Depth to bedrock	0.42	Depth to bedrock	0.42	Droughty	0.97
		Runoff limitation	0.40	Slope	0.04	Depth to bedrock	0.42
LcF: Labu-----	65	Slope	0.04			Too steep for sprinkler application	0.22
		Very limited		Very limited			
		Restricted permeability	1.00	Restricted permeability	1.00	Very limited	
		Slope	1.00	Slope	1.00	Restricted permeability	1.00
						Too steep for surface application	1.00
		Droughty	0.97	Droughty	0.97	Too steep for sprinkler application	1.00
		Depth to bedrock	0.42	Depth to bedrock	0.42	Droughty	0.97
		Runoff limitation	0.40			Depth to bedrock	0.42
		Very limited		Very limited		Very limited	
		Depth to bedrock	1.00	Droughty	1.00	Droughty	1.00
Sansarc-----	35	Droughty	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Slope	1.00	Slope	1.00	Too steep for surface application	1.00
		Runoff limitation	0.40			Too steep for sprinkler application	1.00
		Very limited		Very limited		Very limited	
		Depth to bedrock	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Droughty	1.00	Droughty	1.00	Droughty	1.00
		Slope	1.00	Slope	1.00	Too steep for surface application	1.00
						Too steep for sprinkler application	1.00
Le: Leshara-----	100	Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Flooding	1.00	Depth to saturated zone	1.00
LsC: Lynch-----	100	Flooding	0.60	Depth to saturated zone	1.00	Flooding	0.60
		Very limited		Very limited		Very limited	
LsD: Lynch-----	100	Restricted permeability	1.00	Restricted permeability	1.00	Restricted permeability	1.00
		Droughty	0.92	Droughty	0.92	Droughty	0.92
LyD: Lynch-----	65	Depth to bedrock	0.42	Depth to bedrock	0.42	Depth to bedrock	0.42
		Runoff limitation	0.40			Too steep for surface application	0.08
		Slope	0.04			Too steep for sprinkler application	0.22

AGRICULTURAL WASTE MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food-processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Bristow-----	35	Depth to bedrock	0.42	Depth to bedrock	0.42	Droughty	0.92
		Runoff limitation	0.40	Slope	0.04	Depth to bedrock	0.42
		Slope	0.04			Too steep for sprinkler application	0.22
		Very limited Restricted permeability	1.00	Very limited Droughty	1.00	Very limited Droughty	1.00
		Depth to bedrock	1.00	Restricted permeability	1.00	Restricted permeability	1.00
LyF: Lynch-----	55	Droughty	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Runoff limitation	0.40	Slope	0.04	Too steep for surface application	1.00
		Slope	0.04			Too steep for sprinkler application	0.22
		Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Slope	1.00	Slope	1.00	Too steep for surface application	1.00
Bristow-----	45	Droughty	0.92	Droughty	0.92	Too steep for sprinkler application	1.00
		Depth to bedrock	0.42	Depth to bedrock	0.42	Droughty	0.92
		Runoff limitation	0.40			Depth to bedrock	0.42
		Very limited Restricted permeability	1.00	Very limited Droughty	1.00	Very limited Droughty	1.00
		Depth to bedrock	1.00	Restricted permeability	1.00	Restricted permeability	1.00
MaG: Mariaville-----	55	Droughty	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Slope	1.00	Slope	1.00	Too steep for surface application	1.00
		Droughty	1.00	Droughty	1.00	Too steep for sprinkler application	1.00
		Runoff limitation	0.40			Droughty	1.00
		Very limited Slope	1.00	Very limited Slope	1.00	Too steep for surface application	1.00
Paka-----	45	Restricted permeability	0.43	Restricted permeability	0.32	Too steep for sprinkler application	1.00
						Restricted permeability	0.32
MeE: Meadin-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Droughty	0.89	Droughty	0.89	Too steep for surface application	1.00
		Leaching limitation	0.45	Slope	0.16	Too steep for surface application	0.89
		Slope	0.16	Too acid	0.07	Droughty	0.89
		Too acid	0.02			Too steep for sprinkler application	0.39
NoC: Nora-----	100	Not limited		Not limited		Too acid	0.07
						Somewhat limited Too steep for surface application	0.08

AGRICULTURAL WASTE MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
NoD: Nora-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Too steep for surface application Too steep for sprinkler application	1.00 0.22
Oa: Onawa-----	100	Very limited Restricted permeability Flooding Depth to saturated zone Runoff limitation	1.00 0.60 0.43 0.40	Very limited Restricted permeability Flooding Depth to saturated zone	1.00 1.00 0.43	Very limited Restricted permeability Flooding Depth to saturated zone	1.00 0.60 0.43
Oe: O'Neill-----	100	Very limited Filtering capacity Depth to dense layer Droughty Too acid	1.00 1.00 0.12 0.11	Very limited Filtering capacity Too acid Droughty	1.00 0.42 0.12	Very limited Filtering capacity Too acid Droughty	1.00 0.42 0.12
OeC: O'Neill-----	100	Very limited Filtering capacity Depth to dense layer Droughty Too acid	1.00 1.00 0.12 0.11	Very limited Filtering capacity Too acid Droughty	1.00 0.42 0.12	Very limited Filtering capacity Too acid Droughty Too steep for surface application	1.00 0.42 0.12 0.08
OfD: O'Neill-----	55	Very limited Filtering capacity Depth to dense layer Droughty Too acid	1.00 1.00 0.12 0.11	Very limited Filtering capacity Too acid Droughty	1.00 0.42 0.12	Very limited Filtering capacity Too steep for surface application Too acid Droughty Too steep for sprinkler application	1.00 0.66 0.42 0.12 0.00
Meadin-----	45	Very limited Filtering capacity Droughty Leaching limitation Too acid	1.00 0.89 0.45 0.02	Very limited Filtering capacity Droughty Too acid	1.00 0.89 0.07	Very limited Filtering capacity Droughty Too steep for surface application Too acid Too steep for sprinkler application	1.00 0.89 0.66 0.07 0.00
On: Onita-----	100	Very limited Restricted permeability Very limited Slope Low adsorption	1.00 1.00 1.00	Very limited Restricted permeability Very limited Low adsorption Slope	1.00 1.00 1.00	Very limited Restricted permeability Very limited Low adsorption Too steep for surface application Too steep for sprinkler application	1.00 1.00 1.00
Poorly Drained-----							
Or: Ord-----	100	Very limited Filtering capacity Depth to saturated zone Flooding	1.00 0.95 0.60	Very limited Flooding Filtering capacity Depth to saturated zone	1.00 1.00 0.95	Very limited Filtering capacity Depth to saturated zone Flooding	1.00 0.95 0.60

AGRICULTURAL WASTE MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
PaC: Paka-----	100	Somewhat limited Restricted permeability Filtering capacity	0.43 0.00	Somewhat limited Restricted permeability Filtering capacity	0.32 0.00	Somewhat limited Restricted permeability Too steep for surface application Filtering capacity	0.32 0.08 0.00
Ph: Paka-----	100	Somewhat limited Restricted permeability	0.43	Somewhat limited Restricted permeability	0.32	Somewhat limited Restricted permeability	0.32
PhC: Paka-----	100	Somewhat limited Restricted permeability	0.43	Somewhat limited Restricted permeability	0.32	Somewhat limited Restricted permeability Too steep for surface application	0.32 0.08
PhD: Paka-----	100	Somewhat limited Restricted permeability Slope	0.43 0.04	Somewhat limited Restricted permeability Slope	0.32 0.04	Very limited Too steep for surface application Restricted permeability Too steep for sprinkler application	1.00 0.32 0.22
PoC: Promise-----	100	Very limited Restricted permeability Runoff limitation Sodium content	1.00 0.40 0.02	Very limited Restricted permeability Sodium content	1.00 0.02	Very limited Restricted permeability Too steep for surface application Sodium content	1.00 0.08 0.02
RaC: Ree-----	100	Somewhat limited Restricted permeability	0.43	Somewhat limited Restricted permeability	0.32	Somewhat limited Restricted permeability Too steep for surface application	0.32 0.08
RaD: Ree-----	100	Somewhat limited Restricted permeability Slope	0.43 0.04	Somewhat limited Restricted permeability Slope	0.32 0.04	Very limited Too steep for surface application Restricted permeability Too steep for sprinkler application	1.00 0.32 0.22
RaE: Ree-----	100	Somewhat limited Slope Restricted permeability	0.84 0.43	Somewhat limited Slope Restricted permeability	0.84 0.32	Very limited Too steep for surface application Too steep for sprinkler application Restricted permeability	1.00 0.89 0.32
ReC: Reliance-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability Too steep for surface application	1.00 0.08
ReD: Reliance-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00

AGRICULTURAL WASTE MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food-processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
RfC: Reliance-----	100	Slope	0.04	Slope	0.04	Too steep for surface application	1.00
						Too steep for sprinkler application	0.22
		Somewhat limited Restricted permeability	0.43	Somewhat limited Restricted permeability	0.32	Somewhat limited Restricted permeability	0.32
						Too steep for surface application	0.08
Rw: Riverwash-----	100	Very limited Depth to saturated zone	1.00	Very limited Droughty	1.00	Very limited Droughty	1.00
		Flooding	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Droughty Filtering capacity	1.00	Flooding Filtering capacity	1.00	Flooding Filtering capacity	1.00
		Runoff limitation	0.40				
SaG: Sansarc-----	100	Very limited Slope	1.00	Very limited Droughty	1.00	Very limited Droughty	1.00
		Depth to bedrock	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Droughty	1.00	Slope	1.00	Too steep for surface application	1.00
		Runoff limitation	0.40			Too steep for sprinkler application	1.00
Sc: Scott-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Runoff limitation	0.40	Too acid	0.42	Too acid	0.42
		Too acid	0.11				
Sm: Simeon-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Leaching limitation	0.45	Droughty	0.21	Droughty	0.21
		Droughty	0.21				
SuC: Simeon-----	55	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Leaching limitation	0.45	Droughty	0.10	Droughty	0.10
		Droughty	0.10			Too steep for surface application	0.00
Valentine-----	45	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Droughty	0.71	Droughty	0.71	Droughty	0.71
		Leaching limitation	0.45			Too steep for surface application	0.00
SvF2: Simeon-----	70	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Too steep for surface application	1.00
						Too steep for sprinkler application	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Too steep for sprinkler application	1.00
		Leaching limitation	0.45	Droughty	0.10	Filtering capacity	1.00
Valentine-----	30	Droughty	0.10			Droughty	0.10
		Very limited		Very limited		Very limited	

AGRICULTURAL WASTE MANAGEMENT--Continued
Boyd County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
VaE: Valentine-----	100	Slope	1.00	Slope	1.00	Too steep for surface application	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Too steep for sprinkler application	1.00
		Droughty	0.78	Droughty	0.78	Filtering capacity	1.00
		Leaching limitation	0.45			Droughty	0.78
		Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Too steep for surface application	1.00
		Droughty	0.78	Droughty	0.78	Filtering capacity	1.00
		Slope	0.63	Slope	0.63	Droughty	0.78
		Leaching limitation	0.45			Too steep for sprinkler application	0.77
		Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Droughty	0.71	Droughty	0.71	Droughty	0.71
VbB: Valentine-----	100	Leaching limitation	0.45				
Ve: Verdel-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Runoff limitation	0.40				
WeC: Wewela-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Droughty	0.99	Droughty	0.99	Droughty	0.99
		Depth to bedrock	0.42	Depth to bedrock	0.42	Depth to bedrock	0.42
		Filtering capacity	0.00	Filtering capacity	0.00	Too steep for surface application	0.08
						Filtering capacity	0.00
zwa: Water-----	100	Not rated		Not rated		Not rated	

In this section, hydric soils are defined and described and the hydric soils in the survey area are listed. The three essential characteristics of wetlands are hydrophytic vegetation, hydric soils, and wetland hydrology (Cowardin and others, 1979; U.S. Army Corps of Engineers, 1987; National Research Council, 1995; Tiner, 1985). Criteria for each of the characteristics must be met for areas to be identified as wetlands. Undrained hydric soils that have natural vegetation should support a dominant population of ecological wetland plant species. Hydric soils that have been converted to other uses should be capable of being restored to wetlands.

Hydric soils are defined by the National Technical Committee for Hydric Soils (NTCHS) as soils that formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part (Federal Register, 1994). These soils are either saturated or inundated long enough during the growing season to support the growth and reproduction of hydrophytic vegetation.

The NTCHS definition identifies general soil properties that are associated with wetness. In order to determine whether a specific soil is a hydric soil or nonhydric soil, however, more specific information, such as information about the depth and duration of the water table, is needed. Thus, criteria that identify those estimated soil properties unique to hydric soils have been established (Federal Register, 1995). These criteria are used to identify a phase of a soil series that normally is associated with wetlands. The criteria used are selected estimated soil properties that are described in "Soil Taxonomy" (USDA, 1999) and "Keys to Soil Taxonomy" (USDA, 1998) and in the "Soil Survey Manual" (USDA, 1993).

If soils are wet enough for a long enough period to be considered hydric, they should exhibit certain properties that can be easily observed in the field. These visible properties are indicators of hydric soils. The indicators used to make onsite determinations of hydric soils in this survey area are specified in "Field Indicators of Hydric Soils in the United States" (Hurt and others, 1996).

Hydric soils are identified by examining and describing the soil to a depth of about 20 inches. This depth may be greater if determination of an appropriate indicator so requires. It is always recommended that soils be excavated and described to the depth necessary for an understanding of the redoximorphic processes. Then, using the completed soil descriptions, soil scientists can compare the soil features required by each indicator and specify which indicators have been matched with the conditions observed in the soil. The soil can be identified as a hydric soil if at least one of the approved indicators is present.

Map units in the Hydric Soil Interpretations table meet the definition of hydric soils and, in addition, have at least one of the hydric soil indicators. This list can help in planning land uses; however, onsite investigation is recommended to determine the hydric soils on a specific site (National Research Council, 1995; Hurt and others, 1996).

Map units that are made up of hydric soils may have small areas, or inclusions, of nonhydric soils in the higher positions on the landform, and map units made up of nonhydric soils may have inclusions of hydric soils in the lower positions on the landform.

These map units, in general, do not meet the definition of hydric soils because they do not have one of the hydric soil indicators. A portion of these map units, however, may include hydric soils. Onsite investigation is recommended to determine whether hydric soils occur and the location of the included hydric soils.

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
Boyd County, Nebraska

PAGE 2 of 5

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
Ab: ALBATON SILTY CLAY, 0 TO 2 PERCENT SLOPES	ALBATON	Yes	depression, flood plain	2B3	YES	NO	NO
AnC: ANSELMO FINE SANDY LOAM, 2 TO 6 PERCENT SLOPES	ANSELMO	No	hill	---	---	---	---
AnD: ANSELMO FINE SANDY LOAM, 6 TO 11 PERCENT SLOPES	ANSELMO	No	hill	---	---	---	---
AnF: ANSELMO FINE SANDY LOAM, 11 TO 20 PERCENT SLOPES	ANSELMO	No	hill	---	---	---	---
ArF: ANSELMO-ROCK OUTCROP COMPLEX, 11 TO 20 PERCENT SLOPES	ANSELMO	No	hill	---	---	---	---
	ROCK OUTCROP	Unranked	hill	---	---	---	---
Ba: BARNEY SILT LOAM, 0 TO 2 PERCENT SLOPES	BARNEY	Yes	flood plain	2B3	YES	NO	NO
Bd: BLAKE SILTY CLAY LOAM, 0 TO 2 PERCENT SLOPES	BLAKE	No	flood plain	---	---	---	---
Be: BLENDON FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	ALBATON	Yes	swale	2B3	YES	NO	NO
	BLENDON	No	flat, swale	---	---	---	---
BeC: BLENDON FINE SANDY LOAM, 2 TO 6 PERCENT SLOPES	BLENDON	No	hill, swale, valley side	---	---	---	---
BoD: BOYD SILTY CLAY, 6 TO 11 PERCENT SLOPES	PONDED SOILS	Yes	depression	2B3,3	YES	NO	YES
BrG: BRISTOW SILTY CLAY, 20 TO 40 PERCENT SLOPES	BOYD	No	hill	---	---	---	---
Bs: BROCKSBURG FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	BRISTOW	No	hill, valley side	---	---	---	---
Bt: BROCKSBURG LOAM, 0 TO 2 PERCENT SLOPES	BROCKSBURG	No	flat	---	---	---	---
Cb: CASS FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	SCOTT	Yes	playa	2B3,3	YES	NO	YES
CrE2: CROFTON SILT LOAM, 11 TO 15 PERCENT SLOPES, ERODED	BROCKSBURG	No	flat	---	---	---	---
DuB: DUNDAY LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES	CASS	No	flood plain	---	---	---	---
DuC: DUNDAY LOAMY FINE SAND, 3 TO 6 PERCENT SLOPES	CROFTON	No	loess hill	---	---	---	---
DuD: DUNDAY LOAMY FINE SAND, 6 TO 11 PERCENT SLOPES	DUNDAY	No	hummock	---	---	---	---
DxB: DUNDAY LOAMY FINE SAND, LOAMY SUBSTRATUM, 0 TO 3 PERCENT SLOPES	DUNDAY	No	dune, valley side	---	---	---	---
Et: ELTREE SILT LOAM, 0 TO 2 PERCENT SLOPES	DUNDAY	No	dune, valley side	---	---	---	---
Go: GRIGSTON SILT LOAM, 0 TO 2 PERCENT SLOPES	ELTREE	No	flat	---	---	---	---
	SCOTT	Yes	playa	2B3,3	YES	NO	YES
	GRIGSTON	No	flood plain	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B3	YES	NO	NO

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
Boyd County, Nebraska

PAGE 3 of 5

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
GrB: GRIGSTON SILT LOAM, CHANNELED, 0 TO 3 PERCENT SLOPES	AOWA	No	flood plain	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B3	YES	NO	NO
Ha: HALL SILT LOAM, 0 TO 2 PERCENT SLOPES	HALL	No	stream terrace	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2A	YES	NO	NO
He: HAYNIE SILT LOAM, 0 TO 2 PERCENT SLOPES	HAYNIE	No	flood plain	---	---	---	---
Ifd: INAVALE FINE SAND, 3 TO 11 PERCENT SLOPES	INAVALE	No	flood plain	---	---	---	---
IgB: INAVALE FINE SAND, CHANNELED, 0 TO 3 PERCENT SLOPES	INAVALE	No	flood plain	---	---	---	---
	BARNEY	Yes	swale	2B3	YES	NO	NO
IhB: INAVALE LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES	INAVALE	No	flood plain	---	---	---	---
In: INAVALE FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	INAVALE	No	flood plain	---	---	---	---
Jn: JANSEN LOAM, 0 TO 2 PERCENT SLOPES	JANSEN	No	flat	---	---	---	---
JnC: JANSEN LOAM, 2 TO 6 PERCENT SLOPES	JANSEN	No	hill	---	---	---	---
JnD: JANSEN LOAM, 6 TO 11 PERCENT SLOPES	JANSEN	No	hill	---	---	---	---
LaD: LABU SILTY CLAY, 6 TO 11 PERCENT SLOPES	LABU	No	hill	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B3	YES	NO	NO
LcF: LABU-SANSARC SILTY CLAYS, 11 TO 30 PERCENT SLOPES	LABU	No	hill, valley side	---	---	---	---
	SANSARC	No	hill, valley side	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B3	YES	NO	NO
Le: LESHARA SILT LOAM, 0 TO 2 PERCENT SLOPES	LESHARA	No	flood plain	---	---	---	---
	POORLY DRAINED	Yes	swale	2B3	YES	NO	NO
LsC: LYNCH SILTY CLAY, 2 TO 6 PERCENT SLOPES	LYNCH	No	hill, valley side	---	---	---	---
LsD: LYNCH SILTY CLAY, 6 TO 11 PERCENT SLOPES	LYNCH	No	hill	---	---	---	---
LyD: LYNCH-BRISTOW SILTY CLAYS, 6 TO 11 PERCENT SLOPES	LYNCH	No	hill	---	---	---	---
	BRISTOW	No	hill	---	---	---	---
LyF: LYNCH-BRISTOW SILTY CLAYS, 11 TO 30 PERCENT SLOPES	LYNCH	No	valley side	---	---	---	---
	BRISTOW	No	valley side	---	---	---	---
MaG: MARIIVILLE-PAKA LOAMS, 15 TO 40 PERCENT SLOPES	MARIIVILLE	No	valley side	---	---	---	---
	PAKA WT AT 0-1 FOOT	No Yes	valley side swale	---	---	---	---
MeE: MEADIN SANDY LOAM, 3 TO 17 PERCENT SLOPES	MEADIN	No	hill, valley side	---	---	---	---

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
Boyd County, Nebraska

PAGE 4 of 5

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
NoC: NORA SILT LOAM, 2 TO 6 PERCENT SLOPES	NORA	No	loess hill	---	---	---	---
NoD: NORA SILT LOAM, 6 TO 11 PERCENT SLOPES	NORA	No	loess hill	---	---	---	---
Oa: ONAWA SILTY CLAY, 0 TO 2 PERCENT SLOPES	ONAWA	No	flood plain, swale	---	---	---	---
Oe: O'NEILL FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	ALBATON	Yes	swale	2B3	YES	NO	NO
OeC: O'NEILL FINE SANDY LOAM, 2 TO 6 PERCENT SLOPES	O'NEILL	No	flat	---	---	---	---
OfD: O'NEILL-MEADIN FINE SANDY LOAMS, 3 TO 9 PERCENT SLOPES	O'NEILL	No	hill	---	---	---	---
On: ONITA SILT LOAM, 0 TO 2 PERCENT SLOPES	MEADIN	No	hill	---	---	---	---
Or: ORD FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	ONITA	No	flat, swale	---	---	---	---
	POORLY DRAINED SCOTT	Yes	swale	2B3	YES	NO	NO
	ORD	Yes	playa	2B3,3	YES	NO	YES
	BARNEY	No	interdune, stream terrace	---	---	---	---
PaC: PAKA FINE SANDY LOAM, 2 TO 6 PERCENT SLOPES	Yes	No	swale	2B3	YES	NO	NO
Ph: PAKA LOAM, 0 TO 2 PERCENT SLOPES	PAKA	No	hill	---	---	---	---
PhC: PAKA LOAM, 2 TO 6 PERCENT SLOPES	SCOTT	Yes	playa	2B3,3	YES	NO	YES
PhD: PAKA LOAM, 6 TO 11 PERCENT SLOPES	PAKA	No	flat	---	---	---	---
PoC: PROMISE SILTY CLAY, 2 TO 6 PERCENT SLOPES	SCOTT	Yes	playa	2B3,3	YES	NO	YES
RaC: REE SILT LOAM, 2 TO 6 PERCENT SLOPES	PAKA	No	hill	---	---	---	---
RaD: REE SILT LOAM, 6 TO 11 PERCENT SLOPES	PROMISE	No	hill, valley side	---	---	---	---
RaE: REE SILT LOAM, 11 TO 15 PERCENT SLOPES	REE	No	hill	---	---	---	---
ReC: RELIANCE SILT LOAM, 2 TO 6 PERCENT SLOPES	REE	No	hill	---	---	---	---
ReD: RELIANCE SILT LOAM, 6 TO 11 PERCENT SLOPES	RELANCE	No	loess hill	---	---	---	---
RfC: RELIANCE SILTY CLAY LOAM, 2 TO 6 PERCENT SLOPES	SCOTT	Yes	playa	2B3,3	YES	NO	YES
Rw: RIVERWASH	RELANCE	No	loess hill	---	---	---	---
SaG: SANSARC SILTY CLAY, 20 TO 40 PERCENT SLOPES	SCOTT	Yes	playa	2B3,3	YES	NO	YES
Sc: SCOTT SILT LOAM, 0 TO 1 PERCENT SLOPES	RIVERWASH	Unranked	flood plain	---	---	---	---
Sm: SIMEON LOAMY SAND, 0 TO 2 PERCENT SLOPES	SANSARC	No	hill, valley side	---	---	---	---
	SCOTT	Yes	playa	2B3,3	YES	NO	YES
	SIMEON	No	flat	---	---	---	---

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
SuC: SIMEON-VALENTINE LOAMY SANDS, 0 TO 6 PERCENT SLOPES	SIMEON	No	swale	---	---	---	---
	VALENTINE	No	dune	---	---	---	---
SvF2: SIMEON-VALENTINE COMPLEX, 3 TO 30 PERCENT SLOPES, ERODED	SIMEON	No	hill, valley side	---	---	---	---
	VALENTINE	No	valley side, hill	---	---	---	---
VaE: VALENTINE FINE SAND, 6 TO 17 PERCENT SLOPES	VALENTINE	No	dune	---	---	---	---
VbB: VALENTINE LOAMY SAND, 0 TO 3 PERCENT SLOPES	VALENTINE	No	hummock	---	---	---	---
Ve: VERDEL SILTY CLAY, 0 TO 2 PERCENT SLOPES	VERDEL	No	stream terrace	---	---	---	---
WeC: WEWELA FINE SANDY LOAM, 2 TO 6 PERCENT SLOPES	WEWELA	No	hill	---	---	---	---
zwa: WATER > 40 ACRES	WATER	Unranked	---	---	---	---	---

FOOTNOTE: There may be small areas of included soils or miscellaneous areas that are significant to use and management of the soil; yet are too small to delineate on the soil map at the map's original scale. These may be designated as spot symbols and are defined in the published Soil Survey Report or the USDA-NRCS Technical Guide, Part II. Areas mapped as water or any map unit that contains one of the following conventional symbols is considered a hydric soil map unit: marshes or swamps; wet spots; depressions; streams, lakes and ponds.

1. All Histosols except Folists, or
2. Soils in Aquic suborders, great groups, or subgroups, Albolls suborder, Aquisalids, Pachic subgroups, or Cumulic subgroups that are:
 - a. Somewhat poorly drained with a water table equal to 0.0 foot (ft) from the surface during the growing season, or
 - b. poorly drained or very poorly drained and have either:
 - (1) water table equal to 0.0 ft during the growing season if textures are coarse sand, sand, or fine sand in all layers within 20 inches (in), or for other soils
 - (2) water table at less than or equal to 0.5 ft from the surface during the growing season if permeability is equal to or greater than 6.0 in/hour (h) in all layers within 20 in, or
 - (3) water table at less than or equal to 1.0 ft from the surface during the growing season if permeability is less than 6.0 in/h in any layer within 20 in, or
3. Soils that are frequently ponded for long duration or very long duration during the growing season, or
4. Soils that are frequently flooded for long duration or very long duration during the growing season.

HIGHLY ERODIBLE LANDS REPORT

Survey Area- BOYD COUNTY, NEBRASKA

Map Symbol	Soil Mapunit Name	HEL Classifications		
		C=30	R=100	wnd wat mu
Ab	ALBATON SILTY CLAY, 0 TO 2 PERCENT SLOPES	3	3	3
AnC	ANSELMO FINE SANDY LOAM, 2 TO 6 PERCENT SLOPES	3	3	3
AnD	ANSELMO FINE SANDY LOAM, 6 TO 11 PERCENT SLOPES	3	2	2
AnF	ANSELMO FINE SANDY LOAM, 11 TO 20 PERCENT SLOPES	3	2	2
ArF	ANSELMO-ROCK OUTCROP COMPLEX, 11 TO 20 PERCENT SLOPES	3	2	2
Ba	BARNEY SILT LOAM, 0 TO 2 PERCENT SLOPES	1	3	1
Bd	BLAKE SILTY CLAY LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
Be	BLENDON FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
BeC	BLENDON FINE SANDY LOAM, 2 TO 6 PERCENT SLOPES	3	3	3
BoD	BOYD SILTY CLAY, 6 TO 11 PERCENT SLOPES	3	2	2
BrG	BRISTOW SILTY CLAY, 20 TO 40 PERCENT SLOPES	1	1	1
Bs	BROCKSBURG FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
Bt	BROCKSBURG LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
Cb	CASS FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
CrE2	CROFTON SILT LOAM, 11 TO 15 PERCENT SLOPES, ERODED	3	1	1
DuB	DUNDAY LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES	1	3	1
DuC	DUNDAY LOAMY FINE SAND, 3 TO 6 PERCENT SLOPES	1	3	1
DuD	DUNDAY LOAMY FINE SAND, 6 TO 11 PERCENT SLOPES	1	2	1
DxB	DUNDAY LOAMY FINE SAND, LOAMY SUBSTRATUM, 0 TO 3 PERCENT SLOPES	1	3	1
Et	ELTREE SILT LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
Go	GRIGSTON SILT LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
GrB	GRIGSTON SILT LOAM, CHANNELED, 0 TO 3 PERCENT SLOPES	3	3	3
Ha	HALL SILT LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
He	HAYNIE SILT LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
IfD	INAVALE FINE SAND, 3 TO 11 PERCENT SLOPES	1	2	1
IgB	INAVALE FINE SAND, CHANNELED, 0 TO 3 PERCENT SLOPES	1	3	1
IhB	INAVALE LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES	1	3	1
In	INAVALE FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
Jn	JANSEN LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
JnC	JANSEN LOAM, 2 TO 6 PERCENT SLOPES	3	3	3
JnD	JANSEN LOAM, 6 TO 11 PERCENT SLOPES	3	2	2
LaD	LABU SILTY CLAY, 6 TO 11 PERCENT SLOPES	3	2	2
LcF	LABU-SANSARC SILTY CLAYS, 11 TO 30 PERCENT SLOPES	2	2	2
Le	LESHARA SILT LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
LsC	LYNCH SILTY CLAY, 2 TO 6 PERCENT SLOPES	3	2	2
LsD	LYNCH SILTY CLAY, 6 TO 11 PERCENT SLOPES	3	2	2
LyD	LYNCH-BRISTOW SILTY CLAYS, 6 TO 11 PERCENT SLOPES	2	2	2
LyF	LYNCH-BRISTOW SILTY CLAYS, 11 TO 30 PERCENT SLOPES	2	1	1
MaG	MARIAVILLE-PAKA LOAMS, 15 TO 40 PERCENT SLOPES	3	1	1
MeE	MEADIN SANDY LOAM, 3 TO 17 PERCENT SLOPES	1	2	1
NoC	NORA SILT LOAM, 2 TO 6 PERCENT SLOPES	3	2	2
NoD	NORA SILT LOAM, 6 TO 11 PERCENT SLOPES	3	2	2
Oa	ONAWA SILTY CLAY, 0 TO 2 PERCENT SLOPES	3	3	3
Oe	O'NEILL FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
OeC	O'NEILL FINE SANDY LOAM, 2 TO 6 PERCENT SLOPES	3	3	3
OfD	O'NEILL-MEADIN FINE SANDY LOAMS, 3 TO 9 PERCENT SLOPES	2	2	2
On	ONITA SILT LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
Or	ORD FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
PaC	PAKA FINE SANDY LOAM, 2 TO 6 PERCENT SLOPES	3	3	3
Ph	PAKA LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
PhC	PAKA LOAM, 2 TO 6 PERCENT SLOPES	3	2	2
PhD	PAKA LOAM, 6 TO 11 PERCENT SLOPES	3	2	2
PoC	PROMISE SILTY CLAY, 2 TO 6 PERCENT SLOPES	3	2	2
RaC	REE SILT LOAM, 2 TO 6 PERCENT SLOPES	3	2	2
RaD	REE SILT LOAM, 6 TO 11 PERCENT SLOPES	3	2	2

RaE	REE SILT LOAM, 11 TO 15 PERCENT SLOPES	3	1	1
ReC	RELIANCE SILT LOAM, 2 TO 6 PERCENT SLOPES	3	2	2
ReD	RELIANCE SILT LOAM, 6 TO 11 PERCENT SLOPES	3	2	2
RfC	RELIANCE SILTY CLAY LOAM, 2 TO 6 PERCENT SLOPES	3	2	2
Rw	RIVERWASH	2	2	2
SaG	SANSARC SILTY CLAY, 20 TO 40 PERCENT SLOPES	1	1	1
Sc	SCOTT SILT LOAM, 0 TO 1 PERCENT SLOPES	3	3	3
Sm	SIMEON LOAMY SAND, 0 TO 2 PERCENT SLOPES	1	3	1
SuC	SIMEON-VALENTINE LOAMY SANDS, 0 TO 6 PERCENT SLOPES	1	3	1
SvF2	SIMEON-VALENTINE COMPLEX, 3 TO 30 PERCENT SLOPES, ERODED	1	2	1
VaE	VALENTINE FINE SAND, 6 TO 17 PERCENT SLOPES	1	2	1
VbB	VALENTINE LOAMY SAND, 0 TO 3 PERCENT SLOPES	1	3	1
Ve	VERDEL SILTY CLAY, 0 TO 2 PERCENT SLOPES	3	3	3
WeC	WEWELA FINE SANDY LOAM, 2 TO 6 PERCENT SLOPES	3	2	2